Iridium-Catalyzed Asymmetric Allylic Amination of Allylic
Alcohols Activated by Lewis Acids

Supporting Information

Yasuhiro Yamashita, Apsara Gopalarathnam and John F. Hartwig

*Department of Chemistry, Yale University, P.O. Box 208107, New Haven,
Connecticut 06520-8107 and Department of Chemistry, University of Illinois
Urbana-Champaign, 600 South Mathews Avenue, Urbana, Illinois 61801*

Table of Contents

	Page
1. General	3
2. General procedure for the enantioselective allylic amination	
a. using a stoichiometric amount of niobium(V) ethoxide	4
3. General procedure for the enantioselective allylic amination	
a. using a catalytic amount of triphenylborane	4
4. Spectral characterization of products	5
5. Table S1. Asymmetric allylic substitutions of 1 with 2 in the presence of Ti alkoxides	13
6. Table S2. Screening of Lewis acid activator	14
7. Table S3. Screening of conditions with Ti(O ^{<i>i</i>} Pr) ₄ , Nb(OEt) ₅ and Ta(OEt) ₅ as activator	15
8. Figure S1. ¹ H NMR of 4-chloro- <i>N</i> -(1-(4-methoxyphenyl)allyl)benzenamine	16
9. Figure S2. ¹³ C NMR of 4-chloro- <i>N</i> -(1-(4-methoxyphenyl)allyl)benzenamine	17
10. References	18

General:

All manipulations were carried out under an inert atmosphere using a nitrogen-filled glovebox or standard Schlenk techniques. All glassware was oven dried immediately prior to use. THF was obtained as HPLC grade without inhibitors. Toluene was obtained as HPLC grade. *tert*-Butyl methyl ether (TBME), 1,2-dimethoxyethane (DME), and 1,4-dioxane were obtained as anhydrous grade packaged under nitrogen and used without further purification. THF and toluene were dried with a solvent purification system containing a 1 m column of activated alumina. All reagents were obtained from commercial sources and used without further purification unless otherwise noted. ^1H NMR spectra were obtained at 400 MHz and recorded relative to solvent (chloroform: 7.26 ppm). ^{13}C NMR spectra were obtained at 125 MHz, and chemical shifts were recorded relative to the solvent (Chloroform: 77.0 ppm). As proof of purity of previously unknown compounds, elemental analysis or copies of spectra are provided. $[\text{Ir}(\text{COD})\text{Cl}]_2$ ¹, *O,O'*-(biphenyl-2,2'-diyl)-*N*-cyclododecyl-*N*-(1*R*)-1-phenylethylphosphoamidite (**5a**)², and *O,O'*-(*R*)-(1,1'-binaphthyl-2,2'-diyl)-*N,N*-bis-[(1*R*)-1-phenylethyl]phosphoamidite (**5b**)³ were synthesized according to published procedures. *trans*-Cinnamyl alcohol and *trans*-2-hexenyl alcohol were purchased from commercial sources. *trans*-Cinnamyl alcohol was recrystallized from ether and pentane before use, and *trans*-2-hexenyl alcohol was distilled before use. *trans,trans*-2,4-Hexadien-1-ol was purchased from commercial sources and used without further purification. *trans-o*-Methoxycinnamyl alcohol, *trans*-4-methylpent-2-en-1-ol, and *trans*-3-(2-furyl)-2-propen-1-ol were synthesized from corresponding aldehydes by simple reduction using NaBH_4 in MeOH. *trans-p*-Methoxycinnamyl alcohol was synthesized from *trans-p*-methoxycinnamic acid according to the literature method.⁴ Propylamine (PrNH_2) was purchased from commercial sources, distilled from calcium hydride, and stored under nitrogen. Powdered molecular sieves were purchased from Aldrich Chemical Co. and used after activation by heating at 250 °C for 1 day under vacuum.

General procedure for the enantioselective allylic amination using a stoichiometric amount of niobium(V) ethoxide (Table 2). $[\text{Ir}(\text{COD})\text{Cl}]_2$ (0.0500 mmol) and phosphoramidite ligand **5** (0.100 mmol) were placed in a 20 mL vial with a magnetic stir bar, and dissolved in THF (0.50 mL) and PrNH_2 (0.50 mL). The mixture was heated at 50 °C for 1 h. After heating, all volatile materials were removed under reduced pressure. THF (1.00 mL) was then added to the vial to generate a stock solution of the iridium catalyst (0.100 M, 0.100 mmol Ir catalyst in 1.00 mL of THF).

To a 4 mL vial were added allylic alcohol **1** (1.00 mmol), powdered molecular sieves 4A (50 mg), amine **2** (1.50 mmol), THF (0.30 mL), a portion of the catalyst stock solution 0.20 mL (0.020 mmol) and $\text{Nb}(\text{OEt})_5$ (1.20 mmol). The mixture was stirred at 50 °C for 24 h unless otherwise noted. After cooling to room temperature, the reaction mixture was poured into a saturated sodium bicarbonate solution using methylene chloride. The mixture was stirred vigorously at room temperature for 1 h. The mixture was filtered through Celite[®], and the solid remained was washed well with methylene chloride. After the organic phase was separated, the aqueous phase was extracted with methylene chloride twice. The organic phases were combined and washed with brine, and were dried over anhydrous sodium sulfate. After filtration and concentration *in vacuo*, the crude mixture was analyzed to determine the ratio of **3** to **4** by ^1H NMR spectroscopy using CDCl_3 and one drop of D_2O . The addition of D_2O removed a proton resonance that overlapped with the allylic signals of **3** and **4** used to measure this ratio. The crude mixture was purified with silica gel column chromatography (hexanes-ether) to afford the desired branched product **3** (and linear product **4** in the cases noted in Table 2).

General procedure for the enantioselective allylic amination using a catalytic amount of triphenylborane (Table 3). $[\text{Ir}(\text{COD})\text{Cl}]_2$ (50.4 mg, 0.0750 mmol) and **5a** (75.2 mg, 0.150 mmol) were placed in a 20 mL vial with a magnetic stirrer bar and diluted with THF (0.90 mL) and PrNH_2 (0.90 mL). The mixture was heated at 50 °C for 1 h. After heating, all volatile materials were removed under reduced pressure. 1,4-Dioxane (1.50 mL) was then added to the vial to generate a stock solution of the iridium catalyst (0.1 M, 0.150 mmol Ir catalyst in 1.50 mL 1,4-dioxane).

To a 20 mL vial were added the aniline (**2**) (1.00 mmol), the cinnamyl alcohol (**1**) (1.50 mmol), powdered molecular sieves 4Å (300 mg), 1,4-dioxane (1.0 mL), 0.5 mL of the stock solution of the Ir catalyst (0.050 mmol), and a solution of triphenylborane (19.4 mg, 0.0801 mmol) in 1,4-dioxane (0.50 mL). The mixture was stirred at 50 °C for 24 h. After cooling to room temperature, the reaction mixture was filtered, and poured into saturated sodium bicarbonate. The organic phase was separated, and the aqueous phase was extracted with ether. The ether phases were combined and washed with brine and dried over anhydrous Na₂SO₄. After filtration and concentration *in vacuo*, the crude mixture was purified by silica gel column chromatography (hexane:ether = 50:1 or hexanes: CH₂Cl₂ = 9:1) to afford products **3**.

N-Phenyl-1-phenyl-2-propenylamine (Table 1):⁵ ¹H NMR (400 MHz, CDCl₃) 7.40-7.32 (m, 4H), 7.30-7.24 (m, 1H), 7.17-7.11 (m, 2H), 6.70 (t, 1H, *J* = 7.3 Hz), 6.60 (d, 2H, *J* = 7.9 Hz), 6.04 (ddd, 1H, *J* = 17.1, 10.2, 5.9 Hz), 5.28 (d, 1H, *J* = 17.1 Hz), 5.22 (d, 1H, *J* = 10.2 Hz), 4.93 (d, 1H, *J* = 5.8 Hz), 4.1 (br, 1H); ¹³C NMR (125 MHz, CDCl₃) 147.2, 141.9, 139.1, 129.1, 128.7, 127.4, 127.1, 117.6, 116.0, 113.5, 60.8; EI-MS [*m/z*] 209 (M⁺); HPLC (Daicel CHIRALCEL OJ-H; hexanes:2-propanol = 19:1; detection wavelength = 230 nm; flow rate = 0.6 mL/min) Tr = 30.8 min (major) and 35.7 min (minor).

N-Tolyl-1-phenyl-2-propenylamine (Table 2, entry 1):⁵ ¹H NMR (400 MHz, CDCl₃) 7.39-7.31 (m, 4H), 7.29-7.22 (m, 1H), 6.97-6.92 (m, 2H), 6.55-6.50 (m, 2H), 6.03 (ddd, 1H, *J* = 17.1, 10.2, 5.9 Hz), 5.27 (dt, 1H, *J* = 17.1, 1.3 Hz), 5.21 (dt, 1H, *J* = 10.2, 1.2 Hz), 4.90 (d, 1H, *J* = 5.9 Hz), 3.92 (br, 1H), 2.21 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) 144.9, 142.1, 139.3, 129.6, 128.7, 127.4, 127.1, 126.8, 115.9, 113.7, 61.1, 20.3; EI-MS [*m/z*] 223 (M⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.7:0.03, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 27.9 min (major) and 34.7 min (minor).

***N*-(*p*-Methoxyphenyl)-1-phenyl-2-propenylamine (Table 2, entry 2):**⁵ ¹H NMR (400 MHz, CDCl₃) 7.40-7.31 (m, 4H), 7.29-7.23 (m, 1H), 6.75-6.70 (m, 2H), 6.58-6.53 (m, 2H), 6.03 (ddd, 1H, *J* = 17.1, 10.2, 6.0 Hz), 5.26 (dt, 1H, *J* = 17.1, 1.2 Hz), 5.20 (dt, 1H, *J* = 10.2, 1.2 Hz), 4.85 (d, 1H, *J* = 6.0 Hz), 3.79 (br, 1H), 3.71 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) 152.2, 142.1, 141.4, 139.5, 128.7, 127.3, 127.1, 115.8, 114.9, 114.7, 61.8, 55.7; EI-MS [*m/z*] 239 (*M*⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.7:0.03, detection wavelength = 230 nm, flow rate = 1.0 mL/min) Tr = 19.5 min (major) and 26.4 min (minor).

***N*-(*o*-Methoxyphenyl)-1-phenyl-2-propenylamine (Table 2, entry 3):** ¹H NMR (400 MHz, CDCl₃) 7.41-7.31 (m, 4H), 7.28-7.24 (m, 1H), 6.79-6.74 (m, 2H), 6.65 (ddd, 1H, *J* = 8.3, 7.1, 1.5 Hz), 6.52-6.49 (m, 1H), 6.06 (ddd, 1H, *J* = 17.1, 10.2, 5.9 Hz), 5.26 (dt, 1H, *J* = 17.1, 1.3 Hz), 5.21 (dt, 1H, *J* = 10.2, 1.2 Hz), 4.92 (d, 1H, *J* = 5.8 Hz), 4.66 (br, 1H), 3.86 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) 146.8, 142.1, 139.3, 137.1, 128.7, 127.3, 127.1, 121.1, 116.7, 115.8, 111.2, 109.4, 60.7, 55.4; EI-MS [*m/z*] 239 (*M*⁺); [*α*]_D²⁰ = −4.3 (c 0.69, CHCl₃); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 9:1, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 10.5 min (minor) and 13.3 min (major); Anal. Calcd for C 80.30, H 7.16, N 5.85; Found: C 80.02, H 7.09, N 5.84.

***N*-(*m*-Methoxyphenyl)-1-phenyl-2-propenylamine (Table 2, entry 4):**⁵ ¹H NMR (400 MHz, CDCl₃) 7.39-7.31 (m, 4H), 7.29-7.24 (m, 1H), 7.04 (t, 1H, *J* = 8.1 Hz), 6.26 (dd, 1H, *J* = 8.1, 2.3 Hz), 6.22 (dd, 1H, *J* = 8.0, 2.1 Hz), 6.16 (t, 1H, *J* = 2.3 Hz), 6.03 (ddd, 1H, *J* = 17.1, 10.2, 5.8 Hz), 5.27 (dt, 1H, *J* = 17.2, 1.2 Hz), 5.22 (dt, 1H, *J* = 10.2, 1.2 Hz), 4.92 (d, 1H, *J* = 5.6 Hz), 4.06 (br, 1H), 3.72 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) 160.6, 148.6, 141.8, 139.0, 129.8, 128.7, 127.4, 127.1, 116.1, 106.6, 102.8, 99.6, 60.8, 55.0; EI-MS [*m/z*] 239 (*M*⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 19:1, detection

wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 29.3 min (minor) and 43.1 min (major).

***N*-(*p*-Iodophenyl)-1-phenyl-2-propenylamine (Table 2, entry 6):⁵**

¹H NMR (400 MHz, CDCl₃) 7.39-7.25 (m, 7H), 6.40-6.34 (m, 2H), 6.01 (ddd, 1H, *J* = 17.1, 10.2, 5.8 Hz), 5.25 (dt, 1H, *J* = 17.1, 1.2 Hz), 5.23 (dt, 1H, *J* = 10.2, 1.2 Hz), 4.88 (t, 1H, *J* = 4.7 Hz), 4.08 (br, 1H);

¹³C NMR (125 MHz, CDCl₃) 146.7, 141.2, 138.5, 137.6, 128.8, 127.6, 127.1, 116.3, 115.8, 78.3, 60.6; EI-MS [*m/z*] 335 (*M*⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.7:0.3, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 40.0 min (major) and 57.1 min (minor).

***N*-(*p*-Chlorophenyl)-1-phenyl-2-propenylamine (Table 2, entry 7):⁵**

¹H NMR (400 MHz, CDCl₃) 7.37-7.26 (m, 5H), 7.09-7.04 (m, 2H), 6.53-6.47 (m, 2H), 6.02 (ddd, 1H, *J* = 17.1, 10.2, 5.8 Hz), 5.26 (dt, 1H, *J* = 17.1, 1.1 Hz), 5.23 (dt, 1H, *J* = 10.2, 1.1 Hz), 4.88 (d, 1H, *J* = 5.8 Hz), 4.05 (br, 1H); ¹³C NMR (125 MHz, CDCl₃) 145.7,

141.4, 138.6, 128.9, 128.8, 127.6, 127.1, 122.2, 116.3, 114.6, 60.9; EI-MS [*m/z*] 243 (*M*⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.7:0.3, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 32.5 min (major) and 50.3 min (minor).

***N*-benzyl-1-phenyl-2-propenylamine (Table 2, entry 8):⁶**

¹H NMR (400 MHz, CDCl₃) 7.39-7.21 (m, 10H), 5.95 (ddd, 1H, *J* = 17.1, 10.2, 7.2 Hz), 5.23 (dt, 1H, *J* = 17.1, 1.2 Hz), 5.12 (d, 1H, *J* = 10.2 Hz), 4.22 (d, 1H, *J* = 7.2 Hz), 3.74 (d, 1H, *J* = 13.3 Hz), 3.72 (d, 1H, *J* = 13.3 Hz), 1.63 (br, 1H);

¹³C NMR (125 MHz, CDCl₃) 142.8, 141.0, 140.5, 128.5, 128.3, 128.1, 127.3, 127.2, 126.9, 115.1, 65.1, 51.3; EI-MS [*m/z*] 222 (*M*⁺-1), 223 (*M*⁺);

HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.7:0.3, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 15.6 min (major) and 19.4 min (minor).

***N*-(*p*-Methoxyphenylmethyl)-1-phenyl-2-propenylamine** (Table 2, entry 9):⁶ ¹H NMR (400 MHz, CDCl₃) 7.38-7.31 (m, 4H), 7.28-7.21 (m, 3H), 6.87-6.83 (m, 2H), 5.94 (ddd, 1H, *J* = 17.1, 10.2, 7.2 Hz), 5.22 (d, 1H, *J* = 17.1 Hz), 5.11 (d, 1H, *J* = 10.2 Hz), 4.21 (d, 1H, *J* = 7.2 Hz), 3.80 (s, 3H), 3.67 (d, 1H, *J* = 13.1 Hz), 3.65 (d, 1H, *J* = 13.1 Hz), 1.61 (br, 1H); ¹³C NMR (125 MHz, CDCl₃) 158.6, 142.9, 141.0, 132.6, 129.3, 128.5, 127.3, 127.2, 115.1, 113.8, 65.0, 55.3, 50.7; EI-MS [*m/z*] 253 (M⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.7:0.3, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 24.3 min (major) and 31.5 min (minor).

***N*-(1'-phenyl-2'-propenyl)morpholine** (Table 2, entry 10):⁶ ¹H NMR (400 MHz, CDCl₃) 7.36-7.22 (m, 5H), 5.90 (ddd, 1H, *J* = 17.8, 9.4, 8.3 Hz), 5.24 (d, 1H, *J* = 17.0 Hz), 5.10 (d, 1H, *J* = 10.0 Hz), 3.75-3.65 (m, 4H), 3.62 (d, 1H, *J* = 8.8 Hz), 2.55-2.45 (m, 2H), 2.38-2.28 (m, 2H), ¹³C NMR (125 MHz, CDCl₃) 141.6, 139.7, 128.6, 128.0, 127.2, 116.6, 75.5, 67.1, 52.0; EI-MS [*m/z*] 203 (M⁺); HPLC (Daicel CHIRALCEL OJ-H; hexanes:2-propanol:diethylamine = 99.7:0.3:0.01, detection wavelength = 210 nm, flow rate = 0.5 mL/min) Tr = 19.6 min (minor) and 24.6 min (major).

***N*-Phenyl-1-(*p*-methoxyphenyl)-2-propenylamine** (Table 2, entry 11):⁵ ¹H NMR (400 MHz, CDCl₃) 7.31-7.26 (m, 2H), 7.16-7.11 (m, 2H), 6.90-6.85 (m, 2H), 6.69 (t, 1H, *J* = 7.3 Hz), 6.60 (d, 2H, *J* = 7.7 Hz), 6.02 (ddd, 1H, *J* = 17.1, 10.2, 5.8 Hz), 5.26 (dt, 1H, *J* = 17.1, 1.3 Hz), 5.21 (dt, 1H, *J* = 10.2, 1.2 Hz), 4.88 (d, 1H, *J* = 5.8 Hz), 4.1 (br, 1H), 3.79 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) 158.9, 147.2, 139.2, 133.9, 129.1, 128.3, 117.6, 115.8, 114.1, 113.6, 60.2, 55.3; EI-MS [*m/z*] 239 (M⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 19:1, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 14.7 min (major) and 16.3 min (minor).

N-Phenyl-1-(*o*-methoxyphenyl)-2-propenylamine (Table 2, entry 12):⁵ ¹H NMR (400 MHz, CDCl₃) 7.31 (dd, 1H, *J* = 7.4, 1.6 Hz), 7.24 (ddd, 1H, *J* = 8.0, 7.6, 1.7 Hz), 7.15-7.08 (m, 2H), 6.94-6.87 (m, 2H), 6.67 (t, 1H, *J* = 7.3 Hz), 6.61 (d, 2H, *J* = 7.7 Hz), 6.07 (ddd, 1H, *J* = 17.2, 10.2, 5.4 Hz), 5.34 (d, 1H, *J* = 5.4 Hz), 5.21 (dt, 1H, *J* = 17.2, 1.4 Hz), 5.16 (dd, 1H, *J* = 10.2, 1.4 Hz), 4.4 (br, 1H), 3.86 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) 156.9, 147.3, 138.6, 129.7, 129.0, 128.4, 127.8, 120.8, 117.4, 115.1, 113.5, 110.8, 55.5, 54.7; EI-MS [*m/z*] 239 (M⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.95:0.05, detection wavelength = 230 nm, flow rate = 1.0 mL/min) Tr = 27.4 min (major) and 52.5 min (minor).

N-Phenyl-1-(2-furyl)-2-propenylamine (Table 2, entry 13):⁵ ¹H NMR (400 MHz, CDCl₃) 7.38 (dd, 1H, *J* = 1.7 Hz, 0.7 Hz), 7.20-7.14 (m, 2H), 6.74 (t, 1H, *J* = 7.4 Hz), 6.67 (d, 2H, *J* = 7.7 Hz), 6.33 (dd, 1H, *J* = 3.2, 1.8 Hz), 6.23 (d, 1H, *J* = 3.2 Hz), 6.05 (ddd, 1H, *J* = 17.1, 10.2, 5.7 Hz), 5.34 (dt, 1H, *J* = 17.1, 1.2 Hz), 5.28 (dt, 1H, *J* = 10.2, 1.2 Hz), 5.08 (d, 1H, *J* = 5.6 Hz), 4.3 (br, 1H); ¹³C NMR (125 MHz, CDCl₃) 154.3, 146.8, 142.1, 136.3, 129.1, 118.1, 116.9, 113.7, 110.3, 106.7, 54.5; EI-MS [*m/z*] 199 (M⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 19:1, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 12.1 min (minor) and 14.3 min (major).

3-Phenylamino-1-hexene (Table 2, entry 14):⁵ ¹H NMR (400 MHz, CDCl₃) 7.18-7.12 (m, 2H), 6.68 (t, 1H, *J* = 7.3 Hz), 6.61 (d, 2H, *J* = 7.8 Hz), 5.73 (ddd, 1H, *J* = 17.2, 10.3, 6.2 Hz), 5.20 (dt, 1H, *J* = 17.2, 1.3 Hz), 5.11 (dt, 1H, *J* = 10.3, 1.2 Hz), 3.7 (br, 1H), 3.81 (q, 1H, *J* = 6.4 Hz), 1.63-1.37 (m, 4H), 0.94 (t, 3H, *J* = 7.3 Hz); ¹³C NMR (125 MHz, CDCl₃) 147.7, 140.2, 129.1, 117.1, 114.9, 113.3, 55.7, 38.0, 19.1, 14.0; EI-MS [*m/z*] 175 (M⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.95:0.05, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 30.0 min (major) and 38.1 min (minor).

4-Methyl-3-phenylamino-1-pentene (Table 2, entry 15)⁵: ¹H NMR (400 MHz, CDCl₃) 7.18-7.12 (m, 2H), 6.66 (t, 1H, *J* = 7.3 Hz), 6.60 (d, 2H, *J* = 8.0 Hz), 5.72 (ddd, 1H, *J* = 17.2, 10.3, 6.3 Hz), 5.19 (d, 1H, *J* = 17.1 Hz), 5.16 (d, 1H, *J* = 10.2 Hz), 3.71 (br, 1H), 3.65 (t, 1H, *J* = 5.9 Hz), 1.87 (qqd, 1H, *J* = 6.8, 6.8, 5.7 Hz), 0.99 (d, 3H, *J* = 6.8 Hz), 0.96 (d, 3H, *J* = 6.8 Hz); ¹³C NMR (125 MHz, CDCl₃) 147.9, 137.9, 129.1, 117.1, 116.0, 113.4, 61.5, 32.5, 18.8, 18.5; EI-MS [*m/z*] 175 (*M*⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.95:0.05, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 22.0 min (major) and 24.0 min (minor).

3-Phenylamino-1,4-hexadiene (Table 2, entry 16)⁵: ¹H NMR (400 MHz, CDCl₃) 7.19-7.13 (m, 2H), 6.70 (t, 1H, *J* = 7.3 Hz), 6.63 (d, 2H, *J* = 7.7 Hz), 5.86 (ddd, 1H, *J* = 17.2, 10.3, 5.7 Hz), 5.71 (dq, 1H, *J* = 15.4, 6.4, 1.1 Hz), 5.49 (ddq, 1H, *J* = 15.3, 6.2, 1.5 Hz), 5.25 (dt, 1H, *J* = 17.2, 1.4 Hz), 5.16 (dt, 1H, *J* = 10.3, 1.3 Hz), 4.36 (t, 1H, *J* = 6.0 Hz), 3.93 (br, 1H), 1.71 (dt, 3H, *J* = 6.4, 1.2 Hz); ¹³C NMR (125 MHz, CDCl₃) 147.2, 138.8, 131.0, 129.0, 127.3, 117.4, 115.3, 113.6, 57.6, 17.8; EI-MS [*m/z*] 173 (*M*⁺); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 99.95:0.05, detection wavelength = 230 nm, flow rate = 0.5 mL/min) Tr = 32.9 min (major) and 36.1 min (minor).

***N*-(1-(4-methoxyphenyl)allyl)-4-methylbenzenamine (Table 3, entry 5)**: ¹H NMR (400 MHz, CDCl₃) 7.33 (d, 2H, *J* = 8.8 Hz), 6.99 (d, 1H, *J* = 5.2 Hz), 6.92 (dt, 2H, *J* = 6.8, 1.6 Hz), 6.57 (d, 2H, *J* = 8.4 Hz), 6.05 (ddd, 1H, *J* = 16.8, 10.0, 5.6 Hz), 5.30 (dd, 1H, *J* = 16.8, 1.2 Hz), 5.24 (dd, 1H, *J* = 10.0, 1.2 Hz), 3.92 (br s, 1H), 3.83 (s, 3H), 2.26 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) 158.7, 144.9, 139.1, 134.0, 129.5, 128.2, 126.6, 115.5, 113.9, 113.6, 60.4, 55.2, 20.3; EI-MS [*m/z*] 253 (*M*⁺); [α]_D²⁰ = +20.0 (c 1.5, CHCl₃); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 19:1, detection wavelength = 230 nm, flow rate = 1.0 mL/min) Tr =

25.1 min (major) and 37.2 min (major); Anal. Calcd for C 80.60, H 7.56, N 5.53; Found: C 80.22, H 7.71, N 5.64.

3-methoxy-N-(1-(4-methoxyphenyl)allyl)benzenamine (Table 3, entry 6) : ^1H NMR (400 MHz, CDCl_3) 7.28 (d, 2H, $J = 6.8$ Hz), 7.04 (t, 1H, $J = 8.0$ Hz), 6.88 (d, 2H, $J = 8.0$ Hz), 6.26 (dd, 1H, $J = 8.4, 2.4$ Hz), 6.22 (dd, 1H, $J = 8.0, 2.0$ Hz), 6.15 (t, 1H, $J = 2.4$ Hz), 6.03 (ddd, 1H, $J = 16.0, 10.0, 5.6$ Hz), 5.26 (dt, 1H, $J = 17.2, 1.2$ Hz), 5.20 (dt, 1H, $J = 10.4, 1.6$ Hz), 4.86 (br s, 1H), 4.01 (br s, 1H), 3.79 (s, 3H), 3.77 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) 160.5, 158.8, 148.6, 139.0, 133.8, 129.7, 128.2, 115.8, 114.0, 106.6, 102.5, 99.4, 60.1, 55.2, 55.0; EI-MS [m/z] 269 (M^+); $[\alpha]_{\text{D}}^{20} = +13.5$ (c 3.20, CHCl_3); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 19:1, detection wavelength = 230 nm, flow rate = 1.0 mL/min) Tr = 49.5 min (major) and 61.5 min (minor); Anal. Calcd for C 75.81, H 7.11, N 5.20; Found C 75.32, H 7.24, N 5.20.

4-chloro-N-(1-(4-methoxyphenyl)allyl)benzenamine (Table 3, entry 7) : ^1H NMR (400 MHz, CDCl_3) 7.26 (dt, 2H, $J = 8.4, 2.0$ Hz), 7.07 (dt, 2H, $J = 9.2, 2.4$ Hz), 6.88 (dt, 2H, $J = 8.8, 2.0$ Hz), 6.51 (dt, 2H, $J = 8.8, 2.0$ Hz), 5.99 (ddd, 1H, $J = 16.0, 10.0, 5.6$ Hz), 5.25 (dt, 1H, $J = 13.2, 1.6$ Hz), 5.22 (ddd, 1H, $J = 5.2, 1.2$ Hz), 4.83 (d, 1H, $J = 5.2$ Hz), 4.00 (br s, 1H), 3.80 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) 160.0, 145.7, 138.7, 133.4, 128.8, 128.2, 122.0, 115.9, 114.5, 114.1, 60.0, 55.2; EI-MS [m/z] 273 (M^+); $[\alpha]_{\text{D}}^{20} = 15.5$ (c 1.70, CHCl_3); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 19:1, detection wavelength = 230 nm, flow rate = 1.0 mL/min) Tr = 31.4 min (major) and 44.0 min (minor).

4-methyl-*N*-(1-*p*-tolylallyl)benzenamine (Table 3, entry 8):

^1H NMR (400 MHz, CDCl_3) 7.26 (d, 2H, $J = 8.0$ Hz), 7.15 (d, 2H, $J = 8.0$ Hz), 6.95 (d, 2H, $J = 8.0$ Hz), 6.52 (d, 2H, $J = 8.0$ Hz), 6.02 (ddd, 1H, $J = 16.0, 10.0, 8.0$ Hz), 5.27 (dt, 1H, $J = 16.8, 1.6$ Hz), 5.20 (dt, 1H, $J = 10.4, 1.6$ Hz), 4.87 (d, 1H, $J = 4.0$ Hz), 2.34 (s, 3H), 2.22 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) 144.9, 139.3, 139.0, 137.0, 129.5, 129.4, 127.0, 126.6, 115.6, 113.6, 60.8, 21.1, 20.3; EI-MS [m/z] 237 (M^+); $[\alpha]_{\text{D}}^{20} = 17.1$ (c 1.80, CHCl_3); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 19:1, detection wavelength = 230 nm, flow rate = 1.0 mL/min) Tr = 24.5 min (major) and 32.8 min (minor); Anal. Calcd for C 86.03, H 8.07, N 5.90; Found: C 85.66, H 8.31, N 5.97.

***N*-(1-(4-bromophenyl)allyl)-4-methylbenzenamine (Table 3, entry 9):**

^1H NMR (400 MHz, CDCl_3) 7.46 (dt, 2H, $J = 8.0, 2.0$ Hz), 7.26 (dt, 2H, $J = 8.4, 2.0$ Hz), 6.94 (d, 2H, $J = 8.0$ Hz), 6.48 (d, 2H, $J = 8.0$ Hz), 5.99 (ddd, 1H, $J = 17.2, 10.0, 6.0$ Hz), 5.24 (dt, 1H, $J = 10.0, 1.6$ Hz), 5.21 (dt, 1H, $J = 1.6, 1.6$ Hz), 4.86 (d, 1H, $J = 4.0$ Hz), 3.90 (br s, 1H), 2.21 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) 144.5, 141.0, 138.7, 131.7, 129.6, 128.8, 127.0, 121.1, 116.5, 113.6, 60.6, 20.3; EI-MS [m/z] 301 (M^+); $[\alpha]_{\text{D}}^{20} = 63.0$ (c 2.0, CHCl_3); HPLC (Daicel CHIRALCEL OD-H; hexanes:2-propanol = 19:1, detection wavelength = 230 nm, flow rate = 1.0 mL/min) Tr = 37.1 min (major) and 55.1 min (minor); Anal. Calcd for C 63.59, H 5.34, N 4.63; Found: C 63.20, H 5.37, N 4.59.

Table S1^a Asymmetric allylic substitutions of **1a** with **2a** in the presence of titanium alkoxides

$ \begin{array}{c} \text{Ph}-\text{CH}=\text{CH}-\text{CH}_2\text{OH} + \text{PhNH}_2 \\ \textbf{1a} \quad \quad \quad \textbf{2a} \\ (1.0 \text{ eq.}) \quad \quad (1.2 \text{ eq.}) \end{array} \xrightarrow[\text{Titanium alkoxide Additive, 50 } ^\circ\text{C, 24 h}]{\text{Chiral Ir catalyst (5a) (2 mol\%)}} \begin{array}{c} \text{HN}-\text{Ph} \\ \\ \text{Ph}-\text{CH}-\text{CH}=\text{CH}_2 \\ \textbf{3aa} \end{array} $						
Entry	Titanium alkoxide (eq.)	Solvent	Additive	Isolated yield of 3aa (%)	3aa/4aa^b	ee of 3aa (%)
1	Ti(O ⁱ Pr) ₄ (0.5)	THF	4Å MS	30	92/8	82
2 ^c	Ti(O ⁱ Pr) ₄ (0.5)	THF	4Å MS	41	76/24	57
3	Ti(O ⁱ Pr) ₄ (1.2)	THF	3Å MS	20	91/9	80
4	Ti(O ⁱ Pr) ₄ (1.2)	THF	4Å MS	16	89/11	81
5	Ti(O ⁱ Pr) ₄ (1.2)	THF	5Å MS	7	60/40	74
6	Ti(OEt) ₄ (1.2)	THF	4Å MS	16	94/6	91
7	Ti(O ⁱ Bu) ₄ (0.5)	THF	4Å MS	36	92/8	80
8	Ti(O ⁱ Bu) ₄ (1.2)	THF	4Å MS	30	94/6	89
9 ^d	Ti(O ⁱ Bu) ₄ (1.2)	THF	4Å MS	28	93/7	85
10 ^e	Ti(O ⁱ Bu) ₄ (1.2)	THF	4Å MS	30	95/5	88
11	Ti(O ⁱ Bu) ₄ (3.0)	THF	4Å MS	21	94/6	91
12 ^f	Ti(O ⁱ Bu) ₄ (1.2)	THF	4Å MS	35	92/8	85
13	Ti(O ⁱ Bu) ₄ (1.2)	THF	3Å MS	25	92/8	89
14 ^f	Ti(O ⁱ Bu) ₄ (1.2)	THF	3Å MS	33	93/7	87
15	Ti(O ⁱ Bu) ₄ (0.5)	THF/DMF (4/1)	4Å MS	32	94/6	75
16	Ti(O ⁱ Bu) ₄ (1.2)	THF/DMF (4/1)	4Å MS	26	92/8	84
17	Ti(O ⁱ Bu) ₄ (1.2)	CH ₃ CN	4Å MS	34	93/7	44
18	Ti(O ⁱ Bu) ₄ (1.2)	DMF	4Å MS	32	95/5	77
19	Ti(O ⁱ Bu) ₄ (1.2)	CH ₃ NO ₂	4Å MS	trace	—	—
20	Ti(O ⁱ Bu) ₄ (1.2)	diglyme	4Å MS	35	93/7	90
21 ^g	Ti(O ⁱ Bu) ₄ (1.2)	diglyme	4Å MS	n.p.	— ^h	—
22	Ti(O ⁱ Bu) ₄ (1.2)	toluene	4Å MS	32	90/10	87
23 ^g	Ti(O ⁱ Bu) ₄ (1.2)	toluene	4Å MS	trace	— ^h	—
24	Ti(O ⁱ Bu) ₄ (1.2)	DME	4Å MS	37	94/6	88
25 ⁱ	Ti(O ⁱ Bu) ₄ (1.2)	DME	4Å MS	<44	65/35	62

^a The reaction was performed with **1a** (1.0 mmol) and **2a** (1.2 mmol) in THF (0.5 mL) at 50 °C for 24 h in the presence of the Ir complex (2 mol%) prepared from [Ir(COD)Cl]₂ (0.010 mmol) and **5a** (0.020 mmol), titanium alkoxide and powdered 4Å molecular sieves (50 mg) unless otherwise noted. ^b Ratio of **3aa** and **4aa** determined by ¹H NMR analysis of the crude reaction mixture. ^c For 48 h. ^d 4Å MS (100 mg) was used. ^e 3.0 mmol of **2a** was used. ^f The Ir catalyst (4 mol%) was used. ^g At 110 °C. ^h Linear products (*N*-mono- and *N,N*-dicinnamylaniiline) were obtained. ⁱ At 80 °C. n.p.: No desired product was obtained.

Table S2^a Screening of Lewis acid activator

<div style="text-align: center;"> </div>					
Entry	Activator (eq.)	Solvent	Isolated yield of 3aa (%)	3aa/4aa^b	ee of 3aa (%)
1 ^c	BEt ₃ (1.2)	THF	n.p.	— ^d	—
2 ^{c,e}	BEt ₃ (1.2)	THF	n.p.	—	—
3	BPh ₃ (1.2)	THF	n.p.	Only linear products ^f	
4 ^g	B(O ⁱ Pr) ₃ (1.2)	DME	n.p.	—	—
5 ^{e,g}	B(O ⁱ Pr) ₃ (1.2)	DME	n.p.	—	—
6 ^g	B(O ^t Bu) ₃ (1.2)	DME	n.p.	—	—
7	B ₂ O ₃ (1.2)	THF	n.p.	—	—
8 ^e	B ₂ O ₃ (1.2)	THF	n.p.	—	—
9	MgSO ₄ (2.0)	DME	n.p.	— ^d	—
10	MgSO ₄ (2.0)	toluene	n.p.	— ^d	—
11	CuSO ₄ (2.0)	DME	n.p.	— ^d	—
12	CuSO ₄ (2.0)	toluene	n.p.	— ^d	—
13	Al(OEt) ₃ (1.2)	THF	37	97/3	84
14	Al(O ⁱ Pr) ₃ (1.2)	THF	25	76/24	18
15	Al(O ^t Bu) ₃ (1.2)	THF	10	39/61	0
16	Al(O ^t Bu) ₃ (1.0)	toluene	9	74/26	47
17 ^h	AlMe ₃ (1.2)	hexane:THF=1:1	61 ^f	79/21	1
18 ^{h,i}	AlMe ₃ (1.2)	hexane:THF=1:1	38 ^f	58/42	−2
19 ^j	AlMe ₃ (0.5)	hexane:THF=1:2	20 ^f	73/27	<7
20 ^k	AlMe ₃ (1.0)	hexane:toluene=1:1	35 ^f	57/43	2
21	Si(OEt) ₄ (1.2)	THF	n.p.	—	—
22	Mg(O ^t Bu) ₂ (1.2)	THF	2	35/65	40
23	Mg(OEt) ₂ (1.2)	THF	n.p.	— ^d	—
24 ^k	PhMgBr (1.2)	THF:Et ₂ O = 3:2	n.p.	—	—
25	Y ₅ O(O ⁱ Pr) ₁₃ (0.24)	THF	n.p.	— ^d	—
26	V(O)(O ⁱ Pr) ₃ (1.2)	THF	<5 ^f	8/92	−6
27	Zr(OEt) ₄ (1.2)	THF	<12	93/7	88
28	Zr(O ⁱ Pr) ₄ (1.2)	THF	3	50/50	55
29	Zr(O ^t Bu) ₄ (1.2)	THF	n.p.	—	—
30	Nb(OEt) ₅ (1.2)	THF	82	98/2	92
31	Ta(OEt) ₅ (1.2)	THF	62	96/4	81
32	Ta(O ^t Bu) ₅ (1.2)	THF	54	96/4	84

^a The reaction was performed with **1a** (1.0 mmol) and **2a** (1.2 mmol) in THF (0.5 mL) at 50 °C for 24 h in the presence of the Ir complex (2 mol%) prepared from [Ir(COD)Cl]₂ (0.010 mmol) and **5a** (0.020 mmol), Lewis acid activator and powdered 4Å molecular sieves (50 mg) unless otherwise noted. ^b Ratio of **3aa** and **4aa** determined by ¹H NMR analysis of the crude reaction mixture. ^c The amount of solvent was 1.4 mL. ^d A small amount of linear product (**4aa**) was obtained. ^e Without 4Å MS. ^f *N,N*-dicinnamylaniline was also obtained. ^g At 80 °C. ^h The amount of solvent was 1.2 mL. ⁱ At rt for 72 h. ^j The amount of solvent was 0.75 mL. ^k The amount of solvent was 1.0 mL. n.p.: No desired product was obtained.

Table S3. Screening of conditions for the allylations with Ti(O^{*i*}Pr)₄, Nb(OEt)₅ and Ta(OEt)₅ as activator

Entry	Activator (eq.)	Catalyst	Yield (%) ^b	3aa/4aa ^c	Ee (%)
1	Ti(O ^{<i>i</i>} Pr) ₄ (1.2)	Ir/ 5a	16	89/11	81
2	Ti(O ^{<i>t</i>} Bu) ₄ (1.2)	Ir/ 5a	30	94/6	89
3	Ta(OEt) ₅ (1.2)	Ir/ 5a	62	96/4	81
4	Nb(OEt) ₅ (1.2)	Ir/ 5a	82	98/2	92
5 ^d	Nb(OEt) ₅ (1.2)	Ir/ 5a	47	91/9	92
6	Nb(OEt) ₅ (0.5)	Ir/ 5a	64	84/16	63
7 ^e	Nb(OEt) ₅ (1.2)	Ir/ 5a	85	97/3	92
8 ^f	Nb(OEt) ₅ (1.2)	Ir/ 5b	64	98/2	94
9 ^{e,g}	Nb(OEt) ₅ (1.2)	Ir/ 5a	42	98/2	95

^a The reaction was performed with **1a** (1.0 mmol) and **2a** (1.2 mmol) in THF (0.5 mL) at 50 °C for 24 h in the presence of the Ir complex (2 mol%) prepared from [Ir(COD)Cl]₂ (0.010 mmol) and **5a** or **5b** (0.020 mmol), Lewis acid activator and powdered 4 Å molecular sieves (4 Å MS, 50 mg) unless otherwise noted. ^b Isolated yield of branched product **3aa**. ^c Ratio of **3aa** and **4aa** determined by ¹H NMR analysis of the crude reaction mixture. ^d Without 4 Å MS. ^e 1.5 mmol of **2a** was used. ^f Ligand **5b** was used. ^g The reaction was conducted at room temperature.

Figure S1. ¹H NMR of 4-chloro-*N*-(1-(4-methoxyphenyl)allyl)benzenamine.

Figure S2. ¹³C NMR of 4-chloro-*N*-(1-(4-methoxyphenyl)allyl)benzenamine.

References

- (1) Herde, J. L.; Lambert, J. C.; Senoff, C. V. *Inorg. Synth.* **1974**, *15*, 18.
- (2) Leitner, A.; Shekhar, S.; Pouy, M. J.; Hartwig, J. F. *J. Am. Chem. Soc.* **2005**, *127*, 15506.
- (3) Alexakis, A.; Rosset, S.; Allamand, J.; March, S.; Guillen, F.; Benhaim, C. *Synlett* **2001**, 1375.
- (4) Wu, Z. X.; Minhas, G. S.; Wen, D. Y.; Jiang, H. L.; Chen, K. X.; Zimniak, P.; Zheng, J. *J. Med. Chem.* **2004**, *47*, 3282.
- (5) Shu, C. T.; Leitner, A.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2004**, *43*, 4797.
- (6) Ohmura, T.; Hartwig, J. F. *J. Am. Chem. Soc.* **2002**, *124*, 15164.