Supplementary Information

“Iridium Catalyzed C-C Bond Forming Hydrogenation: Direct Regioselective Reductive Coupling of Alkyl-Substituted Alkynes to Activated Ketones”

Ming-Yu Ngai, Andriy Barchuk and Michael J. Krische*

University of Texas at Austin

Department of Chemistry and Biochemistry

Austin, TX 78712

USA

Table of Contents

I. Experimental Section	
General -----	S2
General Procedure for the Reductive Coupling -----	S2
Detailed Procedure for the Reductive Coupling and Spectral Data -----	S3 – S42

I. Experimental Section

General. All reactions were run under an atmosphere of argon, unless otherwise indicated. All the commercially available substrates (**1a-d**, **2a-e** and **2g**) were purified before used. Anhydrous solvents were transferred by an oven-dried syringe. Reaction vessels (13mm x 180mm test tube) were dried in oven for overnight and cooled under a stream of nitrogen. Dichloroethane (DCE) was distilled from calcium hydride and toluene was distilled from sodium/benzophenone. Ir(cod)₂BARF was used as received from Umicore. BIPHEP and DPPF were used as received from Strem Chemicals. Triphenylacetic acid was purchased from Aldrich and recrystallized from hot methanol before used. Different substituted phenylpropynes (**1e-h**) were synthesized according to Sonogashira Coupling.¹ Phenethylpruvate (**2f**) was prepared according to the literature procedure.² Analytical thin-layer chromatography (TLC) was carried out using 0.2-mm commercial silica gel plates (DC-Fertigplatten Kieselgel 60 F₂₅₄). Preparative column chromatography employing silica gel was performed according to the method of Still.³ Infrared spectra were recorded on a Perkin-Elmer 1600 spectrometer. High-resolution mass spectra (HRMS) were obtained on a Karatos MS9 and are reported as m/z (relative intensity). Accurate masses are reported for the molecular ion (M + 1, M or M-1) or a suitable fragment ion. Proton nuclear magnetic resonance (¹H-NMR) spectra were recorded with a Varian Gemini (400 MHz or 300MHz) spectrometer. Chemical shifts are reported in delta (δ) units, parts per million (ppm) downfield from trimethylsilane. Coupling constants are reported in Hertz (Hz). Carbon-13 nuclear magnetic resonance (¹³C-NMR) spectra were recorded with a Varian Gemini 400 (100 MHz) spectrometer. Chemical shifts are reported in delta (δ) units, ppm relative to the center of the triplet at 77.0 ppm for deuteriochloroform. ¹³C NMR spectra were routinely run with broadband decoupling.

General Procedure for the Coupling of Alkynes and α-Ketoesters.

To a solution of Ir(COD)₂BARF (2 or 5 mol%), DPPF (2 mol%) or BIPHEP (5 mol%) and triphenylacetic acid (2 mol% or 5 mol%) in Toluene (0.5M) or DCE (0.5M) at 60°C or 80°C was added α-ketoester (100 mol%) and alkyne substrates (300 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 60 °C or 80°C under 1 atm of hydrogen until complete consumption of α-ketoester at which point the reaction mixture was concentrated and the product was purified by silica gel chromatography.

¹ (a) Sonogashira, K.; Tohda, Y.; Hagihara, N. *Tetrahedron Lett.* **1975**, 4467. (b) Pschirer, N. G.; Bunz, U. H. F. *Tetrahedron Lett.* **1999**, 2481. (c) Carpita, A.; Lezzi, A.; Rossi, R.; Marchetti, F.; Merlino, S. *Tetrahedron* **1985**, *41*, 621.

² Wong, M.-K.; Chung, N.-W.; He, L.; Wang, X.-C.; Yan, Z.; Tang, Y.-C.; Yang, D. *J. Org. Chem.* **2003**, *68*, 6321.

³ Still, W.C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923.

(E)-ethyl 3-ethyl-2-hydroxy-2-phenylhex-3-enoate (3a).

To a solution of Ir(cod)₂BARF (5.1 mg, 0.004 mmol, 1.9 mol%), DPPF (2.3 mg, 0.0041 mmol, 1.9 mol%), and triphenyl acetic acid (1.3 mg, 0.0045 mmol, 2.1 mol%) in toluene (0.4 mL, 0.5 M) was added ethyl benzoylformate (**2a**) (37.5 mg, 0.21 mmol, 100 mol%) and hexyne (**1a**) (54.6 mg, 0.66 mmol, 316 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 60 °C under 1 atm of hydrogen gas for 1 hour. The title compound (**3a**) was purified by flash column chromatography (R_f = 0.32, 10 %EtOAc/hexane) to afford 51.2 mg as colorless oil (93 % yield).

¹H NMR (400 MHz, CDCl₃): 7.53-7.56 (m, 2H), 7.25-7.35 (m, 3H), 5.28 (t, *J* = 7.2 Hz, 1H), 4.22-4.32 (m, 2H), 3.77 (s, 1H), 2.10 (m, 4H), 1.28 (t, *J* = 7.2 Hz, 3H), 0.98 (t, *J* = 7.2 Hz, 3H), 0.97 (t, *J* = 7.2 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 174.74, 141.13, 140.37, 131.69, 127.67, 127.58, 127.31, 83.13, 62.42, 21.83, 21.27, 14.83, 14.00, 13.97

HRMS Calcd. for C₁₆H₂₂O₃ (M): 262.1569, Found: 262.1570

FTIR(neat): 3511, 2963, 2934, 2874, 2360, 2340, 1718, 1448, 1367, 1299, 1238, 1169, 1122, 1092, 1057, 1032, 1016 cm⁻¹

(E)-ethyl 3-ethyl-2-hydroxy-2-(4-nitrophenyl)hex-3-enoate (3b)

To a solution of Ir(cod)₂BARF (5.0 mg, 0.004 mmol, 2 mol%), DPPF (2.4 mg, 0.0043 mmol, 2.3 mol%), and triphenyl acetic acid (1.3 mg, 0.0045 mmol, 2.2 mol%) in toluene (0.4 mL, 0.5 M) was added ethyl 4-nitrobenzoylformate (**2b**) (98 %, 46.5 mg, 0.2 mmol, 100 mol%) and hexyne (**1a**) (51.1 mg, 0.62 mmol, 310 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 60 °C under 1 atm of hydrogen gas for 1 hour. The title compound (**3b**) was purified by flash column chromatography (R_f = 0.39, 20 %EtOAc/hexane) to afford 52.0 mg as colorless oil (83 % yield).

¹H NMR (400 MHz, CDCl₃): 8.19 (d, *J* = 8.8 Hz, 2H), 7.78 (d, *J* = 8.8 Hz, 2H), 5.25 (t, *J* = 7.2 Hz, 3H), 4.22-4.36 (m, 2H), 3.89 (s, 1H), 1.94-2.16 (m, 4H), 1.29 (t, *J* = 7.2 Hz, 3H), 0.99 (t, *J* = 7.2 Hz, 3H), 0.97 (t, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 173.45, 147.50, 147.43, 140.81, 132.22, 132.24, 128.52, 122.79, 82.99, 63.00, 21.63, 21.28, 14.88, 13.98, 13.89

HRMS Calcd. for C₁₆H₂₂NO₅ (M+1): 308.1498, Found: 308.1504

FTIR(neat): 3481, 2978, 2932, 2361, 2340, 1709, 1520, 1343, 1246, 1171, 1080, 1012 cm⁻¹

(E)-ethyl 3-ethyl-2-hydroxy-2-(4-methoxyphenyl)hex-3-enoate (3c).

3c

To a solution of Ir(cod)₂BARF (4.7 mg, 0.0037 mmol, 2.1 mol%), DPPF (2.4 mg, 0.0043 mmol, 2.4 mol%), and triphenyl acetic acid (1.4 mg, 0.0049 mmol, 2.7 mol%) in toluene (0.4 mL, 0.45 M) was added ethyl 4-methoxybenzoylformate (**2c**) (96 %, 39.8 mg, 0.18 mmol, 100 mol%) and hexyne (**1a**) (49.9 mg, 0.6 mmol, 337 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 60 °C under 1 atm of hydrogen gas for 2 hours. The title compound (**3c**) was purified by flash column chromatography (R_f = 0.21, 10 %EtOAc/hexane) to afford 50.0 mg as colorless oil (93 % yield).

¹H NMR (400 MHz, CDCl₃): 7.46 (d, *J* = 8.8 Hz, 2H), 6.86 (d, *J* = 8.8 Hz, 2H), 5.27 (t, *J* = 7.2 Hz, 1H), 4.21-4.27 (m, 2H), 3.81 (s, 3H), 3.74 (s, 1H), 2.04-2.14 (m, 4H), 1.28 (t, *J* = 7.2 Hz, 3H), 0.98 (t, *J* = 7.2 Hz, 3H), 0.97 (t, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 174.94, 158.97, 141.25, 132.43, 128.52, 112.98, 82.79, 62.34, 55.17, 55.15, 21.77, 21.26, 14.82, 14.02, 13.99

HRMS Calcd. for C₁₇H₂₅O₄ (M+1): 293.1753, Found: 293.1753

FTIR(neat): 3490, 2963, 2934, 2873, 2360, 2336, 1718, 1609, 1508, 1463, 1367, 1299, 1238, 1167, 1123, 1067, 1032, 955 cm⁻¹

(E)-methyl 2-(3-bromophenyl)-3-ethyl-2-hydroxyhex-3-enoate (3d).

To a solution of Ir(cod)₂BARF (4.8 mg, 0.0038 mmol, 1.9 mol%), DPPF (2.3 mg, 0.0041 mmol, 2.1 mol%), and triphenyl acetic acid (1.2 mg, 0.0042 mmol, 2.1 mol%) in toluene (0.4 mL, 0.5 M) was added methyl 3-bromobenzoylformate (**2d**) (48.8 mg, 0.2 mmol, 100 mol%) and hexyne (**1a**) (50.5 mg, 0.61 mmol, 305 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 60 °C under 1 atm of hydrogen gas for 1 hour. The title compound (**3d**) was purified by flash column chromatography (*R_f* = 0.26, 10 %EtOAc/hexane) to afford 60.0 mg as colorless oil (91 % yield).

¹H NMR (400 MHz, CDCl₃): 7.72 (t, *J* = 1.6 Hz, 1H), 7.65 (dt, *J* = 8.0, 1.6 Hz, 1H), 7.41 (ddd, *J* = 8.0, 1.6, 0.8 Hz, 1H), 7.20 (t, *J* = 8.0 Hz, 1H), 5.22 (t, *J* = 7.2 Hz, 1H), 3.81 (s, 3H), 3.75 (s, 1H), 1.98-2.15 (m, 4H), 0.99 (t, *J* = 7.2 Hz, 3H), 0.97 (t, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 174.67, 142.45, 140.86, 131.94, 130.81, 130.41, 129.23, 126.21, 122.06, 82.96, 53.30, 53.27, 21.69, 21.28, 14.85, 13.94

HRMS Calcd. for C₁₅H₁₈O₃Br (M-1): 325.0439, Found: 325.0437

FTIR(neat): 3052, 2964, 2944, 2875, 2360, 2342, 1728, 1582, 1566, 1471, 1436, 1419, 1246, 1171, 1070 cm⁻¹

(E)-ethyl 3-ethyl-2-hydroxy-2-(thiophen-2-yl)hex-3-enoate (3e).

To a solution of Ir(cod)₂BARF (4.8 mg, 0.0038 mmol, 1.8 mol%), DPPF (2.5 mg, 0.0045 mmol, 2.1 mol%), and triphenyl acetic acid (1.2 mg, 0.0042 mmol, 2 mol%) in toluene (0.4 mL, 0.5 M) was added ethyl thiophene-2-glyoxalate (**2e**) (38.5 mg, 0.21 mmol, 100 mol%) and hexyne (**1a**) (49.8 mg, 0.61 mmol, 290 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 60 °C under 1 atm of hydrogen gas for 1 hour. The title compound (**3e**) was purified by flash column chromatography (R_f = 0.26, 10 %EtOAc/hexane) to afford 46 mg as colorless oil (85 % yield).

¹H NMR (400 MHz, CDCl₃): 7.26 (m, 1H), 7.09 (dd, *J* = 3.6, 1.2 Hz, 1H), 6.98 (dd, *J* = 5.2, 3.6 Hz, 1H), 5.34 (t, *J* = 7.2 Hz, 1H), 4.32 (q, *J* = 7.2 Hz, 2H), 4.13 (s, 1H), 2.04-2.11 (m, 4H), 1.33 (t, *J* = 7.2 Hz, 3H), 0.99 (t, *J* = 7.2 Hz, 3H), 0.95 (t, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 173.88, 145.05, 140.64, 131.73, 126.55, 126.06, 125.29, 81.17, 62.81, 21.38, 21.21, 14.67, 14.00, 13.86

HRMS Calcd. for C₁₄H₂₁O₃S (M+1): 269.1211, Found: 269.1210

FTIR(neat): 3496, 2964, 2933, 2874, 2361, 2340, 1723, 1464, 1369, 1247, 1226, 1155, 1117, 1079, 1042, 1014 cm⁻¹

(E)-phenethyl 3-ethyl-2-hydroxy-2-methylhex-3-enoate (3f)

To a solution of Ir(cod)₂BARF (4.9 mg, 0.0039 mmol, 1.8 mol%), DPPF (2.4 mg, 0.0043 mmol, 2.1 mol%), and triphenyl acetic acid (1.4 mg, 0.0049 mmol, 2.3 mol%) in toluene (0.4 mL, 0.5 M) was added phenethyl pyruvate (**2f**) (40.3 mg, 0.21 mmol, 100 mol%) and hexyne (**1a**) (49.1 mg, 0.60 mmol, 285 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 60 °C under 1 atm of hydrogen gas for 8 hours. The title compound (**3f**) was purified by flash column chromatography (R_f = 0.21, 10 %EtOAc/hexane) to afford 47.2 mg as colorless oil (82 % yield).

¹H NMR (400 MHz, CDCl₃): 7.19-7.31 (m, 5H), 5.51 (t, *J* = 7.2 Hz, 1H), 4.32-4.40 (m, 2H), 3.28 (s, 1H), 2.95 (t, *J* = 7.2 Hz, 2H), 2.00-2.09 (m, 4H), 1.47 (s, 3H), 0.98 (t, *J* = 7.2 Hz, 3H), 0.91 (t, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 176.38, 140.73, 137.36, 128.85, 128.75, 128.44, 126.62, 77.15, 66.24, 34.87, 24.27, 21.05, 20.34, 14.54, 14.07

HRMS Calcd. for C₁₇H₂₄O₃ (M): 276.1725, Found: 276.1721

FTIR(neat): 3513, 3028, 2963, 2934, 2874, 2361, 2342, 1722, 1498, 1455, 1375, 1240, 1197, 1148, 1120, 1082, 959, 862 cm⁻¹

(E)-4-deuterio-ethyl 3-ethyl-2-hydroxy-2-phenylhex-3-enoate (*deuterio-3a*).

deuterio-3a

To a solution of Ir(cod)₂BARF (5.0 mg, 0.004 mmol, 1.7 mol%), DPPF (2.3 mg, 0.0041 mmol, 1.8 mol%), and triphenyl acetic acid (1.3 mg, 0.0045 mmol, 2.1 mol%) in toluene (0.4 mL, 0.6 M) was added ethyl benzoylformate (**2a**) (41.4 mg, 0.23 mmol, 100 mol%) and hexyne (**1a**) (53.8 mg, 0.65 mmol, 285 mol%). The system was purged with argon gas followed by deuterium gas (99.6% D₂). The reaction was allowed to stir at 60 °C under 1 atm of deuterium gas for 1.5 hour. The title compound (***deuterio-3a***) was purified by flash column chromatography (R_f = 0.32, 10 %EtOAc/hexane) to afford 55 mg as colorless oil (94 % yield).

¹H NMR (400 MHz, CDCl₃): 7.54-7.56 (m, 2H), 7.25-7.34 (m, 3H), 4.20-4.34 (m, 2H), 3.78 (s, 1H), 2.05-2.13 (m, 4H), 1.28 (t, *J* = 7.2 Hz, 3H), 0.98 (t, *J* = 7.2 Hz, 3H), 0.97 (t, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 174.75, 141.03, 140.37, 127.67, 127.58, 127.31, 83.10, 62.42, 21.80, 21.17, 14.83, 14.01, 13.95

HRMS Calcd. for C₁₆H₂₁DO₃ (M): 263.1632, Found: 263.1631

FTIR(neat): 3500, 2964, 2933, 2873, 2360, 2337, 1718, 1448, 1369, 1241, 1170, 1056, 1014 cm⁻¹

2-Hydroxy-3-methyl-2,4-diphenyl-but-3-enoic acid ethyl ester (4a).

4a

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added ethyl benzoylformate (**2a**) (36 mg, 0.2 mmol, 100 mol%) and 1-phenyl-1-propyne (**1b**) (76 μ L, 69.7 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen for 3 hours (control by TLC). The product (**4a**) was purified by flash column chromatography (4% EtOAc/hexane; *R_f* = 0.15 (10% EtOAc/hexane)) to afford 55 mg as a colorless oil (93% yield).

¹H NMR (300 MHz, CDCl₃): 7.66 (d, *J* = 6.9 Hz, 2H), 7.22-7.45 (m, 8H), 6.49 (s, 1H), 4.42 (dq, *J* = 7.2, 10.8 Hz, 1H), 4.34 (dq, *J* = 7.2, 10.8 Hz, 1H), 4.04 (s, 1H), 1.93 (s, 3H), 1.36 (t, *J* = 7.2 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 174.6, 140.1, 138.9, 137.5, 129.3, 129.1, 128.3, 128.2, 128.1, 127.5, 127.0, 83.2, 63.0, 15.6, 14.4.

HRMS Calcd. for C₁₉H₂₁O₃ (M+1): 297.1491, Found: 297.1487

FTIR (neat): 3492, 3058, 3024, 2981, 1724, 1599, 1491, 1447, 1383, 1367, 1295, 1243, 1152, 1094, 1062, 1029, 1015, 919, 860, 770, 747, 700, 636 cm⁻¹.

2-Hydroxy-3-methyl-2-(4-methoxy-phenyl)-4-phenyl-but-3-enoic acid ethyl ester (4b).

4b

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added ethyl 4-nitrobenzoylformate (**2b**) (44.6 mg, 0.2 mmol, 100 mol%) and 1-phenyl-1-propyne (**1b**) (76 μ L, 69.7 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen for 7 hours (control by TLC). The product (**4b**) was purified by flash column chromatography (5% EtOAc/hexane; R_f = 0.05 (10% EtOAc/hexane)) to afford 53 mg as a colorless oil (78% yield).

¹H NMR (400 MHz, CDCl₃): 8.23(d, *J* = 8.8, 2H), 7.85 (d, *J* = 8.8 Hz, 2H), 7.32-7.38 (m, 2H), 7.23-7.29 (m, 3H), 6.40 (s, 1H), 4.40 (dq, *J* = 7.2, 10.8 Hz, 1H), 4.32 (dq, *J* = 7.2, 10.8 Hz, 1H), 4.10 (s, 1H), 1.84 (d, *J* = 1.4 Hz, 3H), 1.34 (t, *J* = 7.2 Hz 3H).

¹³C NMR (100 MHz, CDCl₃): 173.4, 147.9, 147.1, 138.3, 136.8, 129.5, 129.3, 128.7, 128.5, 127.5, 123.4, 82.9, 63.6, 15.2, 14.3.

HRMS Calcd. for C₁₉H₂₀NO₅ (M+1): 342.1341, Found: 342.1342

FTIR (neat): 3481, 2982, 1728, 1646, 1603, 1521, 1491, 1445, 1348, 1243, 1156, 1078, 1041, 920, 854, 744, 698, 510 cm⁻¹.

2-Hydroxy-2-(4-methoxy-phenyl)-3-methyl-4-phenyl-but-3-enoic acid ethyl ester (4c).

4c

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added ethyl 4-methoxybenzoylformate (**2c**) (41.6 mg, 0.2 mmol, 100 mol%) and 1-phenyl-1-propyne (**1b**) (76 μ L, 69.7 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen for 6 hours (control by TLC). The product (**4c**) was purified by flash column chromatography (4% EtOAc/hexane; R_f = 0.10 (10% EtOAc/hexane)) to afford 63 mg as a colorless oil (97% yield).

¹H NMR (400 MHz, CDCl₃): 7.53 (d, *J* = 9 Hz, 2H), 7.29-7.35 (m, 2H), 7.19-7.28 (m, 3H), 6.90 (d, *J* = 9 Hz, 2H), 6.44 (s, 1H), 4.37 (dq, *J* = 7.2, 10.8 Hz, 1H), 4.30 (dq, *J* = 7.2, 10.8 Hz, 1H), 3.95 (s, 1H), 3.82 (s, 3H), 1.88 (d, *J* = 1.2 Hz, 3H), 1.33 (t, *J* = 7.2 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 174.8, 159.5, 139.1, 137.6, 132.1, 129.3, 129.0, 128.7, 128.3, 126.9, 113.6, 82.9, 62.9, 55.5, 15.5, 14.4.

HRMS Calcd. for C₂₀H₂₃O₄ (M+1): 327.1596, Found: 327.1596

FTIR (neat): 3493, 2980, 2836, 2360, 2342, 1724, 1608, 1509, 1463, 1443, 1366, 1300, 1248, 1177, 1074, 1032, 920, 860, 834, 761, 737, 699, 512 cm⁻¹.

2-(3-Bromo-phenyl)-2-hydroxy-3-methyl-4-phenyl-but-3-enoic acid methyl ester (4d).

4d

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added methyl 3-bromobenzoylformate (**2d**) (49.2 mg, 0.2 mmol, 100 mol%) and 1-phenyl-1-propyne (**1b**) (76 μ L, 69.7 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen or 6 hours (control by TLC). The product (**4d**) was purified by flash column chromatography (4% EtOAc/hexane; R_f = 0.12 (10% EtOAc/hexane)) to afford 53 mg as a colorless oil (73% yield).

¹H NMR (400 MHz, CDCl₃): 7.79 (dd, *J* = 1.8, 1.8 Hz, 1H), 7.54 (ddd, *J* = 8.0, 1.6, 1.2 Hz, 1H), 7.46 (ddd, *J* = 8.0, 2.0, 1.2 Hz, 1H), 7.31-7.61 (m, 2H), 7.21-7.28 (m, 4H), 6.38 (s, 1H), 3.96 (s, 1H), 3.88 (s, 3H), 1.84 (d, *J* = 1.2 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 174.6, 142.1, 138.6, 137.1, 131.4, 130.6, 129.8, 129.32, 129.30, 128.4, 127.3, 126.4, 122.6, 82.9, 53.9, 15.3.

HRMS Calcd. for C₁₈H₁₈O₃Br (M+1): 361.0439, Found: 361.0436

FTIR (neat): 3486, 3058, 3023, 2952, 2359, 2341, 1731, 1592, 1566, 1491, 1472, 1436, 1419, 1248, 1154, 1103, 1072, 1027, 996, 920, 784, 761, 745, 717, 697, 667, 508 cm⁻¹.

2-Hydroxy-3-methyl-4-phenyl-2-thiophen-2-yl-but-3-enoic acid ethyl ester (4e)

4e

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added ethyl thiophene-2-glyoxylate (**2e**) (36.8 mg, 0.2 mmol, 100 mol%) and 1-phenyl-1-propyne (**1b**) (76 μ L, 69.7 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen for 6 hours (control by TLC). The product (**4e**) was purified by flash column chromatography (4% EtOAc/hexane; R_f = 0.17 (10% EtOAc/hexane)) to afford 53 mg as a colorless oil (88% yield).

¹H NMR (400 MHz, CDCl₃): 7.28-7.35 (m, 3H), 7.18-7.26 (m, 4H), 7.02 (dd, *J* = 5.1, 3.5 Hz, 1H), 6.51 (s, 1H), 4.39 (dq, *J* = 10.8, 7.2 Hz, 1H), 4.36 (dq, *J* = 10.8, 7.2 Hz, 1H), 4.30 (s, 1H), 1.91 (d, *J* = 1.4 Hz, 3H), 1.37 (t, *J* = 7.2 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 173.7, 144.4, 138.3, 137.4, 129.3, 129.1, 128.3, 127.1, 127.0, 126.6, 125.9, 81.4, 63.4, 14.9, 14.3.

HRMS Calcd. for C₁₇H₁₉O₃S (M+1): 303.1055, Found: 303.1057

FTIR (neat): 3485, 2980, 2922, 2359, 2341, 1727, 1668, 1558, 1490, 1444, 1367, 1249, 1227, 1141, 1079, 1013, 920, 853, 758, 698, 668, 514 cm⁻¹.

**2-Hydroxy-2,3-dimethyl-4-phenyl-but-3-enoic acid ethyl ester and
2-Hydroxy-2-methyl-3-phenyl-pent-3-enoic acid ethyl ester (4f).**

4f

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added ethyl pyruvate (**2f**) (23.2 mg, 0.2 mmol, 100 mol%) and 1-phenyl-1-propyne (**1b**) (76 μ L, 69.7 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen for 3 hours (control by TLC). The products (**4f**) were purified by flash column chromatography (4% EtOAc/hexane; R_f = 0.10 (10% EtOAc/hexane)) to afford 42 mg as a colorless oil (90% yield, 1:1.45 mixture of regioisomers).

¹H NMR (400 MHz, CDCl₃): 7.20-7.36 (m, 7H), 7.08 (dt, *J* = 6.0, 2.1 Hz), 6.73 and 6.02 (minor: s, major: q, *J* = 6.9 Hz, 1H), 4.27 (minor: q, *J* = 7.2 Hz, 2H) and 4.16 (major: dq, *J* = 7.2, 10.8 Hz, 1H) and 4.09 (major: dq, *J* = 7.2, 10.8 Hz, 1H), 3.49 and 3.32 (minor: s, major: s, 1H), 1.87 (minor, d, *J* = 1.3, Hz, 3H), 1.64 and 1.57 (minor: s, major: s, 3H), 1.46 (major: d, *J* = 6.9 Hz, 3H), 1.31 and 1.20 (minor: t, *J* = 7.2 Hz, major: t, *J* = 7.2 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 176.0, 143.3, 138.8, 137.8, 137.7, 130.0, 129.3, 128.3, 128.2, 127.3, 126.9, 126.1, 124.1, 77.7, 76.9, 62.4, 62.1, 24.3, 24.1, 14.9, 14.4, 14.3.

HRMS Calcd. for C₁₄H₁₉O₃ (M+1): 235.1334, Found: 235.1339

FTIR (neat): 3504, 3080, 3055, 2982, 2937, 2859, 2359, 2341, 1726, 1492, 1443, 1373, 1249, 1134, 1095, 1019, 949, 921, 860, 769, 740, 703, 668, 566 cm⁻¹.

(E)-ethyl 2-hydroxy-2-(4-methoxyphenyl)-3-methylpent-3-enoate (5a).

To a solution of Ir(cod)₂BARF (12.6 mg, 0.010 mmol, 5.2 mol%), BIPHEP (5.5 mg, 0.011 mmol, 5.5 mol%), and triphenyl acetic acid (3.0 mg, 0.010 mmol, 5.2 mol%) in toluene (0.4 mL, 0.45 M) was added ethyl 4-methoxybenzoylformate (**2c**) (96 %, 40.7 mg, 0.19 mmol, 100 mol%) and butyne (**1c**) (excess). The system was purged with argon gas followed by hydrogen gas. Butyne (**1c**) (excess) was introduced along with hydrogen gas and the reaction mixture was allowed to stir at 80 °C for 2 hours (see Setup 1). The title compound (**5a**) was purified by flash column chromatography (R_f = 0.33, 20 %EtOAc/hexane) to afford 41.7 mg as colorless oil (84 % yield).

¹H NMR (400 MHz, CDCl₃): 7.45 (d, *J* = 8.4 Hz, 2H), 6.87 (d, *J* = 8.4 Hz, 2H), 5.38 (q, *J* = 7.2 Hz, 1H), 4.21-4.35 (m, 2H), 3.81 (s, 3H), 3.79 (s, 1H), 1.64 (s, 3H), 1.63, (d, *J* = 7.2 Hz, 3H), 1.29 (t, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 174.76, 159.02, 136.82, 131.95, 128.35, 123.71, 113.11, 82.32, 62.42, 55.17, 14.05, 13.54, 13.17

HRMS Calcd. for C₁₅H₂₀O₄ (M): 264.1362, Found: 264.1363

FTIR(neat): 3503, 2964, 2933, 2875, 2360, 2342, 1728, 1471, 1246, 1171, 1070 cm⁻¹

Figure 1.

(E)-ethyl 2-hydroxy-2-(4-methoxyphenyl)-3,5-dimethylhex-3-enoate (5b).

To a solution of Ir(cod)₂BARF (12.8 mg, 0.010 mmol, 5.3 mol%), DPPF (5.8 mg, 0.010 mmol, 5.3 mol%), and triphenyl acetic acid (2.9 mg, 0.010 mmol, 5.3 mol%) in toluene (0.4 mL, 0.45 M) was added ethyl 4-methoxybenzoylformate (**2c**) (96 %, 40.5 mg, 0.19 mmol, 100 mol%) and 4-methyl-2-pentyne (**1d**) (51.5 mg, 0.63 mmol, 330 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 35 °C under 1 atm of hydrogen gas for 1.5 hour. The title compound (**5b**) was purified by flash column chromatography (R_f = 3.9., 20 %EtOAc/hexane) to afford 46.6 mg as colorless oil (85 % yield).

¹H NMR (400 MHz, CDCl₃): 7.43 (d, *J* = 8.8 Hz, 2H), 6.86 (d, *J* = 8.8 Hz, 2H), 5.17 (d, *J* = 9.2 Hz, 1H), 4.18-4.36 (m, 2H), 3.81 (s, 3H), 3.75 (s, 1H), 2.56 (dseptet, *J* = 9.2, 7.2 Hz, 1H), 1.63 (s, 3H), 1.28 (t, *J* = 7.2 Hz, 3H), 0.95 (d, *J* = 7.2 Hz, 3H), 0.94 (d, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 174.73, 158.98, 136.50, 133.66, 128.42, 113.11, 82.23, 62.31, 55.18, 55.16, 27.26, 22.54, 22.49, 14.06, 13.22

HRMS Calcd. for C₁₇H₂₄O₄ (M): 292.1675, Found: 292.1678

FTIR(neat): 3487, 2957, 2869, 2361, 2341, 1719, 1069, 1509, 1463, 1299, 1240, 1169, 1069, 1033 cm⁻¹

4-(4-Fluoro-phenyl)-2-hydroxy-2-(4-methoxy-phenyl)-3-methyl-but-3-enoic acid ethyl ester (5c).

5c

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added ethyl 4-methoxybenzoylformate (**2c**) (41.6 mg, 0.2 mmol, 100 mol%) and 1-(4-fluorophenyl)-1-propyne (**1e**) (80.5 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen for 3 hours (control by TLC). The product (**5c**) was purified by flash column chromatography (4% EtOAc/hexane; R_f = 0.20 (10% EtOAc/hexane)) to afford 67 mg as a colorless oil (97% yield).

¹H NMR (400 MHz, CDCl₃): 7.51 (d, *J* = 9.0, 2H), 7.21 (dd, *J* = 5.5, 8.4 Hz, 2H), 7.01 (dd, *J* = 8.8, 8.8 Hz, 2H), 6.90 (d, *J* = 9.0 Hz, 2H), 6.38 (s, 1H), 4.36 (dq, *J* = 10.8, 7.2 Hz, 1H), 4.31 (dq, *J* = 10.8, 7.2 Hz, 1H), 3.95 (s, 1H), 3.82 (s, 3H), 1.86 (d, *J* = 1.2 Hz, 3H), 1.33 (t, *J* = 7.2 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 174.7, 163.0, 160.6, 159.5, 139.1, 133.6, 133.5, 131.9, 130.9, 130.8, 128.7, 128.0, 115.3, 115.1, 113.6, 82.8, 62.9, 55.5, 15.5, 14.4.

FTIR (neat): 3490, 2980, 2929, 2836, 1724, 1608, 1508, 1463, 1443, 1366, 1299, 1248, 1177, 1158, 1073, 1034, 864, 833, 779, 746, 580, 524 cm⁻¹.

4-(3-Fluoro-phenyl)-2-hydroxy-2-(4-methoxy-phenyl)-3-methyl-but-3-enoic acid ethyl ester (5d).

5d

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added ethyl 4-methoxybenzoylformate (**2c**) (41.6 mg, 0.2 mmol, 100 mol%) and 1-(3-fluorophenyl)-1-propyne (**1f**) (80.5 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen for 3 hours (control by TLC). The product (**5d**) was purified by flash column chromatography (4% EtOAc/hexane; R_f = 0.22 (10% EtOAc/hexane)) to afford 66 mg as a colorless oil (97% yield).

¹H NMR (400 MHz, CDCl₃): 7.51 (d, *J* = 9.0 Hz, 2H), 7.24-7.31 (m, 1H), 6.92-7.04 (m, 5H), 6.40 (s, 1H), 4.37 (dq, *J* = 10.8, 7.0 Hz, 1H), 4.31 (dq, *J* = 10.8, 7.0 Hz, 1H), 3.96 (s, 1H), 3.82 (s, 3H), 1.88 (d, *J* = 1.4 Hz, 3H), 1.33 (t, *J* = 7.0 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 174.6, 164.0, 161.6, 159.6, 140.3, 139.8, 131.8, 129.8, 128.7, 128.0, 125.1, 116.1, 115.9, 113.9, 113.7, 82.7, 63.0, 55.5, 15.6, 14.4.

HRMS Calcd. for C₂₀H₂₂O₄F (M+1): 345.1502, Found: 345.1502

FTIR (neat): 3488, 2980, 2932, 2836, 1724, 1609, 1580, 1509, 1485, 1442, 1366, 1299, 1249, 1178, 1148, 1074, 1034, 884, 835, 694 cm⁻¹.

(E)-ethyl 2-hydroxy-2,4-bis(4-methoxyphenyl)-3-methylbut-3-enoate (5e)

To a solution of Ir(cod)₂BARF (12.6 mg, 0.010 mmol, 5.3 mol%), BIPHEP (5.7 mg, 0.011 mmol, 5.7 mol%), and triphenyl acetic acid (2.8 mg, 0.010 mmol, 5.3 mol%) in toluene (0.4 mL, 0.45 M) was added ethyl 4-methoxybenzoylformate (**2c**) (96 %, 39.1 mg, 0.19 mmol, 100 mol%) and 1-(4-Methoxyphenyl)-1-propyne (**1g**) (86.8 mg, 0.59 mmol, 312 mol%). The system was purged with argon gas followed by hydrogen gas. The reaction was allowed to stir at 80 °C under 1 atm of hydrogen gas for 3 hours. The title compound (**5e**) was purified by flash column chromatography (R_f = 0.22, 20 %EtOAc/hexane) to afford 64.0 mg as colorless oil (99 % yield).

¹H NMR (400 MHz, CDCl₃): 7.52 (d, *J* = 8.8 Hz, 2H), 7.20 (d, *J* = 8.8 Hz, 2H), 6.88 (d, *J* = 8.8 Hz, 2H), 6.86 (d, *J* = 8.8 Hz, 2H), 6.37 (s, 1H), 4.25-4.40 (m, 2H), 3.94 (s, 1H), 3.81 (s, 3H), 3.80 (s, 3H), 1.88 (s, 3H), 1.32 (t, *J* = 7.2 Hz, 3H)

¹³C NMR (100 MHz, CDCl₃): 174.61, 159.16, 158.31, 137.18, 131.90, 130.31, 129.78, 128.49, 128.20, 113.45, 113.28, 82.71, 62.60, 55.21, 55.19, 15.33, 14.10

HRMS Calcd. for C₂₁H₂₄O₅ (M): 356.1624, Found: 356.1620

FTIR(neat): 3502, 2955, 2836, 2360, 2341, 1720, 1606, 1508, 1463, 1442, 1297, 1243, 1175, 1070, 1030, 913 cm⁻¹

2-Hydroxy-2-(4-methoxy-phenyl)-3-methyl-4-thiophen-2-yl-but-3-enoic acid ethyl ester (5f).

To a solution of Ir(cod)₂BARF (12.7 mg, 0.01 mmol, 5 mol%), BIPHEP (5.2 mg, 0.01 mmol, 5 mol%) and triphenylacetic acid (2.9 mg, 0.01 mmol, 5 mol%) in DCE (0.5 M, 0.4 mL) was added ethyl 4-methoxybenzoylformate (**2c**) (41.6 mg, 0.2 mmol, 100 mol%) and 2-prop-1-ynyl-thiophene (**1h**) (73 mg, 0.6 mmol, 300 mol%). The reaction was allowed to stir at 80 °C under 1 atm of hydrogen for 3 hours (control by TLC). The product (**5f**) was purified by flash column chromatography (4% EtOAc/hexane; R_f = 0.24 (10% EtOAc/hexane)) to afford 58 mg as a colorless oil (87% yield).

¹H NMR (400 MHz, CDCl₃): 7.49 (d, *J* = 8.5 Hz, 2H), 7.32 (d, *J* = 4.6 Hz, 1H), 6.98-7.08 (m, 2H) 6.92 (d, *J* = 8.5 Hz, 2H), 4.39 (dq, *J* = 11.0, 7.2 Hz, 1H), 4.32 (dq, *J* = 11.0, 7.2 Hz, 1H), 3.98 (s, 1H), 3.85 (s, 3H), 2.03 (s, 3H), 1.34 (t, *J* = 7.2 Hz, 3H).

¹³C NMR (100 MHz, CDCl₃): 174.6, 159.5, 140.5, 136.6, 132.1, 128.7, 128.4, 127.1, 125.9, 122.2, 113.7, 83.0, 63.0, 55.5, 16.8, 14.4.

HRMS Calcd. for C₁₈H₂₁O₄S (M+1): 333.1161, Found: 333.1159

FTIR (neat): 3487, 2978, 2835, 1724, 1608, 1509, 1463, 1442, 1365, 1299, 1249, 1176, 1070, 1033, 854, 835, 700, 576, 512 cm⁻¹.