

Supporting Information

KO'Bu-Promoted, Three-Component Domino Reaction of Arenes(indoles/phenols), C₆₀, and (Per/poly)fluoroarenes: Achieve Direct C-C Cross-Coupling of Fullerene with (Per/poly)fluoroarenes

Jun-Xiang Zhang, Ming-Wei Liu, Wan-Yu Wang, Rui-Lin Jia, Man-Qing Yan, Jun Xuan, and Fei Li*

Department of Chemistry, Anhui University; Key Laboratory of Structure and Functional Regulation of Hybrid Materials (Anhui University), Ministry of Education; Anhui Province Key Laboratory of Chemistry for Inorganic/Organic Hybrid Functionalized Materials and Key Laboratory of Functional Inorganic Materials of Anhui Province, Hefei, Anhui 230601, People's Republic of China


lfe@ahu.edu.cn

1. NMR Spectra of 3aa-6ic	S3-S110
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3aa	S3-S5
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ba	S5-S7
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ca	S8-S10
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3da	S10-S12
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ea	S13-S15
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3fa	S15-S17
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ga	S18-S20
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ha	S20-S22
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ia	S23-S25
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ab	S25-S27
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3eb	S28-S30
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3hb	S30-S32
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3jb	S33-S35
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3kb	S35-S37
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ac	S38-S40
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ec	S40-S42
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3gc	S43-S45
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3hc	S45-S47
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ic	S48-S50
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ad	S50-S52
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ed	S53-S55
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ae	S55-S57
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3je	S58-S60
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3le	S60-S62


¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3af	S63-S65
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 3ef	S65-S67
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5aa	S68-S70
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5ba	S70-S72
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5ca	S73-S75
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5bc	S75-S77
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5cc	S78-S80
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5ae	S80-S82
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5be	S83-S85
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5de	S85-S87
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5ee	S88-S90
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5af	S90-S92
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5bf	S93-S95
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5cf	S95-S97
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 5ff	S98-S100
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 6aa	S100-S102
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 6ea	S102-S104
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 6ac	S104-S106
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 6ec	S106-S108
¹ H NMR, ¹⁹ F NMR, and ¹³ C{ ¹ H} NMR spectra of product 6ic	S108-S110
2. Single-Crystal X-Ray Crystallography of 3ca , 3af , and 5ba	S111-S115
3. UV-vis absorption spectra of compounds 3aa-ia , 3ab-ef , 5aa-ff , 6aa-ic	S116-S117

1.NMR Spectra of 3aa-6ic


¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3aa




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3aa




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3aa




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3aa




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3aa




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ba




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ba




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3ba




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ba




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ba




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3ca




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ca




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ca




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ca




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ca




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3da




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3da





¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3da




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3da




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3da




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3ea




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ea




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ea




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ea




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ea




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3fa




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3fa




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3fa




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3fa




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3fa




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3ga




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ga




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ga




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ga




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ga




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 3ha




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ha




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3ha




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ha




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ha




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3ia




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ia




$^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ia




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ia




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ia




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ab





¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ab




¹³C{¹H} NMR (150 MHz, CS₂/d₆-acetone) of compound 3ab




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (150 MHz, CS_2/d_6 -acetone) of compound 3ab




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (150 MHz, CS_2/d_6 -acetone) of compound 3ab




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3eb




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3eb




$^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3eb




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3eb




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3eb




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3hb




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3hb





¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3hb




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3hb




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3hb




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3jb




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3jb




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3jb




Expanded ¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3jb




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3jb




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3kb




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3kb




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3kb




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3kb




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3kb




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3ac




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ac





$^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ac




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ac




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ac




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ec




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3ec




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ec




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ec




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3gc




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3gc




$^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3gc




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3gc




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3gc




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3hc




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3hc




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3hc




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3hc




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3hc




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3ic




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ic




$^{13}\text{C}\{\text{H}\}$ NMR (150 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ic




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (150 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ic




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (150 MHz, CS_2/d_6 -acetone) of compound 3ic




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 3ad




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ad




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 3ad




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ad




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ad




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3ed




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ed




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ed




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3ed




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3ed




^1H NMR (400 MHz, CS_2/d_6 -DMSO) of compound 3ae




¹⁹F NMR (376 MHz, CS₂/d₆-DMSO) of compound 3ae




¹³C{¹H} NMR (100 MHz, CS₂/d₆-DMSO) of compound 3ae




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-DMSO}$) of compound 3ae




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-DMSO}$) of compound 3ae




¹H NMR (400 MHz, CS₂/d₆-DMSO) of compound 3je




¹⁹F NMR (376 MHz, CS₂/d₆-DMSO) of compound 3je




$^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-DMSO}$) of compound 3je




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-DMSO}$) of compound 3je




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-DMSO}$) of compound 3je




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-DMSO}$) of compound 3le




¹⁹F NMR (376 MHz, CS₂/d₆-DMSO) of compound 3le





¹³C{¹H} NMR (100 MHz, CS₂/d₆-DMSO) of compound 3le




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-DMSO}$) of compound 3le




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-DMSO}$) of compound 3le




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 3af




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3af




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3af




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 3af




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 3af




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 3ef




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 3ef




¹³C{¹H} NMR (150 MHz, CS₂/d₆-acetone) of compound 3ef




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (150 MHz, CS_2/d_6 -acetone) of compound 3ef




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (150 MHz, CS_2/d_6 -acetone) of compound 3ef




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 5aa




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5aa




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5aa




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5aa




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5aa




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 5ba




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5ba




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 5ba




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5ba




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5ba




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 5ca




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5ca




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5ca




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5ca




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5ca




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5bc




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5bc




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 5bc




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5bc




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5bc




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 5cc




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5cc




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5cc




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5cc




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5cc




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 5ae




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5ae




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 5ae




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5ae




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5ae




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 5be




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5be




$^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5be




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5be




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5be




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 5de




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5de




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 5de




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5de




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5de




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 5ee




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5ee




$^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5ee




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5ee




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5ee




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 5af




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5af




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 5af




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5af




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 5af




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 5bf




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5bf




$^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5bf




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5bf




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5bf




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 5cf




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5cf




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 5cf




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5cf




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5cf




¹H NMR (400 MHz, CS₂/d₆-acetone) of compound 5ff




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 5ff




$^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5ff




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5ff




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -acetone) of compound 5ff




^1H NMR (400 MHz, CS_2/d_6 -DMSO) of compound 6aa




¹⁹F NMR (376 MHz, CS₂/d₆-DMSO) of compound 6aa




¹³C{¹H} NMR (100 MHz, CS₂/d₆-DMSO) of compound 6aa




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CS_2/d_6 -DMSO) of compound 6aa




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 6ea




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 6ea




¹³C{¹H} NMR (150 MHz, CS₂/d₆-acetone) of compound 6ea




Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (150 MHz, CS_2/d_6 -acetone) of compound 6ea




^1H NMR (400 MHz, CS_2/d_6 -acetone) of compound 6ac




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 6ac




¹³C{¹H} NMR (150 MHz, CS₂/d₆-acetone) of compound 6ac




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (150 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 6ac




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 6ec




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 6ec




¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 6ec




Expanded $^{13}\text{C}\{\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 6ec




^1H NMR (400 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 6ic




¹⁹F NMR (376 MHz, CS₂/d₆-acetone) of compound 6ic



¹³C{¹H} NMR (100 MHz, CS₂/d₆-acetone) of compound 6ic



Expanded $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{CS}_2/d_6\text{-acetone}$) of compound 6ic



2. Single-Crystal X-Ray Crystallography of 3ca, 3af, and 5ba




Figure S1 ORTEP diagrams of **3ca** with 30% thermal ellipsoids.

Black block crystals of **3ca** suitable for X-ray diffraction were obtained from isopropyl alcohol diffusion into its CS₂ solution. Single-crystal X-ray diffraction data were collected on a diffractometer (DECTRIS PILATUS 300K, STOE & Cie GmbH) equipped with a CCD area detector using graphite-monochromated Cu K α radiation ($\lambda = 1.54184 \text{ \AA}$) in the scan range $9.172^\circ < 2\theta < 124.998$. The structure was solved with direct methods using SHELXS-97 and refined with full-matrix least-squares refinement using the SHELXL-97 program within OLEX2. Crystallographic data have been deposited in the Cambridge Crystallographic Data Centre as deposition number CCDC 2177898.

Table S1 Crystal data and structure refinement for 3ca

Identification code	3ca
Empirical formula	C ₇₉ H ₇ F ₈ N ₃
Formula weight	1149.88
Temperature/K	120
Crystal system	monoclinic
Space group	P2 ₁ /n
a/ \AA	10.156
b/ \AA	30.702
c/ \AA	14.665
$\alpha/^\circ$	90
$\beta/^\circ$	90.66

$\gamma/^\circ$	90
Volume/ \AA^3	4572.1
Z	4
$\rho_{\text{calc}} \text{g/cm}^3$	1.671
μ/mm^{-1}	1.016
F(000)	2296.0
Crystal size/ mm^3	0.03 \times 0.02 \times 0.015
Radiation	CuK α ($\lambda = 1.54186$)
2 Θ range for data collection/ $^\circ$	9.172 to 124.998
Index ranges	-9 \leq h \leq 11, -35 \leq k \leq 34, -12 \leq l \leq 16
Reflections collected	13186
Independent reflections	6994 [$R_{\text{int}} = 0.0299$, $R_{\text{sigma}} = 0.0263$]
Data/restraints/parameters	6994/0/620
Goodness-of-fit on F ²	1.158
Final R indexes [I $>= 2\sigma(I)$]	$R_1 = 0.1015$, wR ₂ = 0.2380
Final R indexes [all data]	$R_1 = 0.1186$, wR ₂ = 0.2650
Largest diff. peak/hole / e \AA^{-3}	0.54/-0.46




Figure S2 ORTEP diagrams of **3af** with 30% thermal ellipsoids.

Black block crystals of **3af** suitable for X-ray diffraction were obtained from acetonitrile diffusion into its CS₂ solution. Single-crystal X-ray diffraction data were collected on a diffractometer (DECTRIS PILATUS 300K, STOE & Cie GmbH) equipped with a CCD area detector using graphite-monochromated Cu K α radiation ($\lambda = 1.54184 \text{ \AA}$) in the scan range $11.584^\circ < 2\theta < 139.068$. The structure was solved with direct methods using SHELXS-97 and refined with full-matrix least-squares refinement using the SHELXL-97 program within OLEX2. Crystallographic data have been deposited in the Cambridge Crystallographic Data Centre as deposition number CCDC 2177886.

Table S2 Crystal data and structure refinement for 3af	
Identification code	3af
Empirical formula	C ₉₂ H ₁₇ F ₈ NO ₅ S ₂
Formula weight	1432.19
Temperature/K	293(2)
Crystal system	triclinic
Space group	P-1
a/ \AA	10.7362(7)
b/ \AA	17.4371(13)
c/ \AA	17.4852(12)
$\alpha/^\circ$	118.369(5)
$\beta/^\circ$	93.833(6)
$\gamma/^\circ$	94.424(6)
Volume/ \AA^3	2851.6(4)
Z	2
$\rho_{\text{calcd}}/\text{cm}^3$	1.668
μ/mm^{-1}	1.676
F(000)	1440.0
Crystal size/mm ³	0.17 × 0.15 × 0.10
Radiation	CuK α ($\lambda = 1.54186$)
2 Θ range for data collection/ $^\circ$	11.584 to 139.068
Index ranges	-12 ≤ h ≤ 11, -14 ≤ k ≤ 20, -20 ≤ l ≤ 12
Reflections collected	16586
Independent reflections	10089 [R _{int} = 0.0482, R _{sigma} = 0.0484]
Data/restraints/parameters	10089/0/969
Goodness-of-fit on F ²	1.092
Final R indexes [I>=2σ (I)]	R ₁ = 0.1001, wR ₂ = 0.2474
Final R indexes [all data]	R ₁ = 0.1224, wR ₂ = 0.2726
Largest diff. peak/hole / e \AA^{-3}	0.92/-0.48




Figure S3 ORTEP diagrams of **5ba** with 30% thermal ellipsoids.

Black block crystals of **5ba** suitable for X-ray diffraction were obtained from slow evaporation of its solution in a mixture of CS₂, d₆-acetone and iso-propyl alcohol at room temperature. Single-crystal X-ray diffraction data were collected on a diffractometer (DECTRIS PILATUS 300K, STOE & Cie GmbH) equipped with a CCD area detector using graphite-monochromated Cu K α radiation ($\lambda = 1.54184 \text{ \AA}$) in the scan range $9.23^\circ < 2\theta < 139.01$. The structure was solved with direct methods using SHELXS-97 and refined with full-matrix least-squares refinement using the SHELXL-97 program within OLEX2. Crystallographic data have been deposited in the Cambridge Crystallographic Data Centre as deposition number CCDC 2177889.

Table S3 Crystal data and structure refinement for 5ba

Identification code	5ba
Empirical formula	C ₇₈ H ₈ F ₈ N ₂ O
Formula weight	1140.86
Temperature/K	293(2)
Crystal system	monoclinic
Space group	P2 ₁ /c
a/ \AA	21.6590(9)
b/ \AA	10.0799(4)
c/ \AA	20.0636(8)
$\alpha/^\circ$	90

$\beta/^\circ$	99.028(3)
$\gamma/^\circ$	90
Volume/ \AA^3	4326.0(3)
Z	4
$\rho_{\text{calc}} \text{g/cm}^3$	1.752
μ/mm^{-1}	1.077
F(000)	2280.0
Crystal size/mm ³	0.3 × 0.2 × 0.1
Radiation	CuK α ($\lambda = 1.54186$)
2 Θ range for data collection/°	9.23 to 139.008
Index ranges	-26 ≤ h ≤ 24, -12 ≤ k ≤ 10, -16 ≤ l ≤ 23
Reflections collected	15740
Independent reflections	7644 [R _{int} = 0.0539, R _{sigma} = 0.0496]
Data/restraints/parameters	7644/0/779
Goodness-of-fit on F ²	1.021
Final R indexes [I>=2σ (I)]	R ₁ = 0.0932, wR ₂ = 0.2302
Final R indexes [all data]	R ₁ = 0.1354, wR ₂ = 0.2747
Largest diff. peak/hole / e \AA^{-3}	0.51/-0.46

3. UV-vis absorption spectra of compounds 3aa-ia, 3ab-ef, 5aa-ff, 6aa-ic




Figure S4 UV-vis spectra of 3aa-ia (solvent: CHCl₃)




Figure S5 UV-vis spectra of 3ab-ef (solvent: CHCl₃)




Figure S6 UV-vis spectra of **5aa-ff** (solvent: CHCl₃)




Figure S7 UV-vis spectra of **6aa-ic** (solvent: CHCl₃)