Kinetic Resolutions of Indolines by a Non-Enzymatic Acylation Catalyst

Forrest O. Arp and Gregory C. Fu*

Department of Chemistry, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139

Supporting Information

I. General

Lithium bromide (Alfa Aesar; ultra dry, 99.998%) was crushed into a fine powder under nitrogen, and 2-methylindoline (Aldrich) was purified by distillation from calcium hydride. Toluene was dried by passage through an activated alumina column. All other chemicals were reagent grade and used as received.

HPLC analyses were performed on an Agilent 1100 Series system with Daicel Chiralpak® columns in hexanes/isopropanol solvent mixtures.

All reactions were conducted under nitrogen, unless otherwise noted.

The conversions and selectivity factors were calculated by the method described by Kagan.¹

II. Preparation of Materials

The yields in this section have not been optimized.

Catalyst 1. *n*-Butyllithium (1.6 M in hexanes; 2.1 mL, 3.4 mmol) was added to a solution of penta(3,5-dimethylphenyl)cyclopentadiene (2.00 g, 3.41 mmol) in dry THF (20 mL).

A suspension of crushed FeCl₂ (433 mg, 3.42 mmol) in dry THF (250 mL) was prepared in a 500-mL round-bottomed flask. The flask was fitted with a rubber septum, equipped with a nitrogen-inlet needle, and then cooled to 0 °C in an ice bath. Next, the cyclopentadienyllithium solution was added by cannula into the FeCl₂ solution, and the resulting green mixture was stirred at 0 °C for 15 min.

In a separate flask, 4-dimethylaminopyridine² (0.529 g, 2.84 mmol) was dissolved in dry THF (15 mL) and then treated with *n*-butyllithium (1.6 M in hexanes; 1.8 mL, 2.9 mmol). The mixture was stirred at r.t. for 10 min, and then it was transferred by cannula into the cold mixture that contained the iron complex (previous paragraph). The resulting dark-yellow solution was placed into an oil bath preheated to 60 °C and stirred for 3 h. During that time, the solution turned dark red.

The solvents were removed by rotary evaporation, and the residue was purified by flash chromatography (CH₂Cl₂ and then ethyl acetate). The dark-purple fractions were concentrated to provide the product as a glassy, dark-purple solid (1.68 mg, 72%). Catalyst **1** is air-stable (for months).

¹H NMR (CDCl₃, 400 MHz): δ 8.11 (d, *J* = 5.4 Hz, 1H), 6.72 (s, 5H), 6.57 (s, 10H), 5.73 (d, *J* = 5.4 Hz, 1H), 4.97-4.91 (m, 2H), 4.16 (t, *J* = 2.8 Hz, 1H), 3.53-3.42 (br, 4H), 2.09 (s, 30H), 1.94-1.88 (br, 4H);

¹³C NMR (CDCl₃, 100 MHz): δ 156.5, 152.6, 149.6, 135.3, 135.1, 130.3, 127.2, 96.5, 85.2, 77.9, 77.0, 69.0, 65.1, 49.7, 25.3, 21.2;

IR (film) 3434, 2918, 1642, 1600, 1536, 1485, 1462, 1401, 1339, 1025, 904, 848, 759, 706 cm⁻¹;

HRMS calculated for C₅₇H₅₈FeN₂ 826.3949, found 826.3945;

mp 154-157 °C.

The enantiomers of catalyst **1** were separated by semi-preparative chiral HPLC: Regis (*R,R*)-Whelk-O 2 10/100, 25 cm × 10 mm; diethylamine/CH₂Cl₂/hexanes, 0.4:20:80; 2.50 mL/min flow rate; 25.0 mg catalyst in 0.5 mL per injection. Enantiomer (+)-**1** ([α]_D²³ = +1100 (*c* = 0.021, CHCl₃)) was collected from 8.8 min to 11.1 min, and (–)-**1** was collected from 11.8 min to 16.0 min.

The absolute configuration of (+)-**1** was assigned by X-ray crystallography (see Section IV).

4-tert-Butyl-2-phenyloxazol-5(4H)-one [71953-55-4]. NaOH (3.4 g, 85 mmol) and then benzoyl chloride (4.9 mL, 42 mmol) were added to a solution of *tert*-leucine (5.03 g, 38.3 mmol) in water (200 mL) in a large Erlenmeyer flask. This heterogeneous mixture was stirred overnight, during which time it became homogeneous. The product amide was then precipitated by the slow addition of aqueous HCl (2 M), isolated by filtration, and dried thoroughly under high vacuum. This chalky white solid was dissolved in CH₂Cl₂ (50 mL), and then 1,3-dicyclohexylcarbodiimide (8.05 g, 39.0 mmol) was added. The resulting heterogeneous mixture was stirred open to air at r.t. overnight. The solution was filtered, and then the filtrate was washed with aqueous HCl (2 M; 100 mL), dried (Na₂SO₄), and concentrated. The desired product was purified by flash chromatography through a short pad of silica gel (5% ethyl acetate in hexanes), which afforded a white solid (6.95 g, 84%, two steps).

¹H NMR (CDCl₃, 400 MHz): δ 8.03-8.01 (m, 2H), 7.59-7.55 (m, 1H), 7.50-7.47 (m, 2H), 4.09 (s, 1H), 1.15 (s, 9H);

¹³C NMR (CDCl₃, 125 MHz): δ 176.9, 161.2, 132.6, 128.7, 127.8, 125.9, 74.0, 35.9, 26.1.

4-tert-Butyl-2-phenyloxazol-5-yl acetate. A 500-mL round-bottomed flask was capped with a rubber septum, flushed with nitrogen, and charged with sodium hydride (845 mg, 35.2 mmol) and 4-*tert*-butyl-2-phenyloxazol-5(4H)-one (6.95 g, 32.0 mmol). Dry THF (250 mL) was added, and the resulting heterogeneous mixture was cooled to –15 °C. 15-Crown-5 (7.75 g, 35.2 mmol) was added, and the resulting canary yellow, bubbling solution was stirred at –15 °C for 15 min. Then, acetyl chloride (2.76 g, 35.2 mmol) was added, and the resulting colorless solution was stirred for 20 min. The reaction mixture was then concentrated and purified by flash chromatography (5% ethyl acetate in hexanes) to remove most of the crown ether and sodium salts. The residue was further purified by flash chromatography (2.5% ethyl acetate in hexanes) to afford the product as a white solid (6.74 g, 81%).

¹H NMR (CDCl₃, 400 MHz): δ 7.97-7.95 (m, 2H), 7.43-7.41 (m, 3H), 2.35 (s, 3H), 1.33 (s, 9H);

¹³C NMR (CDCl₃, 100 MHz): δ 168.1, 154.4, 143.4, 132.1, 129.9, 128.5, 127.5, 125.8, 31.0, 28.9, 20.3;

IR (film) 2959, 2870, 1798, 1636, 1556, 1486, 1450, 1371, 1322, 1170, 1093, 1075, 1007, 873, 776, 724, 691 cm^{-1} ;

EIMS (70 eV) m/z : M^+ (calculated for $\text{C}_{15}\text{H}_{17}\text{NO}_3$: 259) 259, 218, 203, 174, 156, 133, 106, 84, 51;

mp 76-78 $^{\circ}\text{C}$.

Reduction by Sodium Cyanoborohydride: General Procedure.³ Sodium cyanoborohydride (3.0 equiv) was carefully dissolved in glacial acetic acid, and the resulting bubbling clear solution was added by pipette to a stirred solution of the indole in glacial acetic acid. This clear, colorless mixture was allowed to stir open to air for 12 h at r.t. Then, the mixture was transferred to an Erlenmeyer flask, and a saturated aqueous K_2CO_3 solution was added until the solution had become basic. The resulting mixture was extracted with diethyl ether (2×100 mL), and the combined organic phases were dried (K_2CO_3), concentrated, and purified by flash chromatography.

2-Propylindoline [76916-57-9]. This indoline was prepared from 2-propylindole⁴ (906 mg, 5.69 mmol) and sodium cyanoborohydride (1.1 g, 18 mmol) via the general reduction procedure. After flash chromatography (5% ethyl acetate in hexanes), the product was obtained as a clear, colorless oil (508 mg, 55%).

^1H NMR (CDCl_3 , 300 MHz): δ 7.10-6.99 (m, 2H), 6.72-6.60 (m, 2H), 3.92-3.82 (m, 2H), 3.14 (dd, $J = 8.6, 15.5$ Hz, 1H), 2.69 (dd, $J = 8.5, 15.5$ Hz, 1H), 1.66-1.57 (m, 2H), 1.49-1.35 (m, 2H), 0.98 (t, $J = 7.2$ Hz, 3H);

^{13}C NMR (CDCl_3 , 75 MHz): δ 151.0, 128.9, 127.2, 124.6, 118.4, 109.0, 59.8, 39.0, 36.1, 19.7, 14.1.

2-(4-Phenylbut-1-ynyl)benzenamine [736184-66-0]. A 100-mL round-bottomed flask was charged with 2-iodoaniline (2.30 g, 10.5 mmol), *trans*-dichlorobis(triphenylphosphine)palladium (73.7 mg, 0.105 mmol), and CuI (100 mg, 0.53 mmol).⁴ The flask was fitted with a rubber septum and purged with nitrogen for 5 min. Triethylamine (20 mL) and then but-3-ynylbenzene (2.39 g, 18.4 mmol) were added by syringe, and the resulting reaction mixture was stirred at r.t. for 24 h. Then, the triethylamine was removed in vacuo, and the resulting brown residue was purified by flash chromatography (5% ethyl acetate in hexanes), which afforded the product as an orange oil (2.25 g, 97%).

^1H NMR (CDCl_3 , 400 MHz): δ 7.37-7.30 (m, 4H), 7.28-7.23 (m, 2H), 7.11-7.07 (m, 1H), 6.69-6.65 (m, 2H), 3.99 (br s, 2H), 2.96 (t, $J = 7.3$ Hz, 2H), 2.82 (t, $J = 7.1$ Hz, 2H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 147.7, 140.6, 131.9, 128.9, 128.6, 128.4, 126.3, 117.7, 114.0, 108.5, 94.6, 77.9, 35.1, 21.7.

2-Phenethylindole [98055-06-2]. A 100-mL round-bottomed flask was charged with 2-(4-phenylbut-1-ynyl)benzenamine (2.25 g, 10.2 mmol), CuI (387 mg, 2.03 mmol), and DMF (10 mL).⁴ The flask was fitted with a reflux condenser, and the system was purged with nitrogen for 5 min. The heterogeneous reaction mixture was stirred at reflux for 12 h. Then, most of the DMF was removed in vacuo, and the residue was purified by flash chromatography (5% ethyl acetate in hexanes), which furnished the product as an off-white crystalline solid (643 mg, 29%).

^1H NMR (CDCl_3 , 400 MHz): δ 7.77 (br s, 1H), 7.56 (d, $J = 7.6$ Hz, 1H), 7.36-7.32 (m, 2H), 7.29-7.24 (m, 4H), 7.16-7.08 (m, 2H), 6.31 (s, 1H), 3.14-3.05 (m, 4H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 141.2, 139.0, 135.8, 128.6, 128.5, 128.4, 126.3, 121.1, 119.8, 119.6, 110.3, 99.8, 35.6, 30.1.

2-Phenethylindoline. This indoline was prepared from 2-phenethylindole (643 mg, 2.91 mmol) and sodium cyanoborohydride (550 mg, 8.8 mmol) via the general reduction procedure. After flash chromatography (5% ethyl acetate in hexanes), the product was obtained as an off-white solid (643 mg, 84%).

^1H NMR (CDCl_3 , 400 MHz): δ 7.33-7.29 (m, 2H), 7.23-7.19 (m, 3H), 7.09 (d, $J = 7.2$ Hz, 1H), 7.02 (t, $J = 7.4$ Hz, 1H), 6.70 (t, $J = 7.4$ Hz, 1H), 6.61 (d, $J = 7.7$ Hz, 1H), 3.93-3.86 (m, 1H), 3.17 (dd, $J = 8.7, 15.5$ Hz, 1H), 2.79-2.68 (m, 3H), 2.00-1.94 (m, 2H), 1.58 (br s, 1H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 150.8, 141.7, 128.7, 128.4, 128.3, 127.2, 125.9, 124.6, 118.5, 109.1, 59.5, 38.4, 36.1, 32.9;

IR (film) 3368, 3022, 2922, 2848, 1607, 1484, 1466, 1454, 1319, 1246, 1017, 745 cm^{-1} ;

EIMS (70 eV) m/z : M^+ (calculated for $\text{C}_{16}\text{H}_{17}\text{N}$: 223) 223, 207, 130, 118, 91, 65;

mp 39-41 $^{\circ}\text{C}$.

2-((*tert*-Butyldimethylsilyloxy)methyl)indoline [321744-15-4]. Lithium aluminum hydride (1.0 M in THF; 37 mL, 37 mmol) was carefully added to a solution of ($\pm$)-indoline-2-carboxylic acid (2.0 g, 12.3 mmol) in dry THF (100 mL). The mixture was

stirred for 1 h at r.t., and then the reaction was carefully quenched by the addition of aqueous sodium potassium tartrate (10% by weight; 100 mL). The resulting mixture was stirred for an additional hour, and then it was filtered. The filtrate was concentrated to a yellow oil, which was purified by flash chromatography (CH₂Cl₂) to afford the aminoalcohol.

The aminoalcohol (1.00 g, 6.70 mmol) was dissolved in CH₂Cl₂ (100 mL), and then imidazole (500 mg, 7.4 mmol) and *tert*-butyldimethylsilyl chloride (1.1 g, 7.4 mmol) were added. The resulting mixture was stirred at r.t. for 2 h, and then it was filtered. The filtrate was concentrated and then purified by flash chromatography (5% ethyl acetate in hexanes), which afforded the product as a clear yellow oil (1.50 g, 46%, two steps).

¹H NMR (CDCl₃, 500 MHz): δ 7.09-7.00 (m, 2H), 6.71-6.63 (m, 2H), 4.23 (br s, 1H), 3.98-3.91 (m, 1H), 3.62-3.52 (m, 2H), 3.11 (dd, *J* = 9.0, 15.7 Hz, 1H), 2.65 (dd, *J* = 5.7, 15.8 Hz, 1H), 0.92 (s, 9H), 0.07 (s, 6H);

¹³C NMR (CDCl₃, 125 MHz): δ 150.5, 128.1, 127.3, 124.7, 118.4, 109.4, 66.6, 60.4, 32.0, 25.9, 18.3, -5.3.

***cis*-5,6,7,8,8a,9-hexahydro-4bH-carbazole [4828-96-0].** This indoline was prepared from 6,7,8,9-tetrahydrocarbazole (1.00 g, 5.84 mmol) and sodium cyanoborohydride (1.1 g, 18 mmol) via the general reduction procedure. After flash chromatography (5% ethyl acetate in hexanes), the product was obtained as a white solid (779 mg, 78%).

¹H NMR (CDCl₃, 300 MHz): δ 7.13-7.03 (m, 2H), 6.80-6.69 (m, 2H), 3.78-3.72 (m, 1H), 3.68 (br s, 1H), 3.13 (app q, *J* = 6.6 Hz, 1H), 1.83-1.77 (m, 2H), 1.83-1.53 (m, 3H), 1.47-1.35 (m, 3H);

¹³C NMR (CDCl₃, 75 MHz): δ 150.7, 133.5, 126.9, 123.1, 118.7, 110.1, 59.6, 40.8, 29.1, 26.9, 22.5, 21.6.

***cis*-5,5a,6,7,8,9,10,10a-octahydrocyclohepta[b]indole [886024-65-3].** This indoline was prepared from 5,6,7,8,9,10-hexahydrocyclohepta[b]indole (Ryan Scientific; 650 mg, 3.51 mmol) and sodium cyanoborohydride (660 mg, 10.5 mmol) via the general reduction procedure. After flash chromatography (5% ethyl acetate in hexanes), an inseparable mixture of the *cis* and *trans* isomers was obtained as a white solid (569 mg; ~9:1).

To facilitate isolation of the *cis* isomer, the mixture was acylated (diethyl ether (25 mL), triethylamine (515 μ L, 1.3 equiv), and acetyl chloride (222 μ L, 1.1 equiv); 15 min at r.t.) and purified by flash chromatography (10% $\rightarrow$ 15% ethyl acetate in hexanes), which furnished the isomerically pure *N*-acetylated *cis* diastereomer.

The title compound was then obtained by dissolving the *N*-acetylated *cis* isomer in dioxane (10 mL), adding concentrated aqueous HCl (5.0 mL), and refluxing the reaction mixture overnight. Then, the mixture was concentrated and purified by flash chromatography (5% ethyl acetate in hexanes), which afforded the desired product as a feathery white solid (482 mg, 73%, three steps).

^1H NMR (CDCl_3 , 300 MHz): δ 7.04-6.99 (m, 2H), 6.73-6.67 (m, 1H), 6.58-6.55 (m, 1H), 4.05 (dt, J = 5.5, 9.8 Hz, 1H), 3.65 (br s, 1H), 3.48 (dt, J = 3.8, 10.6 Hz, 1H), 2.03-1.68 (m, 7H), 1.47-1.29 (m, 3H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 150.1, 133.5, 127.3, 124.1, 118.1, 108.4, 63.4, 46.7, 33.5, 31.31, 31.34, 28.7, 26.0.

***cis*-2,3-Dimethylindoline [10276-90-1] and *trans*-2,3-dimethylindoline [7356-42-5].**

These indolines were prepared from 2,3-dimethylindole (2.28 g, 15.7 mmol) and sodium cyanoborohydride (3.0 g, 48 mmol) via the general reduction procedure. After flash chromatography (3% $\rightarrow$ 5% ethyl acetate in hexanes), *cis*-2,3-dimethylindoline (221 mg, 10%) and *trans*-2,3-dimethylindoline (1.09 g, 47%) were obtained as yellow oils.

***cis*-2,3-Dimethylindoline.**

^1H NMR (CDCl_3 , 500 MHz): δ 7.09-7.01 (m, 2H), 6.76-6.72 (m, 1H), 6.63-6.62 (m, 1H), 3.99-3.93 (m, 1H), 3.65 (br s, 1H), 3.31-3.25 (m, 1H), 1.19 (d, J = 7.2 Hz, 3H), 1.15 (d, J = 6.5 Hz, 3H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 150.1, 134.2, 127.2, 123.8, 118.7, 109.3, 58.3, 39.4, 16.3, 13.6.

***trans*-2,3-Dimethylindoline.**

^1H NMR (CDCl_3 , 500 MHz): δ 7.24-7.21 (m, 2H), 6.95-6.91 (m, 1H), 6.77-6.76 (m, 1H), 3.96 (br s, 1H), 3.61-3.56 (m, 1H), 3.02-2.96 (m, 1H), 1.50-1.47 (m, 6H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 150.3, 134.0, 127.0, 122.9, 118.2, 108.8, 63.6, 44.0, 20.2, 16.9.

Ethyl 2-(5-methoxy-2-methyl-1H-indol-3-yl)acetate [17536-38-8]. A 50-mL round-bottomed flask was charged with bis(2-oxo-3-oxazolidinyl)phosphinic chloride (1.38 g, 5.42 mmol) and 2-(5-methoxy-2-methyl-1H-indol-3-yl)acetic acid (Aldrich; 1.08 g, 4.93 mmol) and then flushed with nitrogen. Dry CH_2Cl_2 (15 mL) and triethylamine (1.51 mL, 10.8 mmol) were added, and the solution was stirred at r.t. for 5 min. Then, dry ethanol (1.0 mL, 17 mmol) was added, and the mixture was stirred overnight. The solution was then extracted with water (2×50 mL), and the organic layer was dried (Na_2SO_4), concentrated, and purified by flash chromatography through a short pad of silica gel (33% ethyl acetate in hexanes), which afforded the desired product as a viscous yellow oil (0.97 g, 86%).

^1H NMR (CDCl_3 , 400 MHz): δ 7.77 (br s, 1H), 7.14 (d, $J = 8.5$ Hz, 1H), 7.02-7.01 (m, 1H), 6.79-6.77 (m, 1H), 4.14 (q, $J = 7.1$ Hz, 2H), 3.87 (s, 3H), 3.65 (s, 2H), 2.39 (s, 3H), 1.26 (t, $J = 7.0$ Hz, 3H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 172.2, 153.9, 133.5, 130.1, 128.8, 110.9, 110.8, 104.3, 100.3, 60.6, 55.8, 30.5, 14.2, 11.6.

cis-Ethyl 2-(5-methoxy-2-methylindolin-3-yl)acetate. A 25-mL round-bottomed flask was charged with ethyl 2-(5-methoxy-2-methyl-1H-indol-3-yl)acetate (657 mg, 2.66 mmol), trifluoroacetic acid (8.0 mL), and triethylsilane (2.12 mL, 13.3 mmol).⁵ The flask was fitted with a reflux condenser, heated to 70 °C, and stirred for 48 h. Then, additional triethylsilane (2.12 mL, 13.3 mmol) was added, and the reaction was stirred for another 48 h. Next, the mixture was carefully transferred by pipette into an aqueous K_2CO_3 solution (5% by weight; 150 mL) and extracted with diethyl ether (2×100 mL). The combined organic layers were dried (K_2CO_3), concentrated, and purified by flash chromatography (15% $\rightarrow$ 20% ethyl acetate in hexanes), which afforded the desired product as a yellow oil (401 mg, 60%; cis isomer), along with recovered starting material (43 mg, 7%).

^1H NMR (CDCl_3 , 400 MHz): δ 6.70 (s, 1H), 6.63-6.61 (m, 1H), 6.61-6.56 (m, 1H), 4.19 (q, $J = 7.1$ Hz, 2H), 4.06 (dq, $J = 6.7, 6.5$ Hz, 1H), 3.75 (s, 3H), 3.68-3.62 (m, 1H), 3.40 (br s, 1H), 2.65-2.53 (m, 2H), 1.28 (t, $J = 7.1$ Hz, 3H), 1.15 (d, $J = 6.5$ Hz, 3H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 172.6, 153.4, 144.0, 133.0, 112.6, 111.0, 110.0, 60.5, 58.2, 55.8, 41.6, 33.7, 16.2, 14.1;

IR (film) 3435, 2981, 2832, 2099, 1727, 1639, 1491, 1232, 1156, 1033 cm^{-1} ;

EIMS (70 eV) m/z : M^+ (calculated for $C_{14}H_{19}NO_3$: 249) 249, 234, 222, 204, 178, 160, 142, 117.

5-Methoxy-2-methylindoline [41568-27-8]. This indoline was prepared from 5-methoxy-2-methylindole (500 mg, 3.10 mmol) and sodium cyanoborohydride (580 mg, 9.2 mmol) via the general reduction procedure. After flash chromatography (15% ethyl acetate in hexanes), the product was obtained as a yellow oil (445 mg, 88%).

1H NMR ($CDCl_3$, 400 MHz): δ 6.73 (s, 1H), 6.61-6.54 (m, 2H), 4.01-3.95 (m, 1H), 3.75 (s, 3H), 3.59 (br s, 1H), 3.12 (dd, J = 8.4, 15.5 Hz, 1H), 2.63 (dd, J = 7.9, 15.5 Hz, 1H), 1.30 (d, J = 6.2 Hz, 3H);

^{13}C NMR ($CDCl_3$, 125 MHz): δ 144.7, 130.7, 130.5, 112.0, 111.6, 109.8, 55.9, 55.7, 38.2, 22.2.

5-Bromo-2-methylindoline [99847-70-8]. This indoline was prepared from 5-bromo-2-methylindole (500 mg, 3.11 mmol) and sodium cyanoborohydride (450 mg, 7.2 mmol) via the general reduction procedure. After flash chromatography (15% ethyl acetate in hexanes), the product was obtained as a clear, colorless oil (454 mg, 90%).

1H NMR ($CDCl_3$, 400 MHz): δ 7.17 (s, 1H), 7.10 (d, J = 8.2 Hz, 1H), 6.46 (d, J = 8.2 Hz, 1H), 4.05-3.96 (m, 1H), 3.78 (br s, 1H), 3.13 (dd, J = 8.6, 15.6 Hz, 1H), 2.63 (dd, J = 7.6, 15.7 Hz, 1H), 1.28 (d, J = 6.2 Hz, 3H);

^{13}C NMR ($CDCl_3$, 125 MHz): δ 149.7, 131.3, 129.8, 127.6, 110.4, 110.1, 55.6, 37.5, 22.1.

III. Kinetic Resolutions

General Procedure. In a nitrogen-filled glovebox, an oven-dried 40-mL vial was charged with catalyst (+)-1 (41 mg, 0.050 mmol), powdered LiBr (130 mg, 1.50 mmol), and the indoline (1.00 mmol). Toluene (8.5 mL) was added, resulting in a light purple, heterogeneous mixture. A solution of 18-crown-6 (0.375 M in toluene; 2.0 mL, 0.75 mmol) was added slowly to the stirred suspension over the course of 1 min, during which time the mixture turned darker. A solution of the acylating agent (0.325 M in toluene; 2.0 mL, 0.65 mmol) was then added. The vial was capped with a rubber septum, removed from the glovebox, and placed into a cold bath. After the indicated time had elapsed, the remaining acylating agent was quenched by the addition of propylamine (0.65 mmol), and the reaction mixture was concentrated and purified by flash chromatography.

The second runs were conducted with catalyst (–)-1.

The ee values were determined on Chirasil® Dex-CB or Daicel Chiralpak® columns.

The catalyst can be recovered in good yield. For an example, see the kinetic resolution of 2-propylindoline (Table 2, entry 2).

The absolute stereochemistry of the indolines resolved in entries 1, 4, and 8 of Table 2 was assigned by comparison with compounds reported in the literature. All other assignments were made by analogy.

Note on reaction times: As indicated in Table 1 (entry 3 vs. entry 12), shorter reaction times can be employed, at the expense of lower selectivity.

(–)-(S)-2-Methylindoline (Table 2, entry 1). This compound was resolved according to the General Procedure using 133 mg (1.00 mmol) of racemic indoline (0 °C for 5 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (5% → 20% ethyl acetate in hexanes) furnished the indoline and the N-acylated indoline.

Recovered indoline: 56.8 mg (43% yield; 94% ee); $[\alpha]_{\text{D}}^{23} = -11$ ($c = 0.11$, benzene) {lit.⁶ $[\alpha]_{\text{D}}^{20} = +11$ ($c = 2.0$, benzene); (*R*)}. The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 11.1 min (major) and 9.6 min (minor).

Acylated product: 88.5 mg (51% yield; 76% ee; white, crystalline solid); $[\alpha]_{\text{D}}^{23} = -45$ ($c = 0.65$, benzene). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 23.7 min (major) and 19.2 min (minor).

Calculated conversion = 55%; **s = 25**.

Second run: conversion = 55%; **s** = **24** (indoline: 53.5 mg, 40% yield, 92% ee; *N*-acyl indoline: 90.4 mg, 52% yield, 77% ee).

2-Propylindoline (Table 2, entry 2). This compound was resolved according to the General Procedure using 161 mg (1.00 mmol) of racemic indoline (0 °C for 10 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (10% → 20% ethyl acetate in hexanes) furnished the indoline and the *N*-acylated indoline.

Catalyst **1** was recovered by flushing the initial short pad of silica gel with a solution of diethylamine/isopropanol/hexanes (10:10:80) and recovering the purple fractions, which contain the catalyst. Purification by flash chromatography (3 column volumes of CH₂Cl₂, then diethylamine/methyl *tert*-butyl ether/hexanes, 2.5:10:87.5) furnished the catalyst in ~80% yield (Run 1: 33.5 mg, 81%; Run 2: 35.3 mg, 85%).

Recovered indoline: 76.0 mg (47% yield; 82% ee); [α]_D²³ = +39 (*c* = 0.56, CHCl₃). The ee was determined by HPLC using a Daicel Chiralpak OJ-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 10.8 min (major) and 17.7 min (minor).

Acylated product: 97.2 mg (48% yield; 82% ee; white, crystalline solid); [α]_D²³ = -79 (*c* = 0.32, CHCl₃). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 21.5 min (major) and 15.3 min (minor).

Calculated conversion = 50%; **s** = **25**.

Second run: conversion = 53%; **s** = **28** (indoline: 59.0 mg, 37% yield, 90% ee; *N*-acyl indoline: 109 mg, 53% yield, 81% ee).

2-Phenethylindoline (Table 2, entry 3). This compound was resolved according to the General Procedure using 223 mg (1.00 mmol) of racemic indoline (r.t. for 14 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (5% → 20% ethyl acetate in hexanes) furnished the indoline and the *N*-acylated indoline.

Recovered indoline: 67.7 mg (30% yield; 98% ee); [α]_D²³ = +12 (*c* = 0.17, CHCl₃). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 24.9 min (major) and 17.0 min (minor).

Acylated product: 148 mg (56% yield; 64% ee; clear, colorless oil); [α]_D²³ = -46 (*c* = 0.38, CHCl₃). The ee was determined by HPLC using a Daicel Chiralpak AS-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 26.9 min (major) and 35.9 min (minor).

Calculated conversion = 60%; **s** = **20**.

Second run: conversion = 54%; **s** = **17** (indoline: 63.5 mg, 28% yield, 85% ee; *N*-acyl indoline: 135 mg, 51% yield, 73% ee).

2-((*tert*-Butyldimethylsilyloxy)methyl)indoline (Table 2, entry 4). This compound was resolved according to the General Procedure using 264 mg (1.00 mmol) of racemic indoline (10 °C for 19 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (5% → 15% ethyl acetate in hexanes) furnished the indoline and the N-acylated indoline.

Recovered indoline: 85.5 mg (32% yield; 84% ee); $[\alpha]_D^{23} = -43$ ($c = 0.20$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 0.5% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 12.0 min (major) and 11.0 min (minor).

Acylated product: 159 mg (52% yield; 62% ee; clear, colorless oil); $[\alpha]_D^{23} = -34$ ($c = 0.30$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 13.8 min (major) and 10.2 min (minor).

Calculated conversion = 57%; **s** = 11.

Second run: conversion = 56%; **s** = 17 (indoline: 97.1 mg, 37% yield, 90% ee; N-acyl indoline: 160 mg, 52% yield, 71% ee).

(-)-(R)-2-Hydroxymethylindoline [77122-21-5]. This indoline was obtained by stirring resolved 2-((*tert*-butyldimethylsilyloxy)methyl)indoline (263 mg, 1.00 mmol) with tetrabutylammonium fluoride (1.0 M in THF; 0.65 mL, 0.65 mmol) in THF (5 mL) at r.t. for 30 min. The mixture was then concentrated and purified by flash chromatography (10% isopropanol in hexanes), which afforded the product as a clear, colorless oil (45.2 mg, 93%).

$[\alpha]_D^{23} = -44$ ($c = 0.31$, ethanol) {lit.⁷ $[\alpha]_D = +53.6$ ($c = 0.89$, ethanol); (S)}.

***cis*-5,6,7,8,8a,9-Hexahydro-4bH-carbazole (Table 2, entry 5).** This compound was resolved according to the General Procedure using 173 mg (1.00 mmol) of racemic indoline (0 °C for 6 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (5% → 20% ethyl acetate in hexanes) furnished the indoline and the N-acylated indoline.

Recovered indoline: 61.6 mg (36% yield; 84% ee); $[\alpha]_D^{23} = -20$ ($c = 0.25$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 10.3 min (major) and 9.5 min (minor).

Acylated product: 108 mg (49% yield; 53% ee; white, crystalline solid); $[\alpha]_D^{23} = -41$ ($c = 0.72$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H

column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 25.6 min (major) and 20.6 min (minor).

Calculated conversion = 58%; **s** = 10.2

Second run: conversion = 64%; **s** = 9.3 (indoline: 53.6 mg, 31% yield, 91% ee; *N*-acyl indoline: 117 mg, 54% yield, 52% ee).

***cis*-5,5a,6,7,8,9,10,10a-Octahydrocyclohepta[b]indole (Table 2, entry 6).** This compound was resolved according to the General Procedure using 187 mg (1.00 mmol) of racemic indoline (0 °C for 10 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (10% → 20% ethyl acetate in hexanes) furnished the indoline and the *N*-acylated indoline.

Recovered indoline: 89.3 mg (48% yield; 86% ee); $[\alpha]_{\text{D}}^{23} = -30$ ($c = 0.52$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 13.0 min (major) and 10.2 min (minor).

Acylated product: 110 mg (48% yield; 80% ee; white, crystalline solid); $[\alpha]_{\text{D}}^{23} = -93$ ($c = 0.48$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 28.2 min (major) and 18.0 min (minor).

Calculated conversion = 52%; **s** = 24.

Second run: conversion = 51%; **s** = 39 (indoline: 85.6 mg, 46% yield, 91% ee; *N*-acyl indoline: 104 mg, 45% yield, 85% ee).

***cis*-2,3-Dimethylindoline (Table 2, entry 7).** This compound was resolved according to the General Procedure using 147 mg (1.00 mmol) of racemic indoline (0 °C for 7 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (10% → 20% ethyl acetate in hexanes) furnished the indoline and the *N*-acylated indoline.

Recovered indoline: 53.5 mg (36% yield; 91% ee); $[\alpha]_{\text{D}}^{23} = +41$ ($c = 0.060$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 0.5% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 34.6 min (major) and 30.5 min (minor).

Acylated product: 95.8 mg (51% yield; 74% ee; white, crystalline solid); $[\alpha]_{\text{D}}^{23} = -66$ ($c = 0.51$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 25.5 min (major) and 19.1 min (minor).

Calculated conversion = 55%; **s** = 21.

Second run: conversion = 55%; **s** = 16 (indoline: 51.7 mg, 35% yield, 85% ee; *N*-acyl indoline: 87.3 mg, 46% yield, 71% ee).

(-)-(2*S*,3*R*)-trans-2,3-Dimethylindoline (Table 2, entry 8). This compound was resolved according to the General Procedure using 147 mg (1.00 mmol) of racemic indoline (r.t. for 20 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (7% → 15% ethyl acetate in hexanes) furnished the indoline and the *N*-acylated indoline.

Recovered indoline: 37.2 mg (25% yield; 94% ee); $[\alpha]_{\text{D}}^{23} = -41$ ($c = 0.13$, CCl_4) {lit.⁸ $[\alpha]_{\text{D}}^{25} = -52.6$ ($c = 1.91$, CCl_4); (2*S*,3*R*)}. The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 0.5% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 19.3 min (major) and 17.8 min (minor).

Acylated product: 93.1 mg (49% yield; 53% ee; yellow oil); $[\alpha]_{\text{D}}^{23} = -2.0$ ($c = 0.26$, CCl_4). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 12.4 min (major) and 11.3 min (minor).

Calculated conversion = 64%; **s = 10.8**.

Second run: conversion = 65%; **s = 8.2** (indoline: 38.4 mg, 26% yield, 90% ee; *N*-acyl indoline: 95.3 mg, 50% yield, 48% ee).

cis-Ethyl 2-(5-methoxy-2-methylindolin-3-yl)acetate (Table 2, entry 9). This compound was resolved according to the General Procedure using 249 mg (1.00 mmol) of racemic indoline (−10 °C for 4 days). After quenching, the reaction mixture was passed through a short pad of silica gel (20% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (15% → 40% ethyl acetate in hexanes) furnished the indoline and the *N*-acylated indoline.

Recovered indoline: 80.4 mg (32% yield; 85% ee); $[\alpha]_{\text{D}}^{23} = +9.6$ ($c = 0.27$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 10% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 17.1 min (major) and 13.5 min (minor).

Acylated product: 159 mg (55% yield; 75% ee; white, crystalline solid); $[\alpha]_{\text{D}}^{23} = -34$ ($c = 0.32$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 10% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 23.6 min (major) and 17.5 min (minor).

Calculated conversion = 53%; **s = 18**.

Second run: conversion = 58%; **s = 19** (indoline: 97.8 mg, 39% yield, 95% ee; *N*-acyl indoline: 142 mg, 49% yield, 68% ee).

5-Methoxy-2-methylindoline (Table 2, entry 10). This compound was resolved according to the General Procedure using 163 mg (1.00 mmol) of racemic indoline (−10 °C for 2.5 days). After quenching, the reaction mixture was passed through a short pad

of silica gel (20% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (15% → 30% ethyl acetate in hexanes) furnished the indoline and the N-acylated indoline.

Recovered indoline: 57.3 mg (35% yield; 92% ee); $[\alpha]_D^{23} = +16$ ($c = 0.17$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 7% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 15.6 min (major) and 8.3 min (minor).

Acylated product: 106 mg (52% yield; 60% ee; clear, colorless oil); $[\alpha]_D^{23} = -28$ ($c = 0.70$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 7% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 19.0 min (major) and 17.0 min (minor).

Calculated conversion = 60%; **s = 12**.

Second run: conversion = 60%; **s = 14** (indoline: 53.7 mg, 33% yield, 92% ee; N-acyl indoline: 111 mg, 54% yield, 63% ee).

5-Bromo-2-methylindoline (Table 2, entry 11). This compound was resolved according to the General Procedure using 212 mg (1.00 mmol) of racemic indoline (r.t. for 23 days). After quenching, the reaction mixture was passed through a short pad of silica gel (10% isopropanol in hexanes) to remove the catalyst and LiBr. Purification by flash chromatography (5% → 20% ethyl acetate in hexanes) furnished the indoline and the N-acylated indoline.

Recovered indoline: 63.0 mg (30% yield; 83% ee); $[\alpha]_D^{23} = -11$ ($c = 0.18$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 16.4 min (major) and 9.6 min (minor).

Acylated product: 121 mg (47% yield; 64% ee; clear, colorless oil); $[\alpha]_D^{23} = +45$ ($c = 0.12$, CHCl_3). The ee was determined by HPLC using a Daicel Chiralpak OD-H column (eluent: 1% isopropanol in hexanes; flow rate: 1.0 mL/min). Retention times: 24.3 min (major) and 21.7 min (minor).

Calculated conversion = 57%; **s = 11**.

Second run: conversion = 60%; **s = 11** (indoline: 55.8 mg, 26% yield, 90% ee; N-acyl indoline: 120 mg, 47% yield, 59% ee).

IV. Absolute Configuration of (+)-1

General. Low-temperature diffraction data were collected on a Siemens Platform three-circle diffractometer coupled to a Bruker-AXS Apex CCD detector with graphite monochromated Mo K α radiation ($\lambda = 0.71073\text{\AA}$), performing φ - and ω -scans. Raw data frames were integrated using the Bruker program SAINT+ for NT version 6.01. All structures were solved by direct methods using SHELXS and refined against F^2 on all data by full-matrix least squares with SHELXL-97 (Sheldrick, G. M. *SHELXL 97*, Universität Göttingen, Göttingen, Germany, 1997). SADABS absorption correction was performed. All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were included in the model at geometrically calculated positions and refined using a riding model. The isotropic displacement parameters of the hydrogen atoms were fixed to 1.2 times the U value of the atoms they are linked to (1.5 times for methyl groups).

Experimental. A dark-purple needle (grown from a mixture of catalyst, acylating agent, 18-crown-6, and LiBr) of dimensions $0.25 \times 0.20 \times 0.10\text{ mm}^3$ was mounted under STP and transferred to a Siemens Platform three-circle diffractometer equipped with a cold stream of N₂ gas. The data that were collected (185924 total reflections, 26781 unique, $R_{\text{int}} = 0.058$) had the following Miller index ranges: (–20 to 20 in h , –27 to 27 in k , and –46 to 46 in l). The structure was solved in the monoclinic space group $P2(1)2(1)2(1)$, $a = 14.9080(4)\text{ \AA}$, $b = 19.7937(6)\text{ \AA}$, $c = 33.7362(10)\text{ \AA}$, $\alpha = 90^\circ$; $\beta = 90^\circ$; $\gamma = 90^\circ$, and refined using standard difference Fourier techniques. Final, full-matrix least squares refinement (26781 data for 1191 parameters) on F^2 yielded residuals of R_1 and wR_2 of 0.0624 and 0.1734 for data $I > 2\sigma(I)$, and 0.0822 and 0.1872, respectively, for all data. Residual electron density amounted to a maximum of $1.001\text{ e/\AA}^3$ and a minimum of $-1.673\text{ e/\AA}^3$. The absolute structure (Flack) parameter for the correct enantiomer is 0.002(8), thus establishing the absolute stereochemistry.

Figure 1. A portion of the unit cell, focusing on (+)-1.

Figure 2. The unit cell: 2 (+)-1•2 LiBr•toluene.

Table 1. Crystal data and structure refinement for 06067.

Identification code	06067	
Empirical formula	C ₁₂ H ₁₂ Br ₂ Fe ₂ Li ₂ N ₄	
Formula weight	1919.73	
Temperature	100(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2(1)2(1)2(1)	
Unit cell dimensions	a = 14.9080(4) Å	$\alpha = 90^\circ$.
	b = 19.7937(6) Å	$\beta = 90^\circ$.
	c = 33.7362(10) Å	$\gamma = 90^\circ$.
Volume	9955.0(5) Å ³	
Z	8	
Density (calculated)	2.476 Mg/m ³	
Absorption coefficient	2.290 mm ⁻¹	
F(000)	7536	
Crystal size	0.25 x 0.20 x 0.10 mm ³	
Theta range for data collection	1.19 to 29.13°.	
Index ranges	-20 ≤ h ≤ 20, -27 ≤ k ≤ 27, -46 ≤ l ≤ 46	
Reflections collected	185924	
Independent reflections	26781 [R(int) = 0.0578]	
Completeness to theta = 29.13°	100.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.8034 and 0.5983	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	26781 / 1362 / 1191	
Goodness-of-fit on F ²	1.041	
Final R indices [I > 2σ(I)]	R1 = 0.0624, wR2 = 0.1734	
R indices (all data)	R1 = 0.0822, wR2 = 0.1872	
Absolute structure parameter	0.002(8)	
Largest diff. peak and hole	1.001 and -1.673 e.Å ⁻³	

V. References

- (1) Kagan, H. B.; Fiaud, J. C. *Top. Stereochem.* **1988**, *18*, 249–330.
- (2) Ruble, J. C.; Fu, G. C. *J. Am. Chem. Soc.* **1998**, *120*, 11532–11533.
- (3) Gribble, G. W.; Hoffman, J. H. *Synthesis* **1977**, 859–860.
- (4) Kuyper, L. F.; Baccanari, D. P.; Jones, M. L.; Hunter, R. N.; Tansik, R. L.; Joyner, S. S.; Boytos, C. M.; Rudolph, S. K.; Knick, V.; Wilson, H. R.; Caddell, J. M.; Friedman, H. S.; Comley, J. C. W.; Stables, J. N. *J. Med. Chem.* **1996**, *39*, 892–903.
- (5) Lanzilotti, A. E.; Littell, R.; Fanshawe, W. J.; McKenzie, T. C.; Lovell, F. M. *J. Org. Chem.* **1979**, *44*, 4809–4813.
- (6) Krasnov, V. P.; Levit, G. L.; Bukrina, I. M.; Andreeva, I. N.; Sadretdinova, L. S.; Korolyova, M. A.; Kodess, M. I.; Charushin, V. N.; Chupakhin, O. N. *Tetrahedron: Asymmetry* **2003**, *14*, 1985–1988.
- (7) Pasquier, C.; Naili, S.; Mortreux, A.; Agbossou, F.; Péliniski, L.; Brocard, J.; Eilers, J.; Reiners, I.; Peper, V.; Martens, J. *Organometallics* **2000**, *19*, 5723–5732.
- (8) Scholl, B.; Hansen, H.-J. *Helv. Chim. Acta* **1980**, *63*, 1823–1832.

STANDARD 1H OBSERVE

Pulse Sequence: s2pu1
 Solvent: CDCl3
 Ambient temperature
 File: 4232p.csm
 INOVA-500 "zippy"

PULSE SEQUENCE
 Relax. delay 1.000 sec
 Pulse 34.1 degrees
 Acq. time 1.995 sec
 Width 4506.5 Hz
 16 repetitions
 OBSERVE H1, 300.0986331 MHz
 DATA PROCESSING
 FT size 32768
 Total time 0 min, 48 sec

STANDARD 1H OBSERVE

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 File: 6237
 INOVA-500 "zippy"

PULSE SEQUENCE
 Relax. delay 1.000 sec
 Pulse 34.1 degrees
 Acq. time 1.995 sec
 Width 4506.5 Hz
 16 repetitions
 OBSERVE H1, 300.0986325 MHz
 DATA PROCESSING
 FT size 32768
 Total time 0 min, 48 sec

Current Data Parameters

NAME 4261sm
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20060720
Time_ 19.13
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16

Current Data Parameters

NAME 4208
EXPNO 1
PROCNO 1

F2 - Acquisition Parameter

Date_ 20060414
Time 16.32
INSTRUM spect
PROBHD 5 mm QNP
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 8

STANDARD 1H OBSERVE

Pulse Sequence: s2pu1
Solvent: CDCl3
Ambient temperature
Mercury-300 "mrhrt"

Relax. delay 1.000 sec
Pulse 34.1 degrees
Acq. time 1.995 sec
Width 4506.5 Hz
16 repetitions
OBSERVE H1, 300.0986331 MHz
DATA PROCESSING
FT size 32768
Total time 0 min, 0 sec

STANDARD PROTON PARAMETERS

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 File: 4228 "zippy"
 INOVA-500
 PULSE SEQUENCE
 Relax. delay 1.000 sec
 Pulse 34.1 degrees
 Acq. time 1.995 sec
 Width 4506.5 Hz
 16 repetitions
 OBSERVE H1, 300.0986320 MHz
 DATA PROCESSING
 FT size 32768
 Total time 0 min, 48 sec

STANDARD PROTON PARAMETERS

Pulse Sequence: s2pu1
 Solvent: CDCl3
 Ambient temperature
 INOVA-500 "bujlwinkle"
 Relax. delay 1.000 sec
 Pulse 89.0 degrees
 Acq. time 3.001 sec
 Width 10504.2 Hz
 8 repetitions
 OBSERVE H1 499.7446543 MHz
 DATA PROCESSING
 FT size 131072
 Total time 0 min, 32 sec

STANDARD PROTON PARAMETERS

Pulse Sequence: s2pu1

Solvent: CDCl3

Ambient temperature

INOVA-500 "bottle"inkle"

Relax. delay 2.000 sec

Pulse 99.0 degrees

Acq. time 3.001 sec

Width 10504.2 Hz

4 repetitions

OBSERVE H1, 499.746521 MHz

DATA PROCESSING

FT size 131072

Total time 0 min, 20 sec

Current Data Parameters
NAME 4251sm
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060702
Time 17.02
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16

Current Data Parameters

NAME 4130-pMeO
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20060128
Time 15.02
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 8

Current Data Parameters

NAME 4133
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20060130
Time 21.13
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 8