Supporting information for

**Kinetics and Mechanistic Aspects of the Heck Reaction Promoted by a CN-
Palladacycle**

*Crestina S. Consorti, Fabrício R. Flores and Jairton Dupont**

Laboratory of Molecular Catalysis – Institute of Chemistry – UFRGS Avenida Bento
Gonçalves, 9500 Porto Alegre 91501-970 RS Brazil

Calibration curve for ART-IR

Figure S1. Calibration curve obtained at 50 °C for the Ph-I stretching vibration at 735 cm⁻¹.
 $Y = -0.1 + 257.7X$ ($R = 0.99$).

Reaction orders determined by the initial velocity method

Table S1. Initial reagents concentrations and initial reaction rates found in the Heck reaction of iodobenzene and *n*-butylacrylate promoted by palladacycle **1** at 80 °C.

NaOAc (mol L ⁻¹)	NBu ₄ Br (mol L ⁻¹)	Acrylate (mol L ⁻¹)	PhI (mol L ⁻¹)	(1) (mmol L ⁻¹)	$-d[\text{PhI}]/dt$ (mol L ⁻¹ h ⁻¹)
0.256	0.036	0.217	0.182	0.018	0.12 ± 0.02
0.255	0.036	0.217	0.182	0.036	0.26 ± 0.04
0.253	0.035	0.215	0.180	0.090	0.49 ± 0.06
0.250	0.035	0.213	0.178	0.178	1.08 ± 0.09
0.073	0.036	0.216	0.181	0.181	1.12 ± 0.10
0.109	0.035	0.215	0.180	0.181	1.13 ± 0.05
0.145	0.035	0.215	0.180	0.180	1.09 ± 0.08
0.252	0.017	0.214	0.179	0.179	1.01 ± 0.10
0.248	0.070	0.211	0.176	0.178	0.95 ± 0.08
0.242	0.137	0.206	0.173	0.173	0.99 ± 0.05
0.257	0.036	0.036	0.183	0.183	0.23 ± 0.04
0.256	0.036	0.054	0.183	0.183	0.35 ± 0.03
0.256	0.036	0.072	0.182	0.182	0.48 ± 0.05
0.253	0.035	0.143	0.180	0.180	1.01 ± 0.08
0.254	0.036	0.216	0.036	0.182	0.60 ± 0.09
0.254	0.036	0.216	0.054	0.181	0.91 ± 0.09
0.253	0.035	0.216	0.072	0.181	0.96 ± 0.05
0.252	0.035	0.215	0.090	0.180	0.98 ± 0.07
0.251	0.035	0.214	0.143	0.179	1.02 ± 0.08
0.250	0.035	0.213	0.178	0.178	1.08 ± 0.09

Table S2. Initial reaction rate found in the Heck reaction of iodobenzene and *n*-butylacrylate promoted by palladacycle **1** at different temperatures. NaOAc (0.250 mol L⁻¹), NBu₄Br (0.035 mol L⁻¹), *n*-butylacrylate (2.13 mol L⁻¹), PhI (0.178 mol L⁻¹), **1** (0.178 mmol L⁻¹).

T (K)	$-d[PhI]/dt$ (mol L ⁻¹ h ⁻¹)
325.15	0.11 ± 0.02
332.15	0.16 ± 0.02
345.15	0.41 ± 0.05
355.15	1.09 ± 0.08
374.15	3.13 ± 0.29

Figure S2. Plot of the logarithm of initial reaction rate versus logarithm of initial concentrations of palladacycle **1** in the Heck reaction of iodobenzene and *n*-butylacrylate at 80 °C. $Y = (8.6 \pm 0.67) + (0.98 \pm 0.072)X$ ($R = 0.99$).

Figure S3. Plot of the logarithm of initial reaction rate versus logarithm of initial concentrations of NBu_4Br in the Heck reaction of iodobenzene and *n*-butylacrylate at 80 °C.
 $Y = (-0.057 \pm 0.12) + (-0.022 \pm 0.041)X$

Figure S4. Plot of the logarithm of initial reaction rate versus logarithm of initial concentrations of NaOAc in the Heck reaction of iodobenzene and *n*-butylacrylate at 80 °C.
 $Y = (-0.0014 \pm 0.21) + (-0.044 \pm 0.099)X$.

Figure S5. Plot of the logarithm of initial reaction rate versus logarithm of initial concentrations of *n*-butylacrylate in the Heck reaction of iodobenzene and *n*-butylacrylate at 80 °C. $Y = (2.1 \pm 0.26) + (1.1 \pm 0.10)X$.

Figure S6. Plot of the logarithm of initial reaction rate versus logarithm of initial concentrations of iodobenzene in the Heck reaction of iodobenzene and *n*-butylacrylate at 80 °C.

Deduction of rate equation using the steady state approximation

$$\frac{d[1]}{dt} = k_3[3] - k_1[1][PhI] = 0 \quad (1)$$

$$\frac{d[3]}{dt} = -k_3[3] + k_2[2][Acr] - k_{-2}[3] = 0 \quad (2)$$

$$\frac{d[2]}{dt} = k_1[1][PhI] - k_2[2][Acr] + k_{-2}[3] = 0 \quad (3)$$

Assuming that $[Pd] = [1] + [2] + [3]$, the following expression may be written:

$$[Pd] = [3] \left(\frac{k_3}{k_1[PhI]} + \frac{k_3 + k_{-2}}{k_2[Acr]} + 1 \right) \quad (4)$$

Or:

$$k_1k_2[PhI][Acr][Pd] = [3](k_2k_3[Acr] + k_1k_3[PhI] + k_1k_{-2}[PhI] + k_1k_2[PhI][Acr]) \quad (5)$$

The rate of product formation is equal to the rate of substrate consumption:

$$\frac{d[P]}{dt} = -\frac{d[PhI]}{dt} = k_3[3] \quad (6)$$

Combining equation (6) and (5):

$$-d[PhI]/dt = \frac{k_1k_2k_3[PhI][Acr][Pd]}{k_2k_3[Acr] + k_1k_2[PhI][Acr] + k_1k_3[PhI] + k_1k_{-2}[PhI]} \quad (7)$$

Deduction of rate equation of two competing halo-arenes

$$\frac{d[1]}{dt} = k_3[3] + k_3'[5] - k_1[1][PhI] - k_1^1[1][RPhI] = 0 \quad (8)$$

$$\frac{d[2]}{dt} = k_1[1][PhI] - k_2[2][Acr] + k_{-2}[3] = 0 \quad (9)$$

$$\frac{d[3]}{dt} = -k_3[3] + k_2[2][Acr] - k_{-2}[3] = 0 \quad (10)$$

$$\frac{d[4]}{dt} = k_1'[1][RPhI] - k_2'[4][Acr] + k_{-2}'[5] = 0 \quad (11)$$

$$\frac{d[5]}{dt} = -k_3'[5] + k_2'[4][Acr] - k_{-2}'[5] = 0 \quad (12)$$

The following relations can be found:

From (10):

$$[2] = \frac{(k_3 + k_{-2})}{k_2[Acr]} [3] \quad (13)$$

From (12):

$$[4] = \frac{(k_3' + k_{-2}')}{k_2'[Acr]} [5] \quad (14)$$

Combining equations (13) and (9):

$$k_1[1][PhI] - k_2 \frac{(k_3 + k_{-2})}{k_2[Acr]} [3][Acr] + k_{-2}[3] = 0 \quad \text{or} \quad [3] = \frac{k_1[PhI][1]}{k_3} \quad (15)$$

Combining equations (14) and (9):

$$k_1'[1][RPhI] - k_2' \frac{(k_3' + k_{-2}')}{k_2'[Acr]} [5][Acr] + k_{-2}'[5] = 0 \quad \text{or} \quad [5] = \frac{k_1'[RPhI][1]}{k_3'} \quad (16)$$

Combining equations (13) and (15):

$$[2] = \frac{(k_3 + k_{-2})}{k_2[Acr]} [3] \quad \text{or} \quad [2] = \frac{(k_3 + k_{-2})}{k_2[Acr]} \frac{k_1[PhI][1]}{k_3} \quad (17)$$

Combining equations (14) and (16):

$$[4] = \frac{(k_3' + k_{-2}')}{k_2'[Acr]} [5] \quad \text{or} \quad [4] = \frac{(k_3' + k_{-2}')}{k_2'[Acr]} \frac{k_1'[RPhI][1]}{k_3'} \quad (18)$$

Assuming that $[Pd] = [1] + [2] + [3] + [4] + [5]$, the following expression may be written:

$$[Pd] = [1] \left(1 + \frac{(k_3 + k_{-2})}{k_2[Acr]} \frac{k_1[PhI]}{k_3} + \frac{k_1[PhI]}{k_3} + \frac{(k_3' + k_{-2}')}{k_2'[Acr]} \frac{k_1'[RPhI]}{k_3'} + \frac{k_1'[RPhI]}{k_3'} \right) \quad (19)$$

$$k_2 k_2' k_3 k_3' [Acr] [Pd] = [1] (k_2 k_2' k_3 k_3' [Acr] + k_1 k_2' k_3' (k_3 + k_{-2}) [PhI] + k_1 k_2 k_3' [Acr] [PhI] + k_1' k_2 k_3 (k_3' + k_{-2}') [RPhI] + k_1' k_2 k_2' k_3 [Acr] [RPhI]) \quad (20)$$

Or:

$$k_2 k_2' k_3 k_3' [Acr] [Pd] = [1] a \quad (21)$$

$$a = k_2 k_3 k_2' k_3' [Acr] + k_1 k_2' k_3' [PhI] (k_{-2} + k_3 + k_2 [Acr]) + k_1' k_2 k_3 [RPhI] (k_{-2}' + k_3' + k_2' [Acr]) \quad (22)$$

The rates of consumption of competing substrates are:

$$\frac{d[PhI]}{dt} = -k_1 [PhI] [1] \quad (23)$$

$$\frac{d[RPhI]}{dt} = -k_1' [RPhI] [1] \quad (24)$$

Combining equation (20) with the above expressions, the reaction rate law for competing substrates can be written:

$$-d[PhI]/dt = \frac{k_1 k_2 k_3 k_2' k_3' [PhI] [Acr] [Pd]}{a} \quad (25)$$

$$-d[RPhI]/dt = \frac{k_1' k_2' k_3' k_2 k_3 [RPhI] [Acr] [Pd]}{a} \quad (26)$$

$$a = k_2 k_3 k_2' k_3' [Acr] + k_1 k_2' k_3' [PhI] (k_{-2} + k_3 + k_2 [Acr]) + k_1' k_2 k_3 [RPhI] (k_{-2}' + k_3' + k_2' [Acr]) \quad (27)$$

The relative reactivity of competing substrates can be obtained by dividing (25) and (26):

$$\frac{d[RPhI]/dt}{d[PhI]/dt} = \frac{k_1'[RPhI]}{k_1[PhI]} \quad (28)$$

Script used in DynaFit for fitting the experimental data

Fitting the time course of PhI
Heck reaction

[task]

```
data = progress
task = fit
model = PhI
```

[mechanism]

```
PdPhac ---> pda + P      : k1
pda + PhI ---> PdPhI      : k2
PdPhI + acr <===> PdPhac : k k-
```

[constants]

k2 = 200 ?, k = 20 ?, k- = 100 ?, k1 = 200 ?

[concentrations]

pda = 0.178, acr = 213.41, PhI = 178.8

[progress]

```
directory    ./examples/heck/data
extension    txt
file h100
concentration PhI = 107.8 | response PhI = 1
file h101
concentration PhI = 143.5 | response PhI = 1
file h64
concentration PhI = 178.8 | response PhI = 1
file h76
concentration acr = 140.7 | response PhI = 1
file h63
concentration pda = 0.08 | response PhI = 1
```

file h61
 concentration pda = 0.018 | response PhI = 1
 file h75
 concentration acr = 60.5 | response PhI = 1
 file h79
 concentration PhI = 72.7 | response PhI = 1

[output]

directory ./examples/heck/outputheck

[settings]

<Filter> | SetSigZero = No | TMax = 200
 <Simulate> | Interpolate = yes | Increment = 2
 <output> | WriteTABtFiles = Yes

[end]

;

Hammett plot obtained with excess bromo-arenes concentration

Figure S7. Hammett correlation for the competitive Heck reaction between bromo arenes and *n*-butylacrylate at 150 °C in DMA promoted by Herrmann palladacycle **2** using the σ_0 constants. Competitive reactions performed with excess *n*-butylacrylate (■) or excess bromo-arene (○). $\rho = 2.4$.

Figure S8. Hammett correlation for the competitive Heck reaction between bromo arenes and *n*-butylacrylate at 150 °C in DMA promoted by palladacycle **1** using the σ_0 constants. Competitive reactions performed with excess *n*-butylacrylate (■) or excess bromo-arene (○). $\rho = 2.5$.