Supporting Information

Lewis Acid Mediated [2,3]-Sigmatropic Rearrangement of Allylic α -Amino Amides.

Jan Blid,[†] Peter Brandt,[‡] Peter Somfai^{,†}*

Department of Chemistry, Organic Chemistry, Royal Institute of Technology, SE-100 44

Stockholm, Sweden, and Department of Structural Chemistry, Biovitrum AB, SE-112 76

Stockholm, Sweden

somfai@kth.se

Content

S2:	Synthesis of 2
S3-S9:	Experimental data for 2 , 7 , 8 , 12 , 14b-e , 15b-e and 16b-e
S10-S40:	NMR spectra for 2 , 3 , 7 , 8 , 9a , 9b , 10a , 10b , 11 , 12 , 14b-e , 15b-e and 16b-e
S41:	References

* Corresponding author: Phone: (+46)-8-790 6960. Fax: (+46)-8-791 2333.

[†] Royal Institute of Technology

[‡] Department of Structural Chemistry, Biovitrum AB

Synthesis of allyl amines 2

Scheme S1^a

^a Reagents and conditions: a) RBr, K₂CO₃, MeCN, rt.¹ b) **12**, pyrrolidine, MgBr₂, THF, rt.²

Table S1^a

Entry	Substrate	R ¹	R ²	Yields [%]			
				14	15	16	2
1	b	Me	H	98	76	93	96
2	c	Ph	H	Quant.	88	91	93
3	d	H	Me	88	71	Quant.	97
4	e	Me	Me	98	90	88	94

^a Reagents and conditions: a) PPh₃, DIAD, THF, rt.;³ b) mercaptoacetic acid, Et₃N, CH₂Cl₂, rt.;³ c) methyl bromoacetate, K₂CO₃, MeCN, rt.;¹ d) pyrrolidine, MgBr₂, THF, rt.²

General Methods.

^1H and ^{13}C NMR spectra were recorded on a 400 MHz or 500 MHz spectrometer in CDCl_3 , CD_2Cl_2 , $\text{DMSO}-d_6$ or CD_3CN using the residual peak of CHCl_3 (^1H NMR δ 7.26, ^{13}C NMR δ 77.0), CH_2Cl_2 (^1H NMR δ 5.32, ^{13}C NMR δ 53.8), DMSO (^1H NMR δ 2.50, ^{13}C NMR δ 39.7), CH_3CN (^1H NMR δ 1.93, ^{13}C NMR δ 118.2, 1.3) or added TMS (δ 0.00) as internal standard. ^{11}B and ^{19}F NMR spectra were recorded on a 500 MHz spectrometer using $\text{BF}_3\cdot\text{OEt}_2$ or trifluorotoluene, respectively, as external standard. Chemical shifts are reported in the δ -scale with multiplicity (b=broad, s=singlet, d=doublet, t=triplet, q=quartet, qn=quintet, m=multiplet), integration and coupling constants (Hz). Analytical thin layer chromatography was performed on silica gel 60 F_{254} plates, the plates were visualized with UV light and phosphomolybdic acid/cerium sulfate staining reagent (ca 5 wt% in ethanol) or I_2/MeOH . Chromatography employed silica gel 60 (35-70 μm). Air and moisture sensitive reactions were carried out in flame-dried, septum-capped flasks under an atmospheric pressure of nitrogen. All liquid reagents were transferred via oven-dried syringes. THF, Et_2O and toluene were distilled from sodium-benzophenone ketyl before use and CH_2Cl_2 and Et_3N from CaH_2 .

1-[(*N*-Allyl-*N*-benzylamino)acetyl]-pyrrolidine (2a). To a stirred solution of **12** (5.37 g, 24.5 mmol) and MgBr_2 (9.11 g, 49.4 mmol) in THF (100 mL) was added pyrrolidine (8.5 mL, 102 mmol). After stirring for 22 h at rt the reaction mixture was cooled to 0 $^\circ\text{C}$, sat. NaHCO_3 (20 mL) at 0 $^\circ\text{C}$ was added and stirring was continued at rt for 30 min. The solvents were evaporated and the residue was diluted with H_2O (100 mL) and extracted with Et_2O (3 $\times$ 30 mL). The dried (K_2CO_3) organic solution was concentrated and chromatographed ($\text{Et}_2\text{O}:\text{MeCN}$ 1:0 $\rightarrow$ 0:1) to provide **2a** (5.0 g, 84%) as a colorless oil. ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.35-7.22 (m, 5H), 5.90 (ddt, 1H, J =17.1, 10.2, 6.6), 5.21 (d, 1H, J =17.1), 5.16 (d, 1H, J =10.2), 3.74 (s, 2H), 3.43 (t, 2H, J =6.7), 3.36 (t, 2H, J =6.7), 3.24 (d, 2H, J =6.6), 3.22 (s, 2H), 1.88 (m, 2H), 1.80 (m, 2H); ^{13}C NMR (CDCl_3 , 100 MHz)

δ_C 169.3, 139.0, 135.7, 129.2, 128.2, 127.0, 117.9, 58.2, 57.4, 56.3, 46.2, 45.7, 26.2, 24.1; IR (neat) 2974, 2875, 1641 cm^{-1} ; HRMS (EI+) calcd for $\text{C}_{16}\text{H}_{22}\text{N}_2\text{O}$ $[\text{M}]^+$: 258.1732, found: 258.1732.

1-[(*E*)-*N*-Benzyl-*N*-2-butenylamino]acetyl]-pyrrolidine (2b). As described for **2a**, starting from **16b** (0.56 g, 2.4 mmol). Pale yellow oil (0.63 g, 96%). ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.32 (d, 2H, $J=7.0$), 7.29 (t, 2H, $J=7.0$), 7.23 (t, 1H, $J=7.0$), 5.62 (dq, 1H, $J=15.2$, 6.3), 5.52 (dt, 1H, $J=15.2$, 6.6), 3.71 (s, 2H), 3.42 (t, 2H, $J=6.9$), 3.37 (t, 2H, $J=6.7$), 3.20 (s, 2H), 3.15 (d, 2H, $J=6.6$), 1.87 (m, 2H), 1.79 (m, 2H), 1.69 (dd, 3H, $J=6.3$, 1.2); ^{13}C NMR (CDCl_3 , 100 MHz) δ_C 169.4, 139.1, 129.14, 129.07, 128.12, 128.07, 126.9, 58.1, 56.5, 56.3, 46.1, 45.6, 26.2, 24.1, 17.8; IR (neat) 2968, 2875, 1643 cm^{-1} ; HRMS (EI+) calcd for $\text{C}_{17}\text{H}_{25}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$: 273.1967, found: 273.1967.

1-[(*E*)-*N*-Benzyl-*N*-(3-phenyl-2-propenyl)amino]acetyl]-pyrrolidine (2c). As described for **2a**, starting from **16c** (0.34 g, 1.2 mmol). Pale yellow oil (0.36 g, 93%). ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.38-7.20 (m, 10H), 6.55 (d, 1H, $J=15.8$), 6.30 (dt, 1H, $J=15.8$, 6.8), 3.81 (s, 2H), 3.43 (t, 2H, $J=6.9$), 3.42 (d, 2H, $J=6.8$), 3.34 (t, 2H, $J=6.7$), 3.27 (s, 2H), 1.86 (m, 2H), 1.78 (m, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ_C 169.2, 138.9, 137.0, 132.9, 129.2, 128.5, 128.2, 127.41, 127.39, 127.0, 126.3, 58.5, 56.7, 56.1, 46.1, 45.7, 26.2, 24.0; IR (neat) 3082, 3057, 2972, 2951, 1641 cm^{-1} ; HRMS (EI+) calcd for $\text{C}_{22}\text{H}_{27}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$: 335.2123, found: 335.2115.

1-[(*Z*)-*N*-Benzyl-*N*-2-butenylamino]acetyl]-pyrrolidine (2d). As described for **2a**, starting from **16d** (0.86 g, 3.7 mmol). Pale yellow oil (1.0 g, 97%). ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.34-7.21 (m, 5H), 5.65 (m, 1H), 5.54 (m, 1H), 3.72 (s, 2H), 3.42 (t, 2H, $J=6.8$), 3.39 (t, 2H, $J=6.8$), 3.24 (d, 2H, $J=6.9$), 3.22 (s, 2H), 1.87 (m, 2H), 1.79 (m, 2H), 1.61 (d, 3H, $J=6.8$); ^{13}C NMR (CDCl_3 , 100 MHz) δ_C 169.4, 139.1, 129.2, 128.1, 127.6, 127.4, 126.9, 58.4, 56.7, 50.5, 46.2, 45.7, 26.2, 24.1, 13.1; IR (neat) 2972, 2873, 1643 cm^{-1} ; HRMS (CI+) calcd for $\text{C}_{17}\text{H}_{25}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$: 273.1967, found: 273.1971.

1-[(*N*-Benzyl-*N*-(3-methyl-2-butenyl)amino)acetyl]-pyrrolidine (2e). As described for **2a**, starting from **16e** (0.26 g, 1.1 mmol). Pale yellow oil (0.28 g, 94%). ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.34-7.20 (m, 5H), 5.29 (t, 1H, $J=7.0$), 3.70 (s, 2H), 3.40 (t, 2H, $J=6.9$), 3.38 (t, 2H, $J=6.7$), 3.20 (s, 2H), 3.16 (d, 2H, $J=6.9$), 1.87 (m, 2H), 1.79 (m, 2H), 1.73 (s, 3H), 1.61 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ_{C} 169.5, 139.3, 135.7, 129.2, 128.1, 126.9, 121.5, 58.4, 56.8, 51.8, 46.2, 45.7, 26.2, 25.9, 24.1, 18.0; IR (neat) 3084, 3060, 2970, 2927, 1641 cm^{-1} ; HRMS (EI+) calcd for $\text{C}_{18}\text{H}_{27}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$: 287.2123, found: 287.2122.

***N*-Allyl-benzylglycine methyl ester (12).**¹ To a suspension of *N*-allyl benzylamine (1.10 g, 7.48 mmol) and K_2CO_3 (2.06 g, 14.9 mmol) in CH_3CN (20 mL) was added methyl bromoacetate (0.90 mL, 8.58 mmol) and the resultant solution stirred for 20 h. The reaction mixture was poured into sat. NaHCO_3 (30 mL) and extracted with CH_2Cl_2 . The combined organic phases were dried (MgSO_4) and concentrated. Purification by chromatography (pentane:EtOAc 6:1) provided **12** (1.45 g, 88%) as a colorless oil. ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.36-7.23 (m, 5H), 5.88 (ddt, 1H, $J=17.1$, 10.1, 6.5), 5.22 (dq, 1H, $J=17.1$, 1.9), 5.16 (ddt, 1H, $J=10.1$, 1.9, 1.2), 3.78 (s, 2H), 3.68 (s, 3H), 3.32 (s, 2H), 3.27 (dt, 2H, $J=6.5$, 1.2); ^{13}C NMR (CDCl_3 , 100 MHz) δ_{C} 171.9, 138.6, 135.5, 129.0, 128.3, 127.1, 118.0, 57.8, 56.9, 53.5, 51.3.

***N*-Allyl-benzylglycine *tert*-butyl ester (7).** As described for **12**. Colorless oil (1.2 g, 71%). ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.37 (d, 2H, $J=7.1$), 7.31 (t, 2H, $J=7.1$), 7.25 (app. t, 1H, $J=7.1$), 5.89 (ddt, 1H, $J=17.2$, 10.2, 6.5), 5.23 (ddt, 1H, $J=17.2$, 2.0, 1.2), 5.15 (ddt, 1H, $J=10.2$, 2.0, 1.2), 3.79 (s, 2H), 3.28 (dt, 2H, $J=6.5$, 1.2), 3.22 (s, 2H), 1.47 (s, 9H); ^{13}C NMR (CDCl_3 , 100 MHz) δ_{C} 170.7, 139.0, 135.8, 128.9, 128.2, 127.0, 117.7, 80.7, 57.6, 56.7, 54.5, 28.2; IR (neat) 3080, 3028, 2980, 2929, 1732 cm^{-1} ; HRMS (FAB+) calcd for $\text{C}_{16}\text{H}_{24}\text{NO}_2$ $[\text{M}+\text{H}]^+$: 262.1807, found: 262.1807.

***N*-Allyl-*N*-benzyl-3-(trimethylsilyl)-2-propyn-1-amine (8).** As described for **12**. Purified by chromatography (heptane:Et₂O 1:0→16:1) to afford **8** (0.61 g, 80%) as a colorless oil. ¹H NMR (CDCl₃, 400 MHz) δ_H 7.33 (m, 4H), 7.25 (m, 1H), 5.88 (ddt, 1H, *J*=17.1, 10.2, 6.5), 5.27 (ddt, 1H, *J*=17.1, 1.9, 1.2), 5.17 (ddt, 1H, *J*=10.2, 1.9, 1.2), 3.63 (s, 2H), 3.30 (s, 2H), 3.16 (dt, 2H, *J*=6.5, 1.2), 0.21 (s, 9H); ¹³C NMR (CDCl₃, 125 MHz) δ_C 138.6, 135.6, 129.2, 128.2, 127.1, 118.0, 100.9, 90.2, 57.2, 56.7, 42.3, 0.2; IR (neat) 3081, 3066, 2960, 2920, 2164, 1250, 844 cm⁻¹; HRMS (FAB+) calcd for C₁₆H₂₄NSi [M+H]⁺: 258.1678, found: 258.1666.

(*E*)-*N*-Benzyl-*N*-2-butenyl-2,4-dinitro-benzenesulfonamide (14b). To a solution of Ph₃P (5.8 g, 22.1 mmol) in THF (100 mL) at 0 °C was added DIAD (4.3 ml, 21.8 mmol). After stirring for 30 min at 0 °C a mixture of **13** (5.22 g, 15.5 mmol) and (*E*)-crotyl alcohol (1.35 g, 18.7 mmol) in THF (20 mL) was added and the resultant mixture was stirred at rt for 1 h. Removal of the solvents and chromatography (pentane:EtOAc 9:1→2:1) gave **14b** (5.95 g, 98%) as a yellow oil, which was used in the subsequent step. A sample of the oil was crystallized (EtOAc:hexane) to provide a yellow solid for characterization purposes. Mp 63.4-63.9 °C. ¹H NMR (CDCl₃, 400 MHz) δ_H 8.47 (d, 1H, *J*=2.5), 8.37 (dd, 1H, *J*=8.7, 2.5), 8.12 (d, 1H, *J*=8.7), 7.31-7.21 (m, 5H), 5.55 (dq, 1H, *J*=15.2, 6.5), 5.25 (dtq, 1H, *J*=15.2, 6.7, 1.5), 4.52 (s, 2H), (d, 2H, *J*=6.7), 1.63 (dd, 3H, *J*=6.5, 1.5); ¹³C NMR (CDCl₃, 125 MHz) δ_C 149.4, 147.8, 139.8, 135.0, 132.7, 132.3, 128.7, 128.4, 128.1, 125.8, 124.0, 119.7, 50.6, 49.5, 17.7; IR (neat) 3102, 3079, 3036, 2940, 1552, 1536, 1370, 1348, 1160, 729 cm⁻¹; HRMS (FAB+) calcd for C₁₇H₁₈N₃O₆S [M+H]⁺: 392.0916, found: 392.0914.

(*E*)-*N*-Benzyl-2,4-dinitro-*N*-(3-phenyl-2-propenyl)-benzenesulfonamide (14c). As described for **14b**, starting from (*E*)-cinnamyl alcohol (0.36 g, 2.7 mmol, 1.2 equiv.). Pale yellow semi-solid (1.02 g, quant.). ¹H NMR (CDCl₃, 400 MHz) δ_H 8.46 (d, 1H, *J*=2.2), 8.33 (dd, 1H, *J*=8.7, 2.2), 8.15 (d, 1H, *J*=8.7), 7.33-7.22 (m, 10H), 6.37 (d, 1H, *J*=15.7), 5.95 (dt, 1H, *J*=15.7, 6.8), 4.58 (s, 2H), 4.07 (d, 2H, *J*=6.8); ¹³C NMR (CDCl₃, 125 MHz) δ_C 149.4, 147.8, 139.7, 135.6, 135.5, 134.8, 132.8,

128.8, 128.7, 128.4(×2), 128.3, 126.4, 125.9, 122.2, 119.7, 51.2, 49.8; IR (neat) 3102, 3033, 2927, 1553, 1538, 1365, 1350, 1164, 749 cm⁻¹; HRMS (FAB+) calcd for C₂₂H₁₉N₃O₆S [M]⁺: 453.0995, found: 453.0995.

(Z)-N-Benzyl-N-2-butenyl-2,4-dinitro-benzenesulfonamide (14d). As described for **14b**, starting from (Z)-crotyl alcohol (0.77 g, 10.7 mmol, 1.2 equiv.). Recrystallization (toluene:hexane) provided **14d** as a pale yellow solid (3.0 g, 88%). Mp 96.4-96.7 °C. ¹H NMR (CDCl₃, 400 MHz) δ_H 8.48 (s, 1H), 8.40 (d, 1H, *J*=8.7 Hz), 8.17 (d, 1H, *J*=8.7 Hz), 7.33-7.23 (m, 5H), 5.64 (dq, 1H, *J*=10.7, 7.0 Hz), 5.20 (dt, 1H, *J*=10.7, 7.1 Hz), 4.52 (s, 2H), 3.95 (d, 2H, *J*=7.1 Hz), 1.44 (d, 2H, *J*=7.0 Hz); ¹³C NMR (CDCl₃, 100 MHz) δ_C 149.5, 147.9, 139.7, 134.9, 132.7, 130.3, 128.8, 128.4, 128.2, 125.9, 123.3, 119.7, 50.9, 43.7, 12.8; IR (neat) 3102, 3027, 2937, 1550, 1535, 1368, 1349, 1158, 700 cm⁻¹; HRMS (FAB+) calcd for C₁₇H₁₈N₃O₆S [M+H]⁺: 392.0916, found: 392.0905.

N-Benzyl-N-(3-methyl-2-butenyl)-2,4-dinitro-benzenesulfonamide (14e). As described for **14b**, starting from prenyl alcohol (0.23 g, 2.7 mmol, 1.2 equiv.). Pale yellow semi-solid (0.87 g, 98%). ¹H NMR (CDCl₃, 400 MHz) δ_H 8.48 (d, 1H, *J*=2.2), 8.39 (dd, 1H, *J*=8.7, 2.2), 8.15 (d, 1H, *J*=8.7), 7.33-7.22 (m, 5H), 4.96 (t, 1H, *J*=7.2), 4.51 (s, 2H), 3.89 (d, 2H, *J*=7.2), 1.63 (s, 3H), 1.42 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz) δ_C 149.4, 147.9, 139.9, 138.8, 135.1, 132.7, 128.7, 128.3, 128.1, 125.8, 119.7, 117.6, 50.9, 45.2, 25.7, 17.7; IR (neat) 3100, 3078, 2968, 2934, 1552, 1536, 1368, 1348, 1158, 732 cm⁻¹; HRMS (FAB+) calcd for C₁₈H₂₀N₃O₆S [M+H]⁺: 406.1073, found: 406.1077.

(E)-N-Benzyl-2-buten-1-amine (15b).¹ To a stirred solution of **14b** (12.3 mg, 0.031 mmol) and Et₃N (9 μL) in CH₂Cl₂ (1 mL) was added mercaptoacetic acid (4 mg, 0.043 mmol). The resultant solution was stirred for 1 h and then diluted with Et₂O (5 mL). The organic phase was washed with sat. NaHCO₃ (3×5 mL) and the aqueous phase was extracted with Et₂O (5 mL). The combined organic phases were dried (K₂CO₃) and the concentrated residue was chromatographed

(EtOAc:MeOH 0:1→1:0 + 1% NH₃) to provide **15b** (4 mg, 76%) as a pale yellow oil. ¹H NMR (CDCl₃, 400 MHz) δ_H 7.34-7.22 (m, 5H), 5.66-5.53 (m, 2H), 3.76 (s, 2H), 3.20 (d, 2H, *J*=4.8 Hz), 1.69 (d, 3H, *J*=5.8 Hz), 1.28 (bs, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ_C 140.9, 129.9, 128.8, 128.6, 127.8, 127.3, 53.8, 51.6, 18.2.

(*E*)-*N*-Benzyl-3-phenyl-2-propen-1-amine (15c). As described for **15b**, starting from **14c** (1.0 mg, 2.2 mmol). Pale yellow oil (0.44 g, 88%). ¹H NMR (CDCl₃, 500 MHz) δ_H 7.37-7.19 (m, 10H), 6.53 (d, 1H, *J*=15.9), 6.31 (dt, 1H, *J*=15.9, 6.0), 3.83 (s, 2H), 3.43 (d, 2H, *J*=6.0 Hz), 1.66 (bs, 1H); ¹³C NMR (CDCl₃, 125 MHz) δ_C 140.1, 137.1, 131.4, 128.5, 128.4, 128.3, 128.2, 127.3, 126.9, 126.2, 53.2, 51.1; IR (neat) 3312, 3081, 3060, 3026, 2914, 2815 cm⁻¹; HRMS (EI+) calcd for C₁₆H₁₇N [M]⁺: 223.1361, found: 223.1361.

(*Z*)-*N*-Benzyl-2-buten-1-amine (15d). As described for **15b**, starting from **14d** (23 mg, 0.059 mmol). Pale yellow oil (7 mg, 71%). ¹H NMR (CDCl₃, 400 MHz) δ_H 7.33-7.22 (m, 5H), 5.49-6.63 (m, 2H), 3.80 (s, 2H), 3.30 (d, 2H, *J*=6.3 Hz), 2.00 (bs, 1H), 1.62 (d, 3H, *J*=6.2 Hz); ¹³C NMR (CDCl₃, 100 MHz) δ_C 139.7, 128.4, 128.3, 128.2, 127.1, 126.8, 53.2, 45.2, 13.1; IR (neat) 3309, 3063, 3023, 2917, 2826 cm⁻¹; HRMS (EI+) calcd for C₁₁H₁₅N [M]⁺: 161.1204, found: 161.1193.

***N*-Benzyl-3-methyl-2-buten-1-amine (15e).** As described for **15b**, starting from **14e** (0.86 g, 2.1 mmol). Yellow oil (0.33 g, 90%). ¹H NMR (CDCl₃, 400 MHz) δ_H 7.33-7.23 (m, 5H), 5.29 (m, 1H), 3.80 (s, 2H), 3.24 (d, 2H, *J*=6.8 Hz), 1.73 (s, 3H), 1.70 (bs, 1H), 1.62 (s, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ_C 140.0, 134.8, 128.4, 128.2, 127.0, 122.5, 53.3, 46.5, 25.8, 17.9; IR (neat) 3310, 3062, 3027, 2914, 2855 cm⁻¹; HRMS (EI+) calcd for C₁₂H₁₇N [M]⁺: 175.1361, found: 175.1353.

(*E*)-*N*-Benzyl-2-butenylglycine methyl ester (16b).¹ As described for **12**, starting from **15b** (0.42 g, 2.6 mmol). Colorless oil (0.58 g, 93%). ¹H NMR (CDCl₃, 400 MHz) δ_H 7.36-7.22 (m, 5H), 5.62

(dtq, 1H, J =15.2, 6.3, 1.1 Hz), 5.51 (dqt, 1H, J =15.2, 6.6, 1.5 Hz), 3.75 (s, 2H), 3.68 (s, 3H), 3.30 (s, 2H), 3.19 (dt, 2H, J =6.6, 1.1 Hz), 1.69 (dq, 3H, J =6.3, 1.5 Hz); ^{13}C NMR (CDCl_3 , 100 MHz) δ_{C} 172.0, 138.7, 129.3, 129.0, 128.2, 128.0, 127.0, 57.9, 56.1, 53.6, 51.3, 17.8.

(*E*)-*N*-Benzyl-3-phenyl-2-propenylglycine methyl ester (16c). As described for **12**, starting from **15c** (0.37 g, 1.7 mmol). Pale yellow oil (0.44 g, 91%). ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.42-7.24 (m, 10H), 6.59 (d, 1H, J =15.9), 6.31 (dt, 1H, J =15.9, 6.7), 3.87 (s, 2H), 3.71 (s, 3H), 3.47 (d, 2H, J =6.7), 3.41 (s, 2H); ^{13}C NMR (CDCl_3 , 100 MHz) δ_{C} 171.9, 138.5, 136.9, 133.0, 129.0, 128.5, 128.3, 127.5, 127.16, 127.14, 126.3, 58.1, 56.4, 53.7, 51.4; IR (neat) 3060, 3027, 2950, 2920, 1739 cm^{-1} ; HRMS (EI $^{+}$) calcd for $\text{C}_{19}\text{H}_{21}\text{NO}_2$ $[\text{M}+\text{H}]^{+}$: 296.1651, found: 296.1648.

(*Z*)-*N*-Benzyl-2-butenylglycine methyl ester (16d). As described for **12**, starting from **15d** (0.6 g, 3.7 mmol). Pale yellow oil (0.87 g, quant.). ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.36-7.22 (m, 5H), 5.66 (dq, 1H, J =11.0, 6.8 Hz), 5.52 (m, 1H), 3.78 (s, 2H), 3.68 (s, 3H), 3.31 (s, 2H), 3.29 (d, 2H, J =7.0 Hz), 1.62 (d, 3H, J =6.8 Hz); ^{13}C NMR (CDCl_3 , 100 MHz) δ_{C} 172.0, 138.8, 129.1, 128.3, 127.9, 127.2, 127.1, 58.0, 53.7, 51.3, 50.0, 13.1; IR (neat) 3062, 3025, 2951, 2918, 1739 cm^{-1} ; HRMS (FAB $^{+}$) calcd for $\text{C}_{14}\text{H}_{18}\text{NO}_2$ $[\text{M}-\text{H}]^{+}$: 232.1338, found: 232.1331.

***N*-Benzyl-3-methyl-2-butenylglycine methyl ester (16e).** As described for **12**, starting from **15e** (0.29 g, 1.7 mmol). Yellow oil (0.36 g, 88%). ^1H NMR (CDCl_3 , 400 MHz) δ_{H} 7.35-7.22 (m, 5H), 5.27 (t, 1H, J =7.1), 3.75 (s, 2H), 3.68 (s, 3H), 3.29 (s, 2H), 3.22 (d, 2H, J =7.1), 1.73 (s, 3H), 1.62 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ_{C} 172.1, 138.9, 136.0, 129.0, 128.2, 127.0, 121.3, 58.1, 53.6, 51.32, 51.28, 25.9, 17.9; IR (neat) 3062, 3028, 2969, 2950, 1739 cm^{-1} ; HRMS (FAB $^{+}$) calcd for $\text{C}_{15}\text{H}_{21}\text{NO}_2$ $[\text{M}+\text{H}]^{+}$: 248.1651, found: 248.1645.

^1H NMR

^{13}C NMR

2b

^1H NMR

^{13}C NMR

¹H NMR

¹³C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

7

^1H NMR

^{13}C NMR

8

^1H NMR

^{13}C NMR

^1H NMR

^{11}B NMR

^{19}F NMR

NB! $^{10}\text{BF}_4^-$ singlet peak also visible!

9b; X=Br or BBr₄

¹H NMR

NB! Contains hexanes from BBr₃ in hexanes solution.

¹¹B NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

11

^1H NMR

^{13}C NMR

12

¹H NMR

¹³C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

^1H NMR

^{13}C NMR

15b

¹H NMR

¹³C NMR

15c

¹H NMR

¹³C NMR

15d

^1H NMR

^{13}C NMR

15e

¹H NMR

¹³C NMR

16b

¹H NMR

¹³C NMR

16c

^1H NMR

^{13}C NMR

16d

^1H NMR

^{13}C NMR

16e

¹H NMR

¹³C NMR

References

- (1) Coldham, I.; Middleton, M. L.; Taylor, P. L. *J. Chem. Soc. Perkin Trans. 1* **1998**, 2817.
- (2) Guo, Z.; Dowdy, E. D.; Li, W.; Polniaszek, R.; Delaney, E. *Tetrahedron Lett.* **2001**, 42, 1843.
- (3) Fukuyama, T.; Cheung, M.; Jow, C. K.; Hidai, Y.; Kan, T. *Tetrahedron Lett.* **1997**, 38, 5831.