SUPPORTING INFORMATION

Title: Lewis Acid Promoted Annulation of *o*-Quinonediimines by Allylstannane: A Facile Synthesis of Quinoxaline Derivatives

Vijay Nair, R. Dhanya, C. Rajesh, M. M. Bhadbhade and K. Manoj

Experimental Section

General: All reactions were carried out in oven-dried glasswares under argon atmosphere. Progresses of reactions were monitored by Thin Layer Chromatography while purification was effected using silica gel column chromatography. NMR data were collected at room temperature in CDCl₃ at operating frequency 300 MHz. IR spectra were recorded in Bomem MB Series FT-IR spectrophotometer. Melting points were recorded on a Buchi melting point apparatus and are uncorrected. Allyltributyltin was purchased from Aldrich and was used as supplied. All the quinoneimides were synthesized according to the literature procedure.

Synthesis of tetrahydroquinoxaline derivatives and allylated amides from *o*-quinonediimides and allyltributyltin **General Procedure:** A solution of *o*-quinonediimide **1** or **4** (0.2 mmol) and allyltributyltin (0.24 mmol) in dry dichloromethane was cooled to 0 °C. To this well cooled solution zinc chloride (0.24 mmol) was added and the resulting mixture was stirred at the same temperature for 30 min. and then quenched with water. The reaction mixture was extracted with dichloromethane (3X10 mL), washed with water and saturated brine solution. It was then concentrated and the residue was subjected to silica gel column chromatography using hexane-ethylacetate solvent mixture afforded the products, which were recrystallized from hexane-dichloromethane solvent mixture.

Tetrahydroquinoxaline **3a**.

To a solution of allyltributylstannane **2a** (79 mg, 0.24 mmol) and 4-chloro-*o*-benzoquinone dibenzenesulfonimide **1a** (84 mg, 0.2mmol) in 4 mL CH₂Cl₂ at 0 °C ZnCl₂ (33 mg, 0.24 mmol) was added and the reaction mixture was stirred at 0 °C for 30

min. It was then processed as described in the general procedure to afford the tetrahydroquinoxaline derivative **3a** (150 mg, 98%).

Colorless crystalline solid Mp 79-80 °C; **IR** (KBr) ν_{max} : 2959, 2922, 2845, 1594, 1485, 1356, 1243, 1171, 1089, 1027, 939, 893, 805, 589 cm^{-1} ; **^1H NMR**: δ 0.87-0.94 (m, 18H), 1.34 -1.51 (m, 11H), 3.01 (dd, 1H, J = 12.4 Hz, 4.0 Hz), 3.75 (dd, 1H, J = 12.4 Hz, J = 3.4 Hz), 4.59 -4.68 (m, 1H), 7.06 (dd, 1H, J = 9.1 Hz, 2.4 Hz), 7.34 -7.48 (m, 6H), 7.53 -7.56 (m, 5H), 7.81 (d, 1H, J = 2.4 Hz); **^{13}C NMR**: δ 9.3, 13.3, 13.7, 27.4, 29.1, 50.3, 53.7, 118.7, 119.9, 123.5, 126.0, 126.2, 126.9, 127.0, 127.2, 127.3, 128.7, 129.2, 129.3, 131.5, 133.1, 133.2, 133.3, 138.2, 139.1.

Elemental analysis calculated for $\text{C}_{33}\text{H}_{45}\text{ClN}_2\text{O}_4\text{S}_2\text{Sn}$: C, 52.71; H, 6.03; N, 3.73; S, 8.53. Found: C, 52.92; H, 6.43; N, 4.00; S, 8.77.

Tetrahydroquinoxaline 3b.

To a solution of allyltributylstannane **2a** (79 mg, 0.24 mmol) and 4-bromo-*o*-benzoquinone dibenzenesulfonimide **1b** (93 mg, 0.2 mmol) in 4 mL CH_2Cl_2 at 0 °C ZnCl_2 (33 mg, 0.24 mmol) was added and the reaction mixture was stirred at 0 °C for 30 min. It was then processed as described in the general procedure to afford the tetrahydroquinoxaline derivative **3b** (115 mg, 72%).

Colorless crystalline solid Mp 80-82 °C; **^1H NMR**: δ 0.83 -0.92 (m, 18H), 1.26 -1.47 (m, 11H), 3.00 (dd, 1H, J = 12.5 Hz, J = 4.0 Hz), 3.75 (dd, 1H, J = 12.5 Hz, 3.4 Hz), 4.65-4.63 (m, 1H), 7.19-7.28 (m, 1H), 7.42-7.48 (m, 6H), 7.52-7.59 (m, 5H), 7.94 (d, 1H, J = 2.4 Hz); **^{13}C NMR**: δ 9.3, 13.2, 13.8, 27.4, 29.7, 50.3, 53.6, 119.6, 120.1, 121.6, 126.2, 126.4, 126.9, 127.2, 127.3, 129.0, 129.1, 129.3, 130.2, 133.1, 133.2, 133.3, 133.4, 138.3, 139.0. Analysis calculated for $\text{C}_{35}\text{H}_{45}\text{BrN}_2\text{O}_4\text{S}_2\text{Sn}$: C, 49.76; H, 5.69; N, 3.52; S, 8.20. Found: C, 50.26; H, 6.02; N, 3.93; S, 8.89.

Tetrahydroquinoxaline 3c.

To a solution of allyltributylstannane **2a** (79 mg, 0.24 mmol) and 4-methyl-*o*-benzoquinone dibenzenesulfonimide **1c** (80 mg, 0.2 mmol) in 4 mL CH_2Cl_2 at 0 °C ZnCl_2 (33 mg, 0.24 mmol) was added and the reaction mixture was stirred at that temperature for 30 min. It was then processed as described in the general procedure to afford the tetrahydroquinoxaline derivative **3c** (107 mg, 72%).

Colorless crystalline solid Mp 48-50 °C; **IR** (neat) ν_{max} : 2995, 2921, 2854, 1499, 1445, 1357, 1162, 1088, 724, 683 cm^{-1} ; **^1H NMR**: δ 0.87-0.95 (m, 18H), 1.26-1.43 (m, 11H), 2.32 (s, 1H), 3.07 (dd, 1H, $J = 12.4$ Hz, 4.1 Hz), 3.65 (dd, 1H, $J = 12.4$ Hz, $J = 3.9$ Hz), 4.67-4.60 (m, 1H), 6.88 (uneven triplet, 1H, $J_1 = 10.2$ Hz, $J_2 = 8.7$ Hz), 7.26-7.38 (m, 6H), 7.94 (d, 1H, $J = 2.4$ Hz); **^{13}C NMR**: δ 9.3, 13.6, 13.7, 20.8, 27.9, 29.2, 50.6, 54.2, 118.8, 119.4, 124.5, 126.9, 127.0, 127.1, 127.3, 127.7, 128.2, 129.0, 129.1, 132.9, 132.9, 133.0, 133.3.

Tetrahydroquinoxaline 3d.

To a solution of allyltributylstannane **2a** (79 mg, 0.24 mmol) and 4,5-dichloro-*o*-benzoquinone dibenzenesulfonimide **1d** (91 mg, 0.2 mmol) in 4 mL CH_2Cl_2 at 0 °C ZnCl_2 (33 mg, 0.24 mmol) was added and the reaction mixture was stirred at that temperature for 30 min. It was then processed as described in the general procedure to afford the tetrahydroquinoxaline derivative **3d** (130 mg, 82%).

Colorless crystalline solid Mp 52-54 °C; **IR** (KBr) ν_{max} : 2964, 2928, 1480, 1362, 1285, 1171, 1084, 1032, 903, 728, 687 cm^{-1} ; **^1H NMR**: δ 0.88-0.92 (m, 18H), 1.34-1.45 (m, 11H), 2.84 (dd, 1H, $J = 12.4$ Hz, 3.8 Hz), 3.79 (dd, 1H, $J = 12.4$ Hz, $J = 2.8$ Hz), 4.59-4.64 (m, 1H), 7.35-7.61 (m, 10H), 7.88 (s, 1H), 7.91 (s, 1H); **^{13}C NMR**: δ 9.3, 13.0, 13.7, 27.8, 29.2, 49.7, 53.0, 119.9, 124.0, 126.8, 127.0, 127.2, 128.4, 129.3, 129.6, 129.9, 133.3, 133.5, 138.0, 138.3.

Analysis calculated for $\text{C}_{33}\text{H}_{44}\text{Cl}_2\text{N}_2\text{O}_4\text{S}_2\text{Sn}$: C, 50.40; H, 5.64; N, 3.56; S, 8.15. Found: C, 50.59; H, 6.06; N, 3.66; S, 8.08.

Tetrahydroquinoxaline 3e.

To a solution of allyltributylstannane **2a** (79 mg, 0.24 mmol) and 4,5-dimethyl-*o*-benzoquinone dibenzenesulfonimide **1e** (82 mg, 0.2 mmol) in 4 mL CH_2Cl_2 at 0 °C ZnCl_2 (33 mg, 0.24 mmol) was added and the reaction mixture was stirred at that temperature for 30 min. It was then processed as described in the general procedure to afford the tetrahydroquinoxaline derivative **3e** (66 mg, 44%).

Colorless crystalline solid Mp 52-54 °C; **IR** (KBr) ν_{max} : 2962, 2928, 1650, 1521, 1485, 1356, 1289, 1197, 1073, 712 cm^{-1} ; **^1H NMR**: δ 0.86-0.91 (m, 18H), 1.28-1.44 (m, 11H), 2.18 (s, 3H), 2.21 (s, 3H), 3.01 (dd, 1H, $J = 12.3$ Hz, $J = 4.1$ Hz), 3.64 (dd, 1H, $J = 12.3$ Hz, 3.8 Hz), 4.57-4.64 (m, 1H), 7.57-7.132 (m, 12H); **^{13}C NMR**: δ 9.3, 13.5, 13.7, 19.2, 20.0, 27.4, 29.1, 50.5, 54.0, 119.8, 122.6, 127.1, 127.3, 128.5, 128.6, 129.1, 132.0, 132.8, 132.9, 134.8, 138.6, 139.7.

Analysis calculated for $\text{C}_{35}\text{H}_{50}\text{N}_2\text{O}_4\text{S}_2\text{Sn}$: C, 56.38; H, 6.76; N, 3.76; S, 8.60. Found: C, 56.57; H, 7.18; N, 3.86; S, 8.63.

Allylated amide 5a.

A solution of *o*-quinonedibenzimide **4a** (62 mg, 0.2 mmol) and allyltributyltin **2a** (79 mg, 0.24 mmol) in 4 mL dry CH_2Cl_2 was cooled to 0 °C. To this solution ZnCl_2 (33 mg, 0.24 mmol) was added the reaction mixture was stirred at 0 °C for 30 min. It was then processed as described in the general procedure to afford **5a** (44 mg, 61%) as a colorless solid.

Mp 186-188 °C; **IR** (KBr) ν_{max} : 3226, 1650, 1602, 1578, 1519, 1472, 910, 823, 705 cm^{-1} ; **^1H NMR**: δ 2.63 (d, 2H, $J = 6.2$ Hz), 4.62-4.75 (m, 2H), 5.289-5.42 (m, 1H), 6.43 (d, 1H, $J = 8.2$ Hz), 7.03 (s, 1H), 7.25 (d, 1H, $J = 8.2$ Hz), 7.51-7.57 (m, 6H), 8.03-8.09 (m, 4H), 9.62 (s, 1H, *exchangeable with D₂O*), 9.69 (s, 1H, *exchangeable with D₂O*); **^{13}C NMR**: δ 38.9, 115.8, 125.5, 126.0, 126.3, 127.8, 127.9, 128.6, 128.8, 130.6, 132.1, 133.5, 133.6, 136.4, 138.1, 166.4. HRMS calculated for $\text{C}_{23}\text{H}_{20}\text{N}_2\text{O}_2$: 356.1567 Found 356.1525.

Allylated amide 5b.

A solution of *o*-quinoneimide **4b** (66 mg, 0.2 mmol) and allyltributyltin **2a** (79 mg, 0.24 mmol) in 4 mL dry CH_2Cl_2 was cooled to 0 °C. To this solution ZnCl_2 (33 mg, 0.24 mmol) was added and the reaction mixture was stirred at 0 °C for 30 min. It was then processed as described in the general procedure to afford **5b** as a colorless solid (50 mg, 68%).

Mp 202-204 °C; **IR** (KBr) ν_{max} : 3265, 2854, 1652, 1604, 1529, 1468, 1266, 997, 892 cm^{-1} ; **^1H NMR**: δ 2.21 (s, 3H), 2.46 (d, 2H, J = 6 Hz), 4.99-5.03 (m, 2H), 5.85-5.88 (m, 1H), 7.44 (s, 1H), 7.50 (s, 1H), 7.54-7.57 (m, 6H), 7.84 (m, 4H, J = 9.0 Hz), 9.60 (s, 1H, *exchangeable with D₂O*), 9.63 (s, 1H, *exchangeable with D₂O*). **^{13}C NMR**: δ 19.4, 37.7, 111.8, 114.8, 114.9, 125.9, 127.2, 127.6, 133.0, 130.7, 135.3, 137.2, 139.4, 162.5, 164.4.

Allylated amide **5c**.

A solution of **4c** (70 mg, 0.2 mmol) and allyltributyltin **2a** (79 mg, 0.24 mol) in 4 mL dry CH_2Cl_2 was cooled to 0 °C and ZnCl_2 (33 mg, 0.24 mmol) was added. The reaction mixture was stirred at 0 °C for 30 min. It was then processed as described in the general procedure to afford **5c** as a colorless solid (53 mg, 68%).

Mp 220-222 °C; **IR** (KBr) ν_{max} : 3265, 2854, 1647, 1600, 1526, 1458, 1276, 993, 872 cm^{-1} ; **^1H NMR**: δ 2.75 (d, 2H, J = 6.3 Hz), 4.59 (d, 1H, J = 17.0 Hz), 4.72 (d, 1H, J = 10.0 Hz), 5.27-5.39 (m, 1H), 7.04 (s, 1H), 7.43 (s, 1H), 7.53-7.58 (m, 6H), 8.02-8.08 (m, 4H), 9.65 (s, 1H, *exchangeable with D₂O*), 9.69 (s, 1H, *exchangeable with D₂O*); **^{13}C NMR**: δ 36.5, 116.4, 126.6, 126.9, 127.7, 127.8, 128.2, 128.9, 129.4, 133.0, 131.3, 132.4, 132.3, 134.4, 135.7, 166.5, 166.6. HRMS (EI) calculated for $\text{C}_{23}\text{H}_{19}\text{N}_2\text{O}_2\text{Cl}$: 390.1135. Found 390.1114.

Allylated amide **5d**.

A solution of **4d** (76 mg, 0.2 mmol) and allyltributyltin **2a** (79 mg, 0.24 mmol) in 4 mL dry CH_2Cl_2 was cooled to 0 °C and ZnCl_2 (33 mg, 0.24 mmol) was added. The reaction mixture was stirred at 0 °C for 30 min. It was then processed as described in the general procedure to afford **5d** as a colorless solid (65 mg, 77%).

Mp 230-232 °C; **IR** (KBr) ν_{max} : 3265, 2921, 2854, 1647, 1600, 1526, 1458, 1276, 993, 872 cm^{-1} ; **^1H NMR**: δ 2.75 (m, 2H), 4.95-5.06 (m, 2H), 5.42-5.50 (m, 1H), 7.53-7.58 (m, 6H), 7.80-7.85 (m, 4H), 8.32 (s, 1H), 9.68 (s, 1H, *exchangeable with D₂O*), 9.71 (s, 1H, *exchangeable with D₂O*); **^{13}C NMR**: δ 31.2, 119.4, 122.8, 123.2, 124.1, 125.9, 126.7, 127.3, 128.6, 130.7, 132.1, 132.8, 132.9, 139.2, 162.3, 164.3.

Allylated amide **5e**.

A solution of **4e** (68 mg, 0.2 mmol) and allyltributyltin **2a** (79 mg, 0.24 mmol) in 4 mL dry CH₂Cl₂ was cooled to 0 °C and ZnCl₂ (33 mg, 0.24 mmol) was added. The reaction mixture was stirred at 0 °C for 30 min. It was then processed as described in the general procedure to afford **5e** as a colorless solid (52 mg, 69%).

Mp 210-212°C; **IR** (KBr) $\nu_{\max}$: 3265, 2921, 2854, 1649, 1603, 1546, 1458, 1276, 993, 872 cm⁻¹; **¹H NMR**: δ 2.16 (s, 6H), 3.52 (m, 2H), 4.93 (d, 1H, J = 17.2 Hz), 5.11 (d, 1H, J = 10.2 Hz), 6.97-5.06 (m, 1H), 7.27-7.56 (m, 7H), 8.86-7.86 (m, 4H), 8.43 (s, 1H, *exchangeable with D₂O*), 9.20 (s, 1H, *exchangeable with D₂O*); **¹³C NMR**: δ 15.6, 20.5, 33.6, 115.9, 125.2, 126.9, 127.4, 127.5, 128.5, 128.9, 130.9, 131.6, 132.1, 133.6, 134.2, 135.5, 136.4, 165.7, 166.5. HRMS (EI) calculated for C₂₅H₂₄N₂ O₂: 384.1838. Found: 384.178

Tetrahydroquinoxaline **3e'**

To a solution of the tetrahydroquinoxaline derivative **3e** (149 mg, 0.2 mmol) in isopropanol at room temperature bromine (24 mg, 0.3 mmol) was added and the reaction mixture was stirred at room temperature for 2h. Excess isopropanol was removed under vacuum. The residue was dissolved in CH₂Cl₂ and cooled to 0 °C and added *m*-CPBA (106 mg, 0.6 mmol). The mixture was allowed to stir for 5 min. at 0 °C, and ammonium solution was added. The mixture was warmed to room temperature, and extracted three times with saturated aqueous sodium carbonate solution, dried over sodium sulfate, and concentrated. Purification was achieved by silica gel column chromatography using hexane-ethyl acetate mixture (85:15) afforded **3e'** as a colorless crystalline solid in 86% yield (82 mg).

Colorless crystalline solid Mp 118-120 °C; **IR** (KBr) $\nu_{\max}$: 3495, 3006, 2944, 1729, 1393, 1268, 1150, 1098, 1025, 570 cm⁻¹; **¹H NMR**: δ 1.25 (s, 1H, *exchangeable with D₂O*), 2.13 (s, 3H), 2.22 (s, 3H), 3.30 (dd, 1H, J = 13.1 Hz, 5.3 Hz), 3.59-3.64 (m, 2H), 3.78 (dd, 1H, J = 13.1 Hz, 5.7 Hz), 4.47-4.55 (m, 1H), 7.13 (s, 1H), 7.26 (s, 1H), 7.40-7.60 (m, 10H); **¹³C NMR**: δ 19.8, 29.8, 45.7, 57.3, 62.1, 121.0, 125.5, 126.8, 127.5, 127.7,

127.8, 128.2, 129.2, 129.4, 129.7, 130.0, 130.3, 132.3, 133.1, 133.3, 135.0. 133.5, 134.7,
HRMS calculated for $C_{23}H_{24}N_2O_5S_2$: 472.1121. Found: 472.1126.

CCDC Deposition Number of Compound **3a**

CCDC 253649