Lewis Acid Tuned Facial Stereodivergent HDA Reactions Using β -substituted *N*-vinylloxazolidinones.

Frédéric Gohier,[†] Keltoum Bouhadjera,¹ Djibril Faye,[†] Catherine Gaulon,[†]
Vincent Maisonneuve,[‡] Gilles Dujardin^{†*} and Robert Dhal^{†*}

[†]UMR 6011 CNRS-Université du Maine, Laboratoire de Synthèse Organique (UCO2M) and [‡]UMR 6010, Laboratoire des Oxydes et Fluorures, Faculté des sciences, Avenue Olivier Messiaen, 72085 Le Mans Cedex 9, France.

SUPPORTING INFORMATION

Table of contents	page
Experimental section	2
General	2
X-ray crystal structure determination of endo α 13e and endo β 13c	2
Experimentals	4
Copy of ¹ H and ¹³ C NMR	13

¹ Present adress : Laboratoire de Chimie organique, Université Aboubekre Belkaid, Tlemcen, Algérie.

Experimental Section

General. NMR spectra were recorded on a 200- or 400-MHz spectrometer. ^{13}C NMR spectra were obtained with broadband proton decoupling. For spectra recorded in CDCl_3 , chemical shifts are recorded relative to the internal TMS (tetramethylsilane) reference signal. For $\text{DMSO}-d_6$ and CD_3SOCD_3 used as solvents, chemical shifts are given relative to the solvent signals. Elemental analyses were obtained from the Service de microanalyse, CNRS ICSN, Gif-sur-Yvette and HRMS analyses were done by the CRMPO, Rennes. All melting points are uncorrected.

X-ray crystal structure determination of *endo* α 13e and *endo* β 13c

A single-crystal, selected using an optical microscope, was mounted on a glass fiber. The crystal cell parameters were obtained from a random reflection search and X-ray diffraction data were collected with a Siemens AED2 four-circle diffractometer equipped with a graphite monochromatized $\text{MoK}\alpha_1$ ($\lambda = 0.71073 \text{ \AA}$). The intensities of the reflections were corrected for Lorentz and polarization effects but no absorption correction was applied. The scattering factors and the anomalous dispersions corrections for all atoms were taken from International Tables for X-ray Crystallography. Table gathers the crystallographic data and information about data collection. The space group was deduced from the observed systematic extinctions. The structure was solved by direct methods of the program SHELXS-86 and successive refinements and difference Fourier maps using SHELXL-97 program. The location of the hydrogen atoms was performed by applying the geometrical constraints using AFIX and DFIX option in SHELXL-97 program.

Table . Crystal data and structure refinement of *endo* α 13e and

endo β 13c

Identification code	<i>endo</i> β 13c	<i>endo</i> α 13e
Empirical formula	C ₁₉ H ₂₄ O ₅ N	C ₂₇ H ₃₁ O ₈ N
Formula mass (g.mol ⁻¹)	346.40	497.55
Unit cell dimensions		
a =	5.377 (1) Å	9.945 (5) Å
b =	10.277 (3) Å	11.10 (1) Å
c =	16.505 (5) Å	24.14 (2) Å
α =	90°	90°
β =	91.25 (2)°	90°
γ =	90°	90°
Volume, Z	911.8 (3) Å ³ , 2	2663 (3) Å ³ , 4
Calculated density (g.cm ⁻³)	1.258	1.241
Crystal system, Space group	Monoclinique, <i>P</i> 2 ₁	Monoclinic, <i>P</i> 2 ₁ 2 ₁ 2 ₁
Temperature (K), scan method	293, 2 θ / ω	293, 2 θ / ω
2 θ range for data collection (°)	2-60	2-55
(hkl) _{min} , (hkl) _{max}	-7 -14 -23, 7 14 23	-12 -14 -31, 12 14 31
Absorption correction, coefficient (mm ⁻¹)	Not applied, 0.09	Not applied, 0.09
Reflections (<i>I</i> > 2 σ (<i>I</i>)) and refined numbers	2749, 230	1191, 191
Goodness of fit (<i>F</i> ²)	1.027	0.98
R ₁ [<i>I</i> > 2 σ (<i>I</i>)], wR ₂ (all data)	0.077, 0.245	0.075, 0.232

General procedure for the synthesis of *N,O*-acetals **8d and **8e**:** A mixture of oxazolidinone **6** or **7** (1 equiv), acetal **2** to **5** (5 equiv), and (*d,l*)-camphorsulfonic acid (0.1 equiv) in dichloromethane was heated overnight at 55 °C. After cooling and dilution with diethyl ether, a saturated aqueous NaHCO₃ solution was added and the organic phases were washed with brine and dried over MgSO₄. Removal of solvent yielded *N,O*-acetals **8a-e** as a crude diastereoisomeric mixture.

8a and **8b** were not isolated. **8c**² was already completely characterised.

(*R*)-3-(1-ethoxypropyl)-4-ethyloxazolidin-2-one **8d (entry 4, table 1):** According to the general procedure, **8d** was obtained as an oil (97 %) used without any purification. ¹H NMR (400 MHz, CDCl₃) δ TMS: 4.99 (1H, t, *J* = 7.0 Hz), 4.33 (1H, t, *J* = 8.6 Hz), 4.01 (1H, dd, *J* = 5.3 and 8.6 Hz), 3.90 and 3.75 (1H, 2m), 3.75 and 3.75 (2H, 2m, OCH₂CH₃), 2.10-1.50 (4H, m), 1.20 (3H, 2t, *J* = 7.7 Hz, OCH₂CH₃), 0.94 (6H, m). IR (KBr): 2970, 1740, 1410, 1220, 1051 cm⁻¹. HRMS calcd for C₈H₁₄NO₃ [M⁻CH₂CH₃] 172.0973, found 172.0982.

(*R*)-3-(3-(benzo[d][1,3]dioxol-6-yl)-2-methoxypropyl)-4-ethyloxazolidin-2-one **8e (entry 5, table 1):** According to the general procedure, **8e** was obtained as an oil (45 %) after column chromatography purification (toluene/AcOEt, 95/5). ¹H NMR (400 MHz, CDCl₃) δ TMS: 6.74 (1H, d, *J* = 7.8 Hz), 6.68 (1H, d, *J* = 1.5 Hz), 6.62 (1H, dd, *J* = 1.5 and 7.9 Hz), 5.93 (2H, s, OCH₂O), 4.98 (1H, 2dd, *J* = 5.8 and 7.6 Hz), 4.38 (minor) and 4.28 (major) (1H, 2t, *J* = 8.6 Hz), 4.02 (1H, 2dd, *J* = 5.6 and 8.6 Hz), 3.76 (1H, m), 3.38 (major) and 3.31 (minor) (1H, 2s, OCH₃), 2.62 (2H, m), 2.04, 1.86 and 1.58 (4H, 2m), 0.91 (minor) et 0.89 (major) (3H, 2t, *J* = 7.3 Hz). IR (KBr): 1739, 1488, 1410, 1242, 1033 cm⁻¹. HRMS calcd for C₁₆H₂₁NO₅ [M⁺] 307.1419, found 307.1419.

² Gaulon, C. ; Dujardin, G. *Synthesis* **2003**, 2269.

General procedure for the synthesis of vinyloxazolidinone 9a-e: To a cooled solution (0°C) of crude *N,O*-acetal **8c-e**, in anhydrous CH₂Cl₂ was added under nitrogen distilled Et₃N (1.5 equiv), then dropwise TMSOTf (1.3 equiv). After slowly warming to r.t. and stirring for 15h, the mixture was treated with basic alumina to remove the excess of TMSOTf. After removal of the solvent, the residue was filtered on silica gel (eluent : ether) to remove the ammonium triflate salts and was purified by chromatography. **9a** and **9c** was already fully characterised.

(R)-4-ethyl-3-styryloxazolidin-2-one (entry 2, table 1) 9b : mp 113.8-114.3°C. ¹H NMR (400 MHz, CDCl₃) δ TMS: 7.32 (4H, m), 7.28 (1H, d, *J* = 15.0 Hz), 7.17 (1H, m), 5.88 (1H, d, *J* = 15.0 Hz), 4.47 (1H, t, *J* = 9.3 Hz), 4.20 (2H, m), 1.84 (2H, m), 0.96 (3H, t, *J* = 7.5 Hz). ¹³C NMR (100 MHz, CDCl₃) δ 155.3, 135.9, 128.6, 126.5, 125.3, 122.8, 111.1, 67.0, 54.6, 23.4, 7.5. [α]_D²⁰ -19.9 (c 1, acetone). IR (KBr): 1737, 1654, 1414, 1221 cm⁻¹. Anal. Calcd for C₁₃H₁₅NO₂ : C, 71.87 ; H, 6.96. Found: C, 72.15 ; H, 6.89.

(R)-4-ethyl-3-((E)-prop-1-enyl)oxazolidin-2-one 9d (entry 4, table 1): ¹H NMR (400 MHz, CDCl₃) δ TMS: 6.51 (1H, dq, *J* = 1.6 and 14.5 Hz), 4.95 (1H, dq, *J* = 6.7 and 14.5 Hz), 4.39 (1H, t, *J* = 8.5 Hz), 4.09 (1H, dd, *J* = 8.5 Hz), 4.03 (1H, m), 1.78 (2H, m), 1.72 (3H, dd, *J* = 1.6 and 6.7 Hz), 0.90 (3H, t, *J* = 7.5 Hz). ¹³C NMR (100 MHz, CDCl₃) δ 155.4, 123.3, 106.4, 66.8, 54.8, 23.4, 7.5. [α]_D²⁰ -39.9 ° (c 1, acetone). IR (KBr): 1735, 1412, 1217, 1052 cm⁻¹. ¹H.RMS calcd for C₈H₁₃NO₂ [M⁺] 155.0946, found 155.0939.

(R)-3-((E)-3-(benzo[d][1,3]dioxol-6-yl)prop-1-enyl)-4-ethyloxazolidin-2-one (entry 5, table 1) 9e : mp 113.8-114.3°C. ¹H NMR (400 MHz, CDCl₃) δ TMS: 6.74 (1H, d, *J* = 7.9 Hz), 6.68 (1H, d, *J* = 1.5 Hz), 6.64 (1H, dd, *J* = 1.5 and 7.9 Hz), 6.62 (1H, dt, *J* = 1.5 and 14.3 Hz), 5.93 (2H, s, OCH₂O), 5.02 (1H, dt, *J* = 7.4 and 14.3 Hz), 4.40 (1H, t, *J* = 8.6 Hz), 4.11 (1H, dd, *J* = 3.4 and 8.4 Hz), 4.02 (1H, m), 3.31 (2H, m), 1.79 et 1.70 (2H, 2m), 0.90 (3H, t, *J* = 7.6 Hz). ¹³C NMR (100 MHz, CDCl₃) δ: 155.5, 147.8, 146.0, 135.3, 123.8,

121.1, 110.4, 108.9, 108.3, 100.9, 67.1, 54.8, 36.2, 23.5, 7.7. $[\alpha]_D^{20}$ -18.8 (c 1, acetone). IR (KBr): 1742, 1666, 1486, 1409, 1242, 1034 cm^{-1} . HRMS calcd for $\text{C}_{15}\text{H}_{17}\text{NO}_4$ $[\text{M}^+]$ 275.1157, found 275.1169.

Heterocycloaddition between dienophiles **9a-e** and dienes **10** and **12**

(2-oxo-1,1-oxazolidin-3-yl)-4,5-diphenyl-5,6-dihydro-4H-pyran-2-carboxylic acid methyl ester endo 11 : To a stirred solution of heterodiene **10** (1 equiv) and *N*-vinyl oxazolidinone **9a** (1 equiv) in solvent (see table 2), a Lewis acid was slowly added (conditions are described in table 2). The treatment depends on the Lewis acid used. For entries 1 and 2 the treatment is like the general procedure A and for the entries 3 to 9 the procedure B has been followed. The crude product was purified by chromatography (toluene/AcOEt 95/5).

^1H NMR (400 MHz, CDCl_3) δ TMS: 6.80-7.24 (m, 10H), 6.27 (d, 1H, J = 2.5Hz), 6.02 (d, 1H, J = 10.4 Hz), 4.29 (dt, 1H, J = 6.1 and 9.0 Hz), 3.99 (dt, 1H, J = 7.6 and 9.0 Hz), 3.84 (s, 3H), 3.75-3.82 (m, 2H), 3.31 (ddd, 1H, J = 6.1, 7.6 and 9.3 Hz), 2.93 (t, 1H, J = 10.4 Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 162.2, 157.3, 143.6, 140.1, 137.1, 125.4-128.5, 115.2, 83.5, 62.0, 52.1, 50.3, 48.5, 39.0. IR (KBr): 1756, 1735, 1660, 1492, 1452, 1413, 1271, 1249, 1226, 1130, 1094 cm^{-1} . Anal. Calcd for $\text{C}_{22}\text{H}_{21}\text{NO}_5$: C, 69.64; H, 5.58. Found: C, 69.55; H, 5.67.

General procedure A : A solution of heterodiene **10** or **12** (1 equiv.), *N*-vinyl oxazolidinone **9b-e** (1 equiv) and $\text{Eu}(\text{fod})_3$ (0.05 equiv) in cyclohexane (5mL) was refluxed under nitrogen. After removal of solvent, the crude product was purified by column chromatography (toluene/AcOEt, 95/5).

General procedure B : To a stirred solution of heterodiene **10** or **12** (1 equiv) and *N*-vinyl oxazolidinone **9b-e** (1 equiv) in dichloromethane, SnCl₄ (1M solution in dichloromethane, 0.5 equiv) was slowly added at -78°C under argon. The mixture was stirred for 3 h at -78 °C then treated with a saturated aqueous solution of NaHCO₃. The resulting solution was allowed to warm up to ambient temperature and the two layers were separated. The organic phase was dried under MgSO₄. After removal of solvent, the crude product was purified by chromatography (toluene/AcOEt 95/5).

(4*R*,5*S*,6*R*)-6-((4*R*)-4-Ethyl-2-oxo-oxazolidin-3-yl)-4,5-diphenyl-5,6-dihydro-4*H*-pyran-2-carboxylic acid methyl ester endo α 13b: According to the general procedure A, **13b** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (57%). mp 52-55 °C. ¹H NMR (400 MHz, CDCl₃) δ TMS: 7.18 (6H, m), 6.87 (4H, m), 6.24 (1H, d, *J* = 2.4 Hz), 5.99 (1H, d, *J* = 10.5 Hz), 3.83 (5H, s and m), 3.80 (1H, dd, *J* = 2.4 and 10.5 Hz), 3.64 (1H, m), 3.12 (1H, t, *J* = 10.6 Hz), 1.82 and 1.70 (2H, 2m), 0.90 (3H, t, *J* = 7.3 Hz). ¹³C NMR (100 MHz, CDCl₃) δ 162.4 , 157.2 , 143.6, 140.3, 137.3, 128.6, 128.3, 127.8, 127.5, 127.1, 114.9, 84.2, 66.9, 54.0, 52.2 , 49.4, 48.6, 26.7, 7.9. [α]_D²⁰ +110.8 (c 1, acetone). IR (KBr): 1731, 1651, 1602, 1269, 1109 cm⁻¹. Anal. Calcd for C₂₄H₂₅NO₅: C, 70.74; H, 6.18. Found: C, 70.69; H, 6.05.

(4*S*,5*R*,6*S*)-6-((4*R*)-4-Ethyl-2-oxo-oxazolidin-3-yl)-4,5-diphenyl-5,6-dihydro-4*H*-pyran-2-carboxylic acid methyl ester endo β 13b. According to the general procedure B, **endo β 13b** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (72%). mp 76-78 °C. ¹H NMR (400 MHz, CDCl₃) δ TMS: 7.17 (6H, m), 6.92 (2H, m), 6.85 (2H, m), 5.76 (sl, 1H), 4.16 (sl, 1H), 3.83 (s, 3H), 3.77 (2H, m), 3.44 (sl, 1H), 1.76 (m, 1H), 1.42 (sl, 1H), 0.72 (t, 3H, *J* = 6.5 Hz). ¹³C NMR (100 MHz, CDCl₃) δ : 162.4, 157.6, 143.6, 140.5, 138.0, 128.4, 128.3, 128.0, 127.1, 127.0, 115.4,

85.4, 67.2, 55.6, 49.6, 48.7, 26.4, 7.8. $[\alpha]_D^{20}$ -145.2 (c 1, acetone). IR (KBr): 1731, 1651, 1602, 1269, 1109 cm^{-1} . HRMS calcd for $\text{C}_{24}\text{H}_{25}\text{NO}_5$ $[\text{M} + \text{Na}]^+$ 430.1630, found 430.1627.

6-((4*R*)-4-Ethyl-2-oxo-oxazolidin-3-yl)-4,5-diphenyl-5,6-dihydro-4*H*-pyran-2-carboxylic acid methyl ester *exo* 13b. According to the general procedure B, **exo 13b** was identified from the crude mixture. ^1H NMR (400 MHz, CDCl_3) δ TMS: 7.12 (6H, m), 6.74 (2H, m), 6.60 (2H, sl), 6.28 (1H, d, J = 5.8 Hz), 6.05 (1H, d, J = 11.1 Hz), 3.84 (3H, s), 3.82 (2H, m), 3.73 (1H, dd, J = 8.3 and 8.7 Hz), 3.61 (1H, dd, J = 6.1 and 11.1 Hz), 3.46 (1H, m), 1.77 (2H, m), 0.91 (3H, t, J = 7.4 Hz).

(4*S*,5*R*,6*R*)-methyl 6-((*R*)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-5-methyl-4-phenyl-4*H*-pyran-2-carboxylate *endo* α 13c: According to the general procedure A, **13c** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (62%). mp 106-108 $^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3) δ TMS: 7.30 (5H, m), 6.02 (1H, d, J = 2.5 Hz), 5.40 (1H, d, J = 10.4 Hz), 4.38 (1H, dd, J = 8.3 and 8.4 Hz), 4.09 (1H, dd, J = 5.4 and 8.4 Hz), 3.89 (1H, m), 3.77 (3H, s), 3.30 (1H, dd, J = 2.5 and 10.1 Hz), 2.17 (1H, m), 1.88 and 1.79 (2H, 2m), 0.98 (3H, t, J = 7.5 Hz), 0.90 (3H, d, J = 6.6 Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 162.6, 157.4, 143.0, 141.3, 128.7, 128.1, 127.4, 114.4, 85.8, 67.1, 54.3, 52.1, 47.0, 36.0, 30.9, 14.2, 8.0. $[\alpha]_D^{20}$ +40.1 (1M, acetone). IR (KBr): 1735, 1650, 1494, 1259, 1056 cm^{-1} . Anal. Calcd for: C, 66.07 ; H, 6.71. Found: C, 65.61; H, 6.58.

(4*R*,5*S*,6*S*)-methyl 6-((*R*)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-5-methyl-4-phenyl-4*H*-pyran-2-carboxylate *endo* β 13c : According to the general procedure B, **13c** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (71%). mp 146-148 $^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3) δ TMS: 7.30 (3H, m), 7.18 (2H, m), 6.05 (1H, d, J = 2.4 Hz), 5.36 (1H, l), 4.47 (1H, dd, J = 6.4 and 8.3 Hz), 4.14 (1H, m), 3.89 (1H, dd, J = 6.4

and 8.3 Hz), 3.79 (3H, s), 3.28 (1H, dd, $J = 2.4$ and 10.1 Hz), 2.16 (1H, m), 1.90 and 1.59 (2H, 2m), 0.95 (3H, d, $J = 6.6$ Hz), 0.93 (3H, t, $J = 7.5$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 157.3, 143.0, 141.3, 128.6, 128.1, 127.2, 115.1, 86.9, 67.5, 54.5, 52.1, 47.2, 38.0, 26.7, 14.1, 8.6. $[\alpha]^{20}_{\text{D}} -77.5$ (1M, acetone). IR (KBr): 1735, 1650, 1494, 1259, 1056 cm^{-1} . HRMS calcd for $\text{C}_{19}\text{H}_{23}\text{NO}_5\text{Na}$ $[\text{M}+\text{Na}]^+$ 368.1474, found 368.1474.

(4*S*,5*R*,6*R*)-methyl 6-((*R*)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-4-(3,4,5-trimethoxyphenyl)-5-methyl-4*H*-pyran-2-carboxylate endo α 13d.

According to the general procedure A, **13d** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (64%). mp 127-128 °C. ^1H NMR (400 MHz, CDCl_3) δ TMS: 6.37 (2H, s), 6.01 (1H, d, $J = 2.5$ Hz), 5.34 (1H, d, $J = 10.4$ Hz), 4.40 (1H, t, $J = 8.6$ Hz), 4.10 (1H, dd, $J = 5.4$ and 8.7 Hz), 3.94 (1H, m), 3.86 (6H, s), 3.84 (3H, s), 3.79 (3H, s), 3.23 (1H, dd, $J = 2.5$ and 10.1 Hz), 2.21 (1H, m), 1.88 and 1.80 (2H, 2m), 0.99 (3H, t, $J = 7.5$ Hz), 0.93 (3H, d, $J = 6.6$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 162.5, 157.4, 153.3, 142.9, 137.2, 136.9, 114.3, 105.10, 86.0, 67.1, 60.8, 56.2, 54.4, 52.1, 47.5, 35.8, 26.7, 14.3, 8.0. $[\alpha]^{20}_{\text{D}} -24.4$ (c 1, acetone). IR (KBr): 2936, 1731, 1650, 1259, 1121 cm^{-1} . HRMS calcd for $\text{C}_{22}\text{H}_{29}\text{NO}_8$ $[\text{M}^+]$ 435.1893, found 435.1850.

(4*R*,5*S*,6*S*)-methyl 6-((*R*)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-4-(3,4,5-trimethoxyphenyl)-5-methyl-4*H*-pyran-2-carboxylate endo β 13d.

According to the general procedure B, **13d** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (67%). mp 70.9-72.9 °C. ^1H NMR (400 MHz, CDCl_3) δ TMS: 6.37 (2H, s), 6.04 (1H, d, $J = 2.5$ Hz), 5.30 (1H, m), 4.48 (1H, t, $J = 8.3$ Hz), 4.13 (1H, m), 4.06 (1H, dd, $J = 6.5$ and 8.3 Hz), 3.86 (6H, s), 3.84 (3H, s), 3.80 (3H, s), 3.20 (1H, dd, $J = 2.5$ and 10.0 Hz), 2.21 (1H, m), 1.89 and 1.61 (2H, 2m), 0.97 (3H, d, $J = 6.6$ Hz), 0.95 (3H, t, $J = 7.2$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 158.3, 153.2, 143.0,

137.0, 136.9, 114.9, 105.0, 87.0, 67.5, 60.7, 56.1, 55.1, 52.1, 47.7, 37.7, 26.7, 14.2, 8.5. $[\alpha]^{20}_D$ -85.7 (c 1, acetone). IR (KBr): 2936, 1731, 1650, 1259, 1121 cm^{-1} . Anal. Calcd for $\text{C}_{22}\text{H}_{29}\text{NO}_8$: C, 60.68; H, 6.71. Found: C, 60.29; H, 6.62.

(4*S*,5*S*,6*R*)-methyl 6-((*R*)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-4-(3,4,5-trimethoxyphenyl)-5-phenyl-4*H*-pyran-2-carboxylate endo α 13e.

According to the general procedure A, **13e** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (59%). mp 159-162°C. ^1H NMR (400 MHz, CDCl_3) δ TMS: 7.23 (3H, m), 6.94 (2H, d, J = 6.0 Hz), 6.28 (1H, d, J = 2.5 Hz), 5.97 (1H, d, J = 10.6 Hz), 5.95 (2H, s), 3.85 (3H, s), 3.84 (2H, m), 3.78 (3H, s), 3.74 (1H, dd, J = 2.5 and 10.5 Hz), 3.66 (1H, m), 3.63 (6H, s), 3.06 (1H, t, J = 10.5 Hz), 1.81 and 1.70 (2H, 2m), 0.90 (3H, t, J = 7.5 Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 157.3, 152.9, 144.0, 137.5, 136.9, 135.7, 128.7, 128.0, 127.5, 114.4, 104.7, 84.3, 66.9, 60.7, 55.9, 54.1, 52.3, 49.3, 48.7, 26.8, 7.9. $[\alpha]^{20}_D$: +132.2 (c 1, acetone). IR (KBr): 1753, 1728, 1640, 1494, 1260, 1126, 1105 cm^{-1} . Anal. Calcd for $\text{C}_{27}\text{H}_{31}\text{NO}_8$: C, 65.18; H, 6.28. Found: C, 64.97; H, 6.18.

(4*R*,5*R*,6*S*)-methyl 6-((*R*)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-4-(3,4,5-trimethoxyphenyl)-5-phenyl-4*H*-pyran-2-carboxylate endo β 13e.

According to the general procedure B, **13e** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (57%). mp 198.5-199.2 °C. ^1H NMR (400 MHz, CDCl_3) δ TMS: 7.21 (3H, m), 6.95 (2H, d, J = 5.8 Hz), 6.28 (1H, d, J = 2.3 Hz), 5.98 (6H, s), 5.76 (1H, sl), 4.17 (1H, sl), 3.84 (3H, s), 3.82 (1H, m), 3.77 (3H, s), 3.69 (1H, dd, J = 10.7 and 2.3 Hz), 3.64 (6H, s), 3.36 (1H, sl), 1.75 (1H, sl), 1.43 (1H, sl), 0.73 (3H, sl). ^{13}C NMR (100 MHz, CDCl_3) δ : 162.4, 157.3, 152.8, 144.0, 138.1, 136.9, 135.9, 128.5, 128.4, 127.2, 115.0, 104.9, 85.4, 67.2, 60.8, 56.0, 52.3, 49.0, 48.7, 26.4, 7.8. $[\alpha]^{20}_D$: -167.9 (c 1, acetone). IR (KBr): 1737, 1724, 1591, 1280, 1125, 993

cm⁻¹. Anal. Calcd for C₂₇H₃₁NO₈: C, 65.18 ; H, 6.28. Found: C, 65.67 ; H, 6.24.

methyl 6-((R)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-4-(3,4,5-trimethoxyphenyl)-5-phenyl-4H-pyran-2-carboxylate exo 13e. According to the general procedure B, **exo 13e** was identified from the crude mixture. ¹H NMR (400 MHz, CDCl₃) δ TMS: 7.14 (3H, m), 6.74 (2H, m), 6.66 (2H, sl), 6.27 (1H, d, *J* = 5.5 Hz), 6.06 (1H, d, *J* = 11.1 Hz), 5.88 (2H, s), 3.86 (3H, s), 3.84 (2H, m), 3.78 (3H, s), 3.76 (1H, m), 3.59 (1H, m), 3.58 (6H, s), 3.47 (m, 1H), 1.78 (2H, m), 0.92 (3H, t, *J* = 7.3 Hz).

(4S,5R,6R)-methyl 5-((benzo[d][1,3]dioxol-5-yl)methyl)-6-((R)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-4-(3,4,5-trimethoxyphenyl)-4H-pyran-2-carboxylate endo α 13f. According to the general procedure A, **13f** was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (65%). mp 74-76 °C. ¹H NMR (400 MHz, CDCl₃) δ TMS: 6.65 (1H, d, *J* = 7.9 Hz), 6.45 (1H, dd, *J* = 1.5 and 7.9 Hz), 6.40 (2H, s), 6.35 (1H, d, *J* = 1.5 Hz), 5.99 (1H, d, *J* = 2.5 Hz), 5.89 and 5.88 (2H, 2d, *J* = 1.5 Hz), 5.60 (1H, d, *J* = 9.8 Hz), 3.88 (6H, s), 3.85 (3H, s), 3.83 (1H, m), 3.77 (3H, s), 3.64 (1H, t, *J* = 8.4 Hz), 3.36 (1H, dd, *J* = 2.5 and 9.8 Hz), 3.28 (1H, m), 2.77 (1H, dd, *J* = 2.0 and 13.8 Hz), 2.48 (1H, dd, *J* = 7.9 and 13.8 Hz), 2.40 (1H, ddd, *J* = 2.0, 7.9 and 9.8 Hz), 1.77 (1H, sext, *J* = 7.4 Hz), 1.67 (1H, dq, *J* = 3.4 and 7.4 Hz), 0.91 (3H, t, *J* = 7.4 Hz). ¹³C NMR (100 MHz, CDCl₃) δ: 162.4, 157.5, 153.4, 147.5, 146.0, 142.2, 137.2, 136.7, 132.7, 121.4, 114.4, 108.8, 108.1, 105.3, 100.9, 85.5, 67.0, 60.8, 56.2, 54.5, 52.1, 46.1, 42.3, 35.7, 26.8, 7.7. [α]_D²⁰: +61.8 (c 1, acetone). IR (KBr): 1735, 1240, 1123 cm⁻¹. HRMS calcd for C₂₉H₃₃NO₁₀Na [M+Na]⁺ 578.2002, found 578.1984.

(4R,5S,6S)-methyl 5-((benzo[d][1,3]dioxol-5-yl)methyl)-6-((R)-4-ethyl-2-oxo-oxazolidin-3-yl)-5,6-dihydro-4-(3,4,5-trimethoxyphenyl)-4H-pyran-2-carboxylate endo β 13f. According to the general procedure B, **13f**

was obtained after column chromatography (toluene/AcOEt 95/5) as small needles (61%). mp 75-77 °C. ^1H NMR (400 MHz, CDCl_3) δ TMS: 6.61 (1H, d, $J = 7.9$ Hz), 6.45 (1H, dd, $J = 1.5$ and 7.9 Hz), 6.39 (1H, d, $J = 1.5$ Hz), 6.34 (2H, s), 5.98 (1H, d, $J = 7.9$ Hz), 5.88 and 5.87 (2H, 2d, $J = 1.3$ Hz), 5.25 (1H, d, $J = 9.6$ Hz), 4.33 (1H, t, $J = 8.3$ Hz), 3.98 (1H, m), 3.85 (6H, s), 3.84 (3H, s), 3.83 (1H, m), 3.77 (3H, s), 3.33 (1H, dd, $J = 2.5$ and 10.0 Hz), 2.76 (1H, m), 2.74 (1H, dd, $J = 4.2$ and 14.6 Hz), 2.65 (1H, dd, $J = 6.6$ and 14.6 Hz), 1.77 and 1.26 (2H, 2m), 0.90 (3H, t, $J = 7.4$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ 162.3, 157.7, 153.3, 147.3, 145.9, 142.0, 137.0, 136.6, 132.2, 121.9, 115.6, 109.4, 107.7, 105.1, 100.9, 86.0, 67.5, 60.7, 56.2, 56.1, 52.1, 45.7, 42.6, 37.9, 26.0, 8.7. $[\alpha]_D^{20}$: -110.8 (1M, acetone). IR (KBr): 1734, 1236, 1121, 1086 cm^{-1} . HRMS calcd for $\text{C}_{29}\text{H}_{33}\text{NO}_{10}\text{Na}$ $[\text{M}+\text{Na}]^+$ 578.2002, found 578.2007.

Copy of ^1H NMR: Figure S-1 (**8d**), Figure S-2 (**8e**), Figure S-3 (**9e**), Figure S-4 (**9d**), Figure S-5 (**13b endo β**), Figure S-6 (**13b exo**), Figure S-7 (**13d endo α**), Figure S-8 (**13e exo**), Figure S-9 (**13f endo α**), Figure S-10 (**13f endo β**), Figure S-11 (**13c endo β**).

Copy of ^{13}C NMR: Figure S-12 (**13c endo β**), Figure S-13 (**13b endo β**), Figure S-14 (**13d endo α**), Figure S-15 (**13f endo α**), Figure S-16 (**13f endo β**).

Figure S-1 : (8d)

Current Data Parameters
NAME Jan17-2006
EXPNO 40
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060117
Time 14.42
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 128
DW 60.400 use
DE 10.00 use
TE 295.8 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.1300030 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-2 : (8e)

Current Data Parameters
NAME Jan25-2006
EXPNO 40
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060125
Time 14.46
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 32
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 128
DW 60.400 use
DE 10.00 use
TE 294.3 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.1300077 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-3 : (9e)

Current Data Parameters
NAME Jan30-2006
EXPNO 100
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060130
Time 15.42
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 228.1
DW 60.400 use
DE 10.00 use
TE 296.9 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.1300083 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-4 : (9d)

Current Data Parameters
NAME Jan27-2006
EXPNO 170
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060127
Time 15.09
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDC13
NS 16
DS 2
SWH 4139.073 Hz
FIDRES 0.063157 Hz
AQ 7.9167986 sec
RG 11585.2
DW 120.800 use
DE 10.00 use
TE 300.0 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 9.50 use
PL1 -6.00 dB
SFO1 200.1312359 MHz

F2 - Processing parameters
SI 32768
SF 200.1300070 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-5 : (13b endo beta)

Current Data Parameters
NAME May04-2006
EXPNO 40
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060504
Time 15.08
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDC13
NS 32
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 456.1
DW 60.400 use
DE 10.00 use
TE 296.3 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.1300090 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-6 : (13b exo)

Current Data Parameters
NAME Apr27-2006
EXPNO 20
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060427
Time 11.45
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDC13
NS 16
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 128
DW 60.400 use
DE 10.00 use
TE 295.2 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.1300088 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-7 : (13d endo alpha)

Current Data Parameters
NAME Dec14-2005
EXPNO 50
PROCNO 1

F2 - Acquisition Parameters
Date_ 20051214
Time 15.14
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDC13
NS 32
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 322.5
DW 60.400 use
DE 10.00 use
TE 295.1 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.130073 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-8 ; (13e exo)

Current Data Parameters
NAME Apr11-2006
EXPNO 90
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060411
Time 14.46
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDC13
NS 16
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 128
DW 60.400 use
DE 10.00 use
TE 296.3 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.130064 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-9 : (13f endo alpha)

13f endo α

Current Data Parameters
NAME Feb09-2006
EXPNO 90
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060209
Time 17.30
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDC13
NS 32
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 406.4
DW 60.400 use
DE 10.00 use
TE 297.8 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.1300080 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-10 : (13f endo beta)

Current Data Parameters
NAME Feb09-2006
EXPNO 80
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060209
Time 17.21
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 32
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 406.4
DW 60.400 use
DE 10.00 use
TE 298.2 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.1300081 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-11 : (13c endo beta)

Current Data Parameters
NAME Jan20-2006
EXPNO 20
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060120
Time 14.26
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 32
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 287.4
DW 60.400 use
DE 10.00 use
TE 296.2 K
D1 1.00000000 sec
TD0 1

===== CHANNEL f1 =====
NUC1 1H
P1 10.00 use
PL1 -4.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.130083 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Figure S-12 : (13c endo beta)

Current Data Parameters
NAME Feb27-2006
EXPNO 140
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060227
Time 20.08
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TD 65536
SOLVENT CDC13
NS 1536
DS 2
SWH 23980.814 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 1149.4
DW 20.850 use
DE 10.00 use
TE 295.5 K
D1 2.00000000 sec
d11 0.03000000 sec
DELTA 1.89999998 sec
TD0 1

===== CHANNEL f1 =====
NUC1 13C
P1 9.00 use
PL1 -1.00 dB
SFO1 100.6228298 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 use
PL2 -4.00 dB
PL12 14.00 dB
PL13 14.00 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127690 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Figure S-13 : (13b endo beta)

Current Data Parameters
NAME Jul06-2006
EXPNO 130
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060706
Time 13.43
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TD 65536
SOLVENT CDC13
NS 1536
DS 4
SWH 11990.407 Hz
FIDRES 0.182959 Hz
AQ 2.7329011 sec
RG 456.1
DW 41.700 use
DE 10.00 use
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
DELTA 1.89999998 sec
TD0 1

===== CHANNEL f1 =====
NUC1 13C
P1 6.00 use
PL1 -6.00 dB
SFO1 50.3277608 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 use
PL2 -6.00 dB
PL12 17.00 dB
PL13 20.00 dB
SFO2 200.1308005 MHz

F2 - Processing parameters
SI 32768
SF 50.3227290 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Figure S-14 : (13d endo alpha)

Current Data Parameters
NAME Mar06-2006
EXPNO 50
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060306
Time 13.34
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TD 65536
SOLVENT CDC13
NS 2048
DS 2
SWH 23980.814 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 1149.4
DW 20.850 use
DE 10.00 use
TE 296.6 K
D1 2.00000000 sec
d11 0.03000000 sec
DELTA 1.89999998 sec
TD0 1

===== CHANNEL f1 =====
NUC1 13C
P1 9.00 use
PL1 -1.00 dB
SFO1 100.6228298 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 use
PL2 -4.00 dB
PL12 14.00 dB
PL13 14.00 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127745 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Figure S-15 : (13f endo alpha)

Current Data Parameters
NAME Feb13-2006
EXPNO 130
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060214
Time 8.09
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TD 65536
SOLVENT CDC13
NS 2048
DS 2
SWH 23980.814 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 362
DW 20.850 use
DE 10.00 use
TE 294.1 K
D1 2.00000000 sec
d11 0.03000000 sec
DELTA 1.89999998 sec
TD0 1

===== CHANNEL f1 =====
NUC1 13C
P1 9.00 use
PL1 -1.00 dB
SFO1 100.6228298 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 use
PL2 -4.00 dB
PL12 14.00 dB
PL13 14.00 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127690 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Figure S-16 : (13f endo beta)

Current Data Parameters
NAME Feb14-2006
EXPNO 40
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060214
Time 12.59
INSTRUM spect
PROBHD 5 mm QNP 1H/13
PULPROG zgpg30
TD 65536
SOLVENT CDC13
NS 1536
DS 2
SWH 23980.814 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 287.4
DW 20.850 use
DE 10.00 use
TE 294.8 K
D1 2.00000000 sec
d11 0.03000000 sec
DELTA 1.89999998 sec
TD0 1

===== CHANNEL f1 =====
NUC1 13C
P1 9.00 use
PL1 -1.00 dB
SFO1 100.6228298 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 use
PL2 -4.00 dB
PL12 14.00 dB
PL13 14.00 dB
SFO2 400.1316005 MHz

F2 - Processing parameters
SI 32768
SF 100.6127690 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40