Supporting Information

Low Temperature *n*-Butyllithium-Induced Rearrangement of Allyl 1,1-Dichlorovinyl Ethers.

Aaron Christopher, Dahniel Brandes, Stephen Kelly, and Thomas Minehan

*Department of Chemistry, California State University–Northridge,
Northridge, California 91330*

Table of Contents:

General Methods and Procedures	S2-S3
Characterization data for new compounds	S3-S16
¹ H NMR spectra of compounds	S17-S38
NOESY spectrum of 8	S39
¹ H NMR spectra of compounds 3am , 3an	S40-S41

General Methods. All reactions were conducted in flame-dried glassware under balloon pressure of argon. Reaction temperatures refer to external bath temperatures. Tetrahydrofuran was dried by distillation from the sodium ketyl of benzophenone. Liquid reagents or reagent solutions were added by syringe or cannula. Organic extracts were dried over Na_2SO_4 , filtered, and concentrated using a rotary evaporator at aspirator pressure (20-30mmHg). Chromatography refers to flash chromatography and was carried out on SiO_2 (silica gel 60, 230-400 mesh) according to Still's method. ^1H - and ^{13}C NMR spectra were recorded on a Varian DPX-400 spectrometer using Me_4Si as internal standard. Chemical shifts are reported in ppm downfield (δ) from Me_4Si . Only noteworthy IR absorptions are listed. GC-MS spectra were obtained using a Hewlett Packard 5890-5971 GC-Mass spectrometer.

Representative Procedure for the Preparation of Allyl Dichlorovinyl Ethers from Allylic Alcohols:

To a stirred solution of allylic alcohol (1 mmol) in pyridine (5 mL) at 0°C under argon was added freshly prepared acetic formic anhydride* dropwise until tlc indicated complete disappearance of the alcohol. The solution was diluted with ether (100 mL) and washed with saturated NaHCO_3 solution (2 x 50 mL) and saturated copper sulfate solution (2x 50 mL). The organic layer was dried over Na_2SO_4 , filtered, and evaporated to dryness. The crude formate was taken up in THF (10 mL) and PPh_3 (786 mg, 3 mmol, 3 eq.) was added. The solution was warmed to 60°C under argon and CCl_4 (1.0 mL, 10 mmol, 10 eq) was added slowly dropwise (via addition funnel) over 8-10 hours. The red-brown solution was cooled to room temperature, diluted with ether (100 mL) and washed with saturated NaHCO_3 solution (2 x 50 mL). The ether extract was dried over Na_2SO_4 , filtered, and evaporated to dryness. Flash chromatography (hexanes) afforded the pure allyl dichlorovinyl ether.

*Acetic formic anhydride was prepared by stirring a solution of formic acid (8.6 mL of a 96% aqueous solution) and acetic anhydride (20.4 mL) at 60°C in a flask equipped with a drying tube. The solution could be used directly after cooling to room temperature. See main text ref. # 10.

Representative Procedure for the Rearrangement of Allyl Dichlorovinyl Ethers: To a stirred solution of allyl dichlorovinyl ether (0.3 mmol) in THF (2mL) under argon at -78°C is added $n\text{BuLi}$ (1.6 M in hexanes, 0.42 mL, 0.66 mmol, 2.2 eq.). The solution was stirred at -78°C for 0.5 h and was then warmed to -42°C (dry ice-acetonitrile bath) for 0.5 h. Excess ethanol (0.5 mL) was then added and the reaction was warmed to room temperature. The solution was diluted with ether (50 mL) and washed with saturated NaHCO_3 solution (2 x 25 mL). The ethereal extract was dried over Na_2SO_4 , filtered and concentrated in vacuo. Purification of the residue by flash chromatography afforded the rearrangement products.

Representative Procedure for Rearrangement with α -substitution:

To a stirred solution of allyl dichlorovinyl ether (0.31 mmol) in THF (2.5 mL) under argon at -78°C is added *n*-BuLi (1.6 M in hexanes, 0.42 mL, 0.66 mmol, 2.2 eq.). The solution was stirred at -78°C for 0.5 h and was then warmed to -42°C (dry ice-acetonitrile bath) for 0.5 h. The solution was cooled again to -78°C , and either TMSCl (0.08 mL, 0.65 mmol, 2 eq.) or methyl triflate (0.07 mL, 0.65 mmol, 2 eq.) was added. The solution was stirred at -78°C for 15 minutes, and was then warmed to -40°C (dry ice-acetonitrile bath) for 0.5 h. Excess ethanol (0.5 mL) was then added and the reaction was warmed to room temperature. The solution was diluted with ether (50 mL) and washed with saturated NaHCO_3 solution (2 x 25 mL). The ethereal extract was dried over Na_2SO_4 , filtered and concentrated in vacuo. Purification of the residue by flash chromatography afforded the rearranged α -substitution products.

Note: none of the dichlorovinyl ethers (2a-2f, 5a, 5b, 7, 9) gave satisfactory results on exact mass analysis due to degradation processes occurring in the mass spectrometer. A possible mechanism may involve thermal rearrangement (injector temperatures $>200^{\circ}\text{C}$) of the allyl-dichlorovinyl ether and subsequent decomposition of the resultant aldehyde. (see reference 11 in main text for thermal rearrangement of allyl 1,1-dichlorovinyl ethers)

We supply instead low-resolution mass analysis and GC data for these compounds.

Purification: SiO_2 (hexanes) affording **2a** (85%) as a colorless oil.

See spectrum S-17

^1H NMR: (400 MHz, CDCl_3)

6.55 (s, 1H); 5.79 (dt, $J=6.8, 15.6$ Hz, 1H); 5.57 (dtt, $J=1.2, 6.4, 16.4$ Hz, 1H); 4.32 (d, $J=6.4$ Hz, 1H); 2.09 (q, $J=7.6$ Hz, 2H); 1.25-1.44 (m, 6H); 0.90 (t, $J=6.8$ Hz, 3H).

^{13}C NMR: (100 MHz, CDCl_3)

142.7; 138.1; 124.6; 96.5; 74.3; 32.6; 31.7; 28.8; 22.8; 14.4.

LRMS: (CI) calculated for $\text{C}_{10}\text{H}_{16}\text{Cl}_2\text{O}$ 223.0651, found $m/z=223$ (MH^+)

GC: $t_r=11.35$ min. (injector 250°C , 10.5 psi, column gradient 100°C for 2 minutes, $25^{\circ}\text{C}/\text{min}$ to 250°C , 250°C for 10 minutes)

IR: 2956 cm^{-1} , 2927, 2857, 1640.

TLC: R_f 0.76 (SiO_2 , hexanes:EtOAc 10:1)

Purification: SiO₂(2% Et₂O in hexanes) affording **3a** (91%) as a pale yellow oil.

¹H NMR: (400 MHz, CDCl₃)
 5.63 (ddd, *J*=8.4, 10.0, 17.2 Hz, 1H); 5.04 (m, 1H); 4.97-5.00 (m, 1H); 4.13 (q, *J*=7.2 Hz, 2H); 2.50 (m, 1H); 2.34, 2.27 (dd, *J*=6.4, 14.4 Hz, 2H); 1.30 (m, 8H); 1.24 (t, *J*=7.2 Hz, 3H); 0.88 (t, *J*=7.2 Hz, 3H)

¹³C NMR: (100 MHz, CDCl₃)
 173.0; 141.6; 115.2; 60.5; 40.8; 40.5; 34.8; 32.1; 26.9; 22.9; 14.6; 14.4

HRMS: (CI) calculated for C₁₂H₂₃O₂ 199.1698, found *m/z*=199.1707 (MH⁺)

IR: 2957 cm⁻¹, 2928, 2857, 1737, 1640.

TLC: R_f 0.66 (SiO₂, hexanes:EtOAc 10:1)

For a previous synthesis of **3a**, see: Boldrini, G.P.; Mengoli, M.; Tagliavini, E.; Trombini, C.; Umani-Ronchi, A. *Tetrahedron Lett.* **1986**, 27, 4223.

See spectrum S-18

Purification: SiO₂(1% Et₂O in hexanes) affording **3am** (85%) as a pale yellow oil.

¹H NMR: (400 MHz, CDCl₃)
 5.62 (ddd, *J*=8.4, 10.4, 17.6 Hz, 1H); 5.04 (m, 1H); 5.00 (m, 1H); 3.64 (s, 3H); 2.53 (m, 1H); 2.36 (dd, *J*=8.4, 14.8 Hz, 2H); 1.33 (m, 8H); 0.90 (t, *J*=7.2 Hz, 3H)

GCMS: t_R=4.79 min. (Agilent 6890; injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes) Calculated for C₁₁H₂₀O₂ 184.1463, found *m/z*=184 (M⁺).

See spectrum S-40

Purification: SiO₂(1% Et₂O in hexanes) affording **3an** (72%) as a pale yellow oil

¹H NMR: (400 MHz, CDCl₃)

5.63 (ddd, *J*=8.4, 10.0, 17.2 Hz, 1H); 5.04 (m, 1H); 4.97-5.01 (m, 1H); 4.67 (tdd, *J*=2.0, 4.4, 10.8 Hz, 1H); 2.50 (m, 1H); 2.30 (dd, *J*=6.4, 14.4 Hz, 2H); 1.97 (m, 1H); 1.88 (m, 1H); 1.66 (m, 2H); 1.50-1.20 (m, 11H); 1.10-0.85 (m, 10H); 0.74 (dd, *J*= 4.0, 6.8 Hz, 3H).

GCMS: *t_R*=8.56 min. (Agilent 6890; injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes) Calculated for C₂₀H₃₆O₂ 308.2715, found *m/z*=308 (M⁺)

See spectrum S-41

((*E*)-4-(2,2-dichlorovinyl)oxy)but-2-enyloxy(*tert*-butyl)diphenylsilane 2b

Purification: SiO₂(hexanes) affording **2b** (80%) as a yellow oil.

¹H NMR: (400 MHz, CDCl₃)

7.62 (m, 4H); 7.34 (m, 6H); 6.48 (s, 1H); 5.81 (m, 2H); 4.33 (d, *J* = 4.0 Hz, 2H); 4.18 (d, *J* = 2.4 Hz, 2H); 1.02 (s, 9H).

¹³C NMR: (100 MHz, CDCl₃)

142.9; 135.9; 134.9; 133.8; 130.2; 128.2; 124.2; 104.7; 73.6; 63.8; 27.2; 19.7

LRMS: (CI⁺) calculated for C₂₂H₃₀Cl₂NO₂Si 438.1423, found *m/z*= 438 (MNH₄⁺)

GC: *t_R*=22.51 min (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 350°C, 350°C for 15 minutes)

IR: 2958 cm⁻¹, 2930, 2857, 1643, 1589.

TLC: *R_f* 0.55 (SiO₂, hexanes:EtOAc 10:1)

See spectrum S-19.

((Z)-4-(2,2-dichlorovinyl)oxy)but-2-en-1-yl tert-butyl diphenylsilane **2c**

Purification: SiO₂(hexanes) affording **2c** (85%) as a pale yellow oil

¹H NMR: (400 MHz, CDCl₃)

7.71 (m, 4H); 7.43 (m, 6H); 6.45 (s, 1H); 5.87 (qt, *J* = 6.0, 1.6 Hz, 1H); 5.60 (qt, *J* = 6.9, 2.0 Hz, 1H); 4.34 (d, *J* = 6.8 Hz, 2H); 4.28 (d, *J* = 6.0 Hz, 2H); 1.09 (s, 9H).

¹³C NMR: (100 MHz, CDCl₃)

142.8; 135.9; 134.2, 133.7, 130.3, 128.2, 96.6, 69.2, 60.8, 27.2, 19.6

LRMS: (CI⁺) calculated for C₂₂H₃₀Cl₂NO₂Si 438.1423, found *m/z* = 438 (MNH₄⁺)

GC: *t_R* = 22.36 min (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 350°C, 350°C for 15 minutes)

IR: 3070 cm⁻¹, 2929, 2856, 1643, 1589.

TLC: *R_f* 0.58 (SiO₂, hexanes:EtOAc 10:1)

See spectrum S-20.

3b/c

Purification: SiO₂(95:5 hexanes/ether) affording **3b/3c** (90%) as a colorless oil

¹H NMR: (400 MHz, CDCl₃)

7.68 (d, *J* = 7.6 Hz, 4H); 7.41 (m, 6H); 5.79 (ddd, *J* = 17.6, 10.4, 8.0 Hz, 1H); 5.09 (m, 2H); 4.14 (q, *J* = 7.2 Hz, 2H); 3.70 (dd, *J* = 9.6, 5.2 Hz, 1H); 3.60 (dd, *J* = 9.6, 6.4 Hz, 1H); 2.86 (m, 1H); 2.71 (dd, *J* = 15.6, 6.0 Hz, 1H); 2.40 (dd, *J* = 15.2, 8.4 Hz, 1H); 1.25 (t, *J* = 7.2 Hz, 3H); 1.09 (s, 9H)

¹³C NMR: (100 MHz, CDCl₃)

172.9; 138.5; 136.0; 133.9; 130.0; 128.1; 116.6; 96.5; 66.8; 60.6; 43.1; 36.6; 27.3; 19.7; 14.6

HRMS: (CI) calculated for C₂₄H₃₃O₃Si 397.2198, found *m/z* = 397.2206 (MH⁺)

IR: 2930 cm^{-1} , 2857, 1731, 1641, 1587.

TLC: R_f 0.50 (SiO_2 , hexanes:EtOAc 10:1)

For a previous synthesis of **3b/c**, see: Sabol, J.S.; Flynn, G.A.; Friedrich, D.; Huber, E.W. *Tetrahedron Lett.* **1997**, 38, 3687.

See spectrum S-21.

3-(2,2-dichlorovinyl)oxy)cyclohex-1-ene 2d

Purification: SiO_2 (hexanes) affording **2d** (75%) as a pale yellow oil.

¹H NMR: (400 MHz, CDCl_3)

6.56 (s, 1H); 5.91 (dt, $J = 10.4, 3.6$ Hz, 1H); 5.65 (dq, $J = 11.2, 2.0$ Hz, 1H); 4.28 (m, 1H); 1.96 (m, 2H); 1.78 (m, 1H); 1.70 (m, 2H); 1.52 (m, 1H)

¹³C NMR: (100 MHz, CDCl_3)

142.2, 133.9, 125.9, 104.4, 77.0, 29.3, 25.3, 19.1

LRMS: (CI) calculated for $\text{C}_8\text{H}_{14}\text{Cl}_2\text{NO}$ 210.0452, found $m/z=210$ (MNH_4^+)

GC: $t_r=9.63$ min. (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes)

IR: 3031 cm^{-1} , 2939, 1640

TLC: R_f 0.71 (SiO_2 , hexanes:EtOAc 10:1)

See spectrum S-22.

ethyl 2-(cyclohex-2-enyl)acetate 3d

Purification: SiO_2 (95:5 hexanes/ether) affording **3d** (92%) as a colorless oil.

¹H NMR: (400 MHz, CDCl_3)

5.57 (dq, $J=2.0, 10.0$ Hz, 1H); 5.41 (dq, $J = 2.0, 12.0$ Hz, 1H); 4.01 (q, $J = 7.2$ Hz, 2H); 2.46 (m, 1H); 2.14 (dq, $J= 6.8, 15.2$ Hz, 2H); 1.85 (m, 2H); 1.70 (m, 1H); 1.58 (m, 1H); 1.44 (m, 1H); 1.16 (m, 1H); 1.14 (t, $J = 7.2$ Hz, 3H)

¹³C NMR: (100 MHz, CDCl_3)

172.9, 130.5, 128.3, 60.4, 41.2, 32.6, 29.2, 25.4, 21.4, 14.6

LRMS: (EI+) calculated for $\text{C}_{10}\text{H}_{16}\text{O}_2$ 168.1150, found $m/z= 168.0$ (M^+)

GC: $t_r=4.89$ min. (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes)

IR: 3019 cm^{-1} , 2980, 2930, 2862, 1736, 1648

TLC: R_f 0.51 (SiO_2 , hexanes:EtOAc 10:1)

For previous routes to this compound, see: Jones, G.B.; Huber, R.S.; Chau, S. *Tetrahedron* **1993**, 49, 369, and references therein.
See spectrum S-23.

Purification: SiO₂(hexanes) affording **2e** (72%) as a yellow oil.

¹H NMR: (400 MHz, CDCl₃)
 6.54 (s, 1H); 5.31 (t, *J*=7.2 Hz, 1H); 4.41 (d, *J*=7.6 Hz, 2H); 2.15-2.21 (m, 4H); 1.58 (m, 6H).
¹³C NMR: (100 MHz, CDCl₃)
 148.4; 142.9; 115.9; 96.5; 69.1; 37.5; 29.5; 28.7; 28.2; 26.9
LRMS: (CI) calculated for C₁₀H₁₈Cl₂NO 238.0765, found *m/z*=238 (MNH₄⁺)
GC: t_r 10.93 min. (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes)
IR: 2931 cm⁻¹, 2854, 1640.
TLC: R_f 0.83 (SiO₂, hexanes:EtOAc 10:1)

See spectrum S-24.

Purification: SiO₂(95:5 hexanes/ether) affording **3e** (85%) as a colorless oil.

¹H NMR: (400 MHz, CDCl₃)
 5.69 (dd, *J* = 17.6, 10.8 Hz, 1H); 5.01 (dt, *J* = 1.2, 11.2 Hz, 1H); 4.90 (dt, *J* = 1.2, 17.6 Hz, 1H); 3.99 (q, *J* = 7.2 Hz, 2H); 2.22 (s, 2H); 1.57 (m, 2H); 1.40 (m, 7H); 1.26 m, 1H); 1.14 (t, *J* = 7.2 Hz, 3H)
¹³C NMR: (100 MHz, CDCl₃)
 172.0, 145.3, 113.5, 60.2, 46.2, 39.6, 36.0, 26.6, 22.5, 14.7
HRMS: (EI+) calculated for C₁₂H₂₀O₂ 196.1463, found *m/z*= 196.1469 (M⁺)
IR: 3082 cm⁻¹, 2979, 2931, 2856, 1738, 1732, 1638
TLC: R_f 0.63 (SiO₂, hexanes:EtOAc 10:1)
 For a previous preparation of **3e**, see: Srikrishna, A.; Nagaraju, S.; Kondaiah, P. *Tetrahedron* **1995**, 51, 1809.

See spectrum S-25.

(Z)-1-(2,2-dichlorovinyloxy)-3,7-dimethylocta-2,6-diene 2f

Purification: SiO₂(hexanes) affording **2f** (93%) as a pale yellow oil.

¹H NMR: (400 MHz, CDCl₃)
 6.46 (s, 1H); 5.29 (t, *J* = 6.8 Hz, 1H); 5.01 (t, *J* = 6.8 Hz, 1H); 4.33 (d, *J* = 7.2 Hz, 2H); 2.03 (m, 4H); 1.63 (s, 3H); 1.62 (s, 3H); 1.54 (s, 3H)

¹³C NMR: (100 MHz, CDCl₃)
 143.6, 142.9, 132.3, 123.9, 119.2, 104.2, 69.9, 40.0, 26.6, 26.1, 18.1, 17.0

LRMS: (CI) calculated for C₁₂H₂₂Cl₂NO 266.1078, found *m/z*=266 (MNH₄⁺)

GC: *t*_r=13.23 min. (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes)

IR: 2926 cm⁻¹, 1640.

TLC: R_f 0.69 (SiO₂, hexanes:EtOAc 10:1)

See spectrum S-26.

ethyl 3,7-dimethyl-3-vinyloct-6-enoate 3f

Purification: SiO₂(95:5 hexanes/ether) affording **3f** (75%) as a colorless oil.

¹H NMR: (400 MHz, CDCl₃)
 5.74 (dd, *J* = 17.2, 10.8 Hz, 1H); 4.99 (tq, *J* = 5.6, 1.2 Hz, 1H); 4.91 (m, 2H); 4.02 (q, *J* = 7.2 Hz, 2H); 2.23 (s, 2H); 1.83 (m, 2H); 1.58 (s, 3H); 1.50 (s, 3H); 1.33 (m, 2H); 1.16 (t, *J* = 7.2 Hz, 3H); 1.05 (s, 3H)

¹³C NMR: (100 MHz, CDCl₃)
 172.0, 146.0, 131.7, 124.9, 112.4, 60.3, 45.3, 41.0, 39.6, 26.1, 23.7, 23.3, 17.9, 14.7

HRMS: (CI) calculated for C₁₄H₂₅O₂ 225.1854, found *m/z*=225.1861 (MH⁺)

IR: 3083 cm⁻¹, 2968, 2928, 1736, 1732, 1639

TLC: R_f 0.56 (SiO₂, hexanes:EtOAc 10:1)

For a previous preparation of **3f**, see: Srikrishna, A.; Nagaraju, S.; Kondaiah, P. *Tetrahedron* **1995**, 51, 1809.

See spectrum S-27.

(4R,6R)-6-(2,2-dichlorovinyl)-1-methyl-4-(prop-1-en-2-yl)cyclohex-1-ene 5a

Purification: SiO₂(hexanes) affording **5a** (78%) as a pale yellow oil.

¹H NMR: (400 MHz, CDCl₃)
6.61 (s, 1H); 5.66 (m, 1H); 4.76 (d, *J*=6.0 Hz, 2H); 4.45 (m, 1H); 2.24 (m, 1H); 2.06-2.19 (m, 2H); 1.97 (m, 1H); 1.74 (s, 3H); 1.73 (q, *J*=1.2 Hz); 1.61 (qd, *J*=10.4, 12.8 Hz, 1H).

¹³C NMR: (100 MHz, CDCl₃)
148.4; 141.8; 133.2; 127.7; 110.0; 104.3; 82.6; 40.9; 35.1; 31.2; 20.8; 19.2.

LRMS: (EI+) calculated for C₁₂H₁₆Cl₂O 246.0578, found *m/z*=246 (M⁺)

GC: *t_r*=7.18 min. (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes)

IR: 2969 cm⁻¹, 2941, 2918, 2859, 1643

TLC: R_f 0.75 (SiO₂, hexanes)

See spectrum S-28.

ethyl 2-((1R,5S)-2-methyl-5-(prop-1-en-2-yl)cyclohex-2-enyl)acetate 6a

Purification: SiO₂(98:2 hexanes/ether) affording **6a** (85%) as a colorless oil.

¹H NMR: (400 MHz, CDCl₃)
5.48 (m, 1H); 4.69 (m, 2H); 4.14 (q, *J*=7.2 Hz, 2 H); 2.64 (dd, *J*=4.8, 16.0 Hz, 2H); 2.02-2.17 (m, 3H); 1.91 (dq, *J*=2.4, 12.4 Hz, 2H), 1.71 (s, 3H); 1.65 (q, *J*=1.2Hz, 3H); 1.26 (t, *J*=6.8 Hz, 3H); 1.25 (m, 1H)

¹³C NMR: (100 MHz, CDCl₃)
173.5; 150.0; 135.1; 123.9; 109.0; 60.6; 41.7; 39.4; 37.7; 35.4; 31.5; 21.3; 21.0; 14.6

HRMS: (EI+) calculated for C₁₄H₂₃O₂ 223.1698, found *m/z*=223.1704 (MH⁺)

IR: 2966 cm⁻¹, 2915, 1731, 1737, 1644.

TLC: R_f 0.60 (SiO₂, 20:1 hexanes/ethyl acetate)

The spectroscopic data reported above matched the literature data reported in reference 15b (main text) for cis ethyl ester 6a

See spectrum S-30.

(4R,6S)-6-(2,2-dichlorovinyl)-1-methyl-4-(prop-1-en-2-yl)cyclohex-1-ene 5b

Purification: SiO₂(hexanes) affording **5b** (71%) as a yellow oil.

¹H NMR: (400 MHz, CDCl₃)
6.64 (s, 1H); 5.76 (m, 1H); 4.77 (m, 1H); 4.73 (s, 1H); 4.11 (m, 1H); 2.38 (m, 1H); 2.18-2.25 (m, 1H); 2.01 (dq, *J*=2.4, 14.0 Hz, 1H); 1.87 (m, 1H); 1.78 (m, 3H); 1.75 (s, 3H); 1.59 (ddd, *J*=4.0, 12.8, 14 Hz, 1H).

¹³C NMR: (100 MHz, CDCl₃)
148.9; 142.9; 130.8; 128.8; 109.7; 105.2; 80.8; 35.5; 34.1; 31.2; 21.3; 21.2

LRMS: (EI+) calculated for C₁₂H₁₆Cl₂O 246.0578, found *m/z*=246.0 (M⁺)

GC: *t*_r=6.99 min. (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes)

IR: 2968 cm⁻¹, 2916, 1640

TLC: R_f 0.60 (SiO₂, hexanes)

See spectrum S-29.

ethyl 2-((1S,5S)-2-methyl-5-(prop-1-en-2-yl)cyclohex-2-enyl)acetate 6b

Purification: SiO₂(98:2 hexanes/ether) affording **6b** (72%) as a colorless oil.

¹H NMR: (400 MHz, CDCl₃)
5.43 (m, 1H); 4.69 (m, 2H); 4.14 (q, *J*= 6.8 Hz, 2H); 2.53 (dd, *J*=4.0, 15.6 Hz, 2H); 2.25 (dd, *J*=11.2, 15.6 Hz, 1H); 2.04-2.16 (m, 2H); 1.86 (m, 1H); 1.71 (s, 3H); 1.66 (s, 3H); 1.64 (m, 1H); 1.57 (td, *J*= 6.8, 12.4 Hz, 1H); 1.25 (t, *J*= 7.2 Hz, 3H)

¹³C NMR: (100 MHz, CDCl₃)
173.4; 149.7; 135.3; 123.3; 109.1; 60.6; 38.3; 36.5; 36.2; 32.6; 31.2; 22.1; 21.2; 14.6.

HRMS: (EI+) calculated for C₁₄H₂₃O₂ 223.1698, found *m/z*=223.1706 (MH⁺)

IR: 3082 cm⁻¹, 2965, 2917, 1736, 1732, 1644.

TLC: R_f 0.65 (SiO₂, 20:1 hexanes/ethyl acetate)

The spectroscopic data reported above matched the literature data reported in reference 15b (main text) for trans ethyl ester 6b.

See spectrum S-31.

Purification: SiO₂(95:5 hexanes/ethyl acetate) affording **7** (79%) as a pale yellow oil

¹H NMR: (400 MHz, CDCl₃)
7.26-7.38 (m, 10H); 6.71 (s, 1H); 6.51 (dd, $J=1.6, 6.4$ Hz, 1H); 4.76-4.79 (m, 2H); 4.66 (m, 2H); 4.59 (m, 2H); 4.03 (ddd, $J=2.8, 4.0, 9.6$ Hz, 1H); 3.77-3.93 (m, 3H).

¹³C NMR: (100 MHz, CDCl₃)
147.0; 141.6; 138.1; 137.9; 128.9; 128.8; 128.7; 128.6; 128.4; 128.3; 128.2; 105.3; 99.3; 96.5; 80.8; 74.6; 74.3; 68.5.

IR: 3063 cm⁻¹, 3030, 2869, 1640, 1600

TLC: R_f 0.65 (SiO₂, hexanes:EtOAc 10:1)

Due to extensive thermal decomposition occurring in the mass spectrometer, a low resolution mass spectrum of **7** could not be obtained; GC-MS analysis showed multiple decomposition products.

See spectrum S-32.

Purification: SiO₂(90:10 hexanes/ethyl acetate) affording **8** (65%) as a colorless oil

¹H NMR: (400 MHz, CDCl₃)
7.32 (m, 10 H); 5.95 (d, $J=10.0$ Hz, 1H); 5.82 (d, $J=10.4$ Hz, 1H); 4.46-4.65 (m, 5 H); 4.18 (q, $J=6.8$ Hz, 2 H); 4.05 (d, $J=3.2$ Hz, 1H); 3.77 (d, $J=9.6$ Hz, 1H); 3.67 (m, 2H); 2.66 (dd, $J=6.8, 15.6$ Hz, 1 H); 2.49 (dd, $J=6.8, 15.2$ Hz, 1H); 1.26 (t, $J=7.2$ Hz, 3 H).

¹³C NMR: (100 MHz, CDCl₃)

171.0; 138.8; 138.5; 128.7; 128.3; 128.2; 128.1; 127.9; 127.4; 77.8; 72.0;
71.6; 70.7; 60.9; 40.9; 14.6

HRMS: (ES-MS) calculated for $C_{24}H_{28}O_5Na$ 419.1834, found $m/z=419.1829$ ($M+Na^+$)

IR: 2856 cm^{-1} , 1734, 1600.

TLC: R_f 0.55 (SiO_2 , hexanes:EtOAc 10:1)

These data closely match those reported in the literature for **8**; see reference 16(main text)

See spectra S-33 and S-39.

9

Purification: SiO_2 (95:5 hexanes/ethyl acetate) affording **9** (87%) as a yellow oil

1H NMR: (400 MHz, $CDCl_3$)

6.75 (s, 1H); 6.40 (dd, $J = 6.4, 1.6$ Hz, 1H); 4.77 (dd, $J = 6.0, 2.0$ Hz, 1H);
4.49 (dt, $J = 7.6, 2.0$ Hz, 1H); 5.93 (ddd, $J = 10.0, 6.0, 4.0$ Hz, 1H); 5.68 (dd, $J = 10.0, 2.0$ Hz, 1H); 3.45 (m, 1H); 2.09 (m, 2H); 2.02 (m, 1H); 1.72 (m, 1H);
1.54 (m, 2H)

^{13}C NMR: (100 MHz, $CDCl_3$)

148.0; 141.0; 131.6; 130.9; 129.9; 129.7; 129.5; 129.1; 123.3 (q, $J = 4.0$ Hz);
42.1; 32.9; 25.3; 21.4

GCMS: $t_R=7.35$ min. (Agilent 6890; injector $250^\circ C$, 10.5 psi, column gradient $100^\circ C$ for 2 minutes, $25^\circ C/min$ to $250^\circ C$, $250^\circ C$ for 10 minutes) Calculated for $C_{11}H_{14}Cl_2O_4$ 281.0342, found $m/z=281$ (MH^+)

IR: 3022 cm^{-1} , 2930, 2859, 1652, 1614

TLC: R_f 0.55 (SiO_2 , hexanes:EtOAc 10:1)

See spectrum S-34.

10 a/b

1.4:1 mixture of diastereomers

Purification: SiO₂(5% ether in hexanes) affording **10a,b** (56%) as a colorless oil

¹H NMR: (400 MHz, CDCl₃)

7.67 (m, 8H); 7.42 (m, 12H); 5.83 (ddd, $J = 17.2, 10.8, 9.2$ Hz, 1H); 5.68 (ddd, $J = 17.2, 10.4, 9.2$ Hz, 1H); 5.09 (m, 4H); 4.09 (q, $J=7.2$ Hz, 4H) 3.69 (m, 4H); 2.80 (m, 2H); 2.68 (m, 1H); 2.41-2.47 (m, 1H); 1.22 (t, $J=6.8$ Hz, 6H); 1.15 (d, $J=6.8$ Hz, 3 H); 1.10 (d, $J=7.2$ Hz, 3H); 1.07 (s, 18H)

¹³C NMR: (100 MHz, CDCl₃)

175.8; 137.7; 136.8; 136.0; 134.0; 130.0; 128.0; 60.5; 49.9; 40.5; 27.3; 19.7; 14.7; 14.6.

LRMS: (CI) calculated for C₂₅H₃₅O₃Si 411.2355, found $m/z = 411$ (MH⁺)

GC: $t_r=21.75$ min. (injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 350°C, 350°C for 15 minutes)

IR: 3071 cm⁻¹, 2930, 2857, 1731, 1640, 1589.

TLC: R_f 0.70 (SiO₂, hexanes:EtOAc 20:1)

For a previous preparation of **10a/b** see: Boldrini, G.P.; Mengoli, M.; Tagliavini, E.; Trombini, C.; Umani-Ronchi, A. *Tetrahedron Lett.* **1986**, 27, 4223.

See spectrum S-35.

11 a/b

1.25:1 mixture of diastereomers

Purification: SiO₂(2% ether in hexanes) affording **11a,b** (73%) as a colorless oil

¹H NMR: (400 MHz, CDCl₃)

7.67 (m, 8H); 7.41 (m, 12H); 6.10 (dt, $J = 16.8, 9.6$ Hz, 1H); 5.77 (dt $J = 17.2, 10.0$ Hz 1H); 5.01-5.22 (m, 4H); 4.08 (m, 4H) 3.54-3.77 (m, 4H); 2.83 (m, 1H); 2.59 (m, 1H); 2.44 (d, $J=6.4$ Hz, 1H); 2.33 (d, $J=11.2$ Hz, 1H); 1.25 (t,

$J=7.6$ Hz, 3 H); 1.21 (t, $J=7.2$ Hz, 3H); 1.08 (s, 9H); 1.07 (s, 9H); 0.11 (s, 9H); 0.06 (s, 9H).

¹³C NMR: (100 MHz, CDCl₃)

181.5; 140.2; 136.1; 134.2; 134.1; 130.0; 128.0; 119.4; 117.2; 67.8; 60.1; 59.9; 46.4; 39.3; 38.2; 27.3; 19.7; 15.9; 14.8; -1.0

GCMS: $t_R=13.51$ min. (Agilent 6890; injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes); calculated for C₂₇H₄₀O₃Si₂ 468.2516, found $m/z=468.0$ (M⁺)

IR: 3071 cm⁻¹, 2957, 2930, 2857, 1716, 1640, 1589.

TLC: R_f 0.65 (SiO₂, hexanes:EtOAc 20:1)

See spectrum S-36.

Procedure for Preparation of lactones **12a** and **12b** from **10a/b**:

To a solution of **10a/b** (80 mg, 0.19 mmol) in CH₃OH (3mL) was added acetyl chloride (10□L), and the mixture was stirred at room temperature overnight. Saturated sodium bicarbonate solution (20 mL) was added and the aqueous layer was extracted with ether (3x20mL). The organic layer was dried over magnesium sulfate, filtered and concentrated in vacuo. The residue was purified by flash chromatography (SiO₂, 10:1 pentane:ether) to afford cis-**12a** (13mg, 54%) and trans **12b** (10 mg, 41%)

cis-3-methyl-4-vinyl-dihydrofuran-2(3H)-one (±)-12a

Purification: SiO₂(9:1 pentane:ether) affording **12a** (54%) as a colorless oil

¹H NMR: (400 MHz, CDCl₃)

5.74 (ddd, $J=8.8, 10.0, 16.8$ Hz, 1H); 5.16-5.24 (m, 2H); 4.35 (dd, $J= 6.4, 9.2$ Hz, 1H); 4.14 (dd, $J=3.6, 9.2$ Hz, 1H); 3.16 (m, 1H); 2.75 (q, $J=7.6$ Hz, 1H); 1.16 (d, $J=7.2$ Hz, 3H)

¹³C NMR: (100 MHz, CDCl₃)

179.6; 134.0; 118.9; 71.1; 44.5; 38.5; 11.2

GCMS: $t_R=3.78$ min. (Agilent 6890; injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes); calculated for C₇H₁₀O₂ 126.0681, found $m/z=126.0$ (M⁺)

IR: 2954 cm⁻¹, 2917, 2854, 1774, 1654.

TLC: R_f 0.2 (SiO₂, hexanes:EtOAc 4:1)

See spectrum S-37.

The spectroscopic data reported above for **12a** matched the literature data reported in reference 18 (main text) for cis-3-methyl-4-vinyl-dihydrofuran-2(3H)-one

trans-3-methyl-4-vinyl-dihydrofuran-2(3*H*)-one ($\pm$)-12b

Purification: SiO₂(9:1 pentane:ether) affording **12b** (40%) as a colorless oil

¹H NMR: (400 MHz, CDCl₃)
 5.72 (ddd, *J*=8.0, 10.0, 16.8 Hz, 1H); 5.17-5.26 (m, 2H); 4.37 (dd, *J*=7.6, 9.2 Hz, 1H); 3.91 (dd, *J*=8.8, 10.4 Hz, 1H); 2.77 (m, 1H); 2.36 (q, *J*=7.2 Hz, 1H); 1.25 (d, *J*=6.8 Hz, 3H).

¹³C NMR: (100 MHz, CDCl₃)
 179.1; 135.1; 119.2; 70.3; 49.2; 40.5; 13.2

GCMS: *t_R*=3.50 min. (Agilent 6890; injector 250°C, 10.5 psi, column gradient 100°C for 2 minutes, 25°C/min to 250°C, 250°C for 10 minutes) Calculated for C₇H₁₀O₂ 126.0681, found *m/z*=126.0 (*M*⁺).

IR: 2954 cm⁻¹, 2919, 2858, 1778, 1651.

TLC: *R_f* 0.25 (SiO₂, hexanes:EtOAc 4:1)

See spectrum S-18.

The spectroscopic data reported above for **12b** matched the literature data reported in reference 18 (main text) for trans-3-methyl-4-vinyl-dihydrofuran-2(3*H*)-one

Pages S17-S41:

¹HNMR spectra for compounds 2a, 3a, 2b, 2c, 3b/c, 2d, 3d, 2e, 3e, 2f, 3f, 5a, 5b, 6a, 6b, 7, 8, 9, 10a/b, 11a/b, 12a, 12b: S-17 - S-38

2-D NMR spectrum (NOESY) of compound 8: S-39

¹HNMR spectra for compounds 3am, 3an :S-40, S-41