Macropodumines D and E, Two Novel Alkaloids With Unusual Skeletons From *Daphniphyllum macropodum* Miq.

Zhen-Yu Li[†], Ping Chen[†], Hong-Gui Xu[‡], Yi-Ming Yang[†], Shu-Ying Peng[†],

Zhong-Zhen Zhao[‡] and Yue-Wei Guo^{*,†}

*Shanghai Institute of Material Medica, Chinese Acedemy of Science, Shanghai
201203, People's Republic of China, and School of Chinese Medicine, Hong Kong
Baptist University, Kowloon Tong, Hong Kong 999077*

Corresponding author, Phone: (+86) 21-50803318, Fax: (+86) 21-50807088. Email address:

ywguo@mail.shcnc.ac.cn

Index

Experimental Section	S2
HR-ESIMS spectrum of macropodumine D (1)	S3-S4
¹ H NMR spectrum of macropodumine D (1) (600MHz, CDCl ₃)	S5-S6
¹³ C NMR spectrum of macropodumine D (1) (100MHz, CDCl ₃)	S7-S8
¹ H- ¹ H COSY spectra of macropodumine D (1)	S9-S10
HSQC spectra of macropodumine D (1)	S11
HMBC spectra of macropodumine D (1)	S12-S13
ROESY spectra of macropodumine D (1)	S14-S15
HR-ESIMS spectrum of macropodumine E (2)	S16-S17
¹ H NMR spectrum of macropodumine E (2) (600MHz, CDCl ₃)	S18
¹³ C NMR spectrum of macropodumine E (2) (100MHz, CDCl ₃)	S19-S20
¹ H- ¹ H COSY spectra of macropodumine E (2)	S21-S22
HSQC spectra of macropodumine E (2)	S23-S24
HMBC spectra of macropodumine E (2)	S25-S27
ROESY spectra of macropodumine E (2)	S28-S29

[†] Shanghai Institute of Material Medica

[‡] Hong Kong Baptist University.

Experimental Procedure

General: Commercial silica gel (Qing Dao Hai Yang Chemical Group Co., 200-300 mesh) and amino silica gel (Merck, LiChroprep NH₂40-63 μ m) was used for column chromatography. Precoated silica gel plates (Yan Tai Zi Fu Chemical Group Co., G60 F-254) were used for analytical thin-layer chromatography (TLC). The NMR spectra were recorded at 293 K on a Varian Mercury-600 and 400 spectrometers (600 MHz for ¹H and 100 MHz for ¹³C). Chemical shifts are reported with the residual CHCl₃ (δ_{H} 7.26 ppm) as the internal standard for ¹H NMR spectrometry, and CDCl₃ (δ_{C} 77.0 ppm) for ¹³C NMR spectrometry. ¹H and ¹³C NMR assignments were supported by ¹H-¹H COSY, HSQC, HMBC and ROESY experiments. Optical Rotations were measured on a PERKIN-ELMER 341 polarimeter at the sodium D-line. Infrared spectra were recorded on a Nicolet Magna FT-IR 750 spectrophotometer; peaks are reported in cm⁻¹. The mass spectra and high resolution mass spectra were performed on a Q-TOF Micro LC-MS-MS mass spectrometer.

Plant material: The barks and leaves of *Daphniphyllum macropodum* Miq. were collected in the Emei Mountain, Sichuan Province of P. R. China, in April, 2005, and identified by Professor Honggui Xu of Hongkong Baptist University. A voucher specimen is available for inspection at the Herbarium of the Institute of Matria Medica, Chinese Academy of Science.

Extraction and isolation: The air-dried powdered leaves and barks (1.4 Kg) of *D. macropodum* were extracted with 95% EtOH (three times, each 7 day) at room temperature. Evaporation of the solvent gave a residue, which was suspended in water (1 L) and adjusted to pH 4-5 with 2 N H₂SO₄. The acidic mixture was defatted with EtOAc (3 $\times$ 1 L), and the aqueous layer was basified to pH 9-10 with saturated NaCO₃, and then extracted with CHCl₃ (3 $\times$ 1 L) to yield the crude alkaloids (7.1g). The crude alkaloids were subjected to silica gel chromatography eluted with a CHCl₃/CH₃OH/Et₂NH (50:1:0.1 to 1:1:0.1) gradient to give fractions Fr. 1 to Fr. 10. Further amino silica gel chromatography (CHCl₃/CH₃OH, 33:1) on Fr. 7 and Fr. 2 afforded macropodumines D (**1**) (5.1 mg) and E (**2**) (9.3mg), respectively.

HR-ESIMS spectrum of Macropodumine D (1)

HR-ESIMS spectrum of Macropodumine D (1) (Continued)

Elemental Composition Report

Page

Tolerance = 20.0 PPM / DBE: min = -1.5, max = 50.0

Isotope cluster parameters: Separation = 1.0 Abundance = 1.0%

Monoisotopic Mass, Odd and Even Electron Ions

13 formula(e) evaluated with 1 results within limits (up to 20 closest results for each mass)

DM-Mix-3

20060623-4 738 (13.867) AM (Cen,2, 80.00, Ht,5000.0,362.93,1.00); Sm (Mn, 2x3.00); Sm (Mn, 2x3.00); Cm (719:739)

TOF MS ES+
335

Minimum: -1.5
Maximum: 50.0

Mass	Calc. Mass	mDa	PPM	DBE	Score	Formula
402.2286	402.2280	0.6	1.4	8.5	1	C23 H32 N O5

¹H NMR spectrum of Macropodumine D (**1**) in CDCl₃

¹H NMR spectrum of Macropodumine D (**1**) in CDCl₃ (Enlarged)

^{13}C NMR spectrum of Macropodumine D (**1**) in CDCl_3

DML-14 CDCl₃ 88+DEPT-135 May 29 2006

^{13}C NMR spectrum of Macropodumine D (**1**) in CDCl_3 (Enlarged)

^1H - ^1H COSY spectrum of Macropodumine D (**1**) in CDCl_3

DML-14 CDCl_3 COSY

Solvent: CDCl_3
 Temp: 25.0°C / 298.1 K
 INOVA-600 "s1mm600"
 Relax. delay 1.000 sec
 Acq. time 0.171 sec
 SFO: 599.995 MHz
 20 Width 599.6 Hz
 2 repetitions
 512 increments
 OBSERVED F2: 599.6560144 MHz
 Data file: 20110501.D
 SFO: sine bell 0.085 sec
 F1 DATA PROCESSING
 SFO: sine bell 0.040 sec
 F1: 599.995 MHz
 Total time 21 min, 14 sec

^1H - ^1H COSY spectrum of Macropodumine D (**1**) in CDCl_3 (Enlarged)

HSQC spectrum of Macropodumine D (**1**) in CDCl₃

DML-IT 0003 120000
1_06Jun2006

HMBC spectrum of Macropodumine D (**1**) in CDCl₃

HMBC spectrum of Macropodumine D (**1**) in CDCl₃ (Enlarged)

DML-14 CDCl₃ HMBC

ROESY spectrum of Macropodumine D (**1**) in CDCl₃

DML-14 CDCl₃ ROESY

Solvent: CDCl₃
 Temp.: 25.0 C / 298.1 K
 INOVA-600 "sinn600"
 Relax. delay 1.000 sec
 Aq. time 0.250 sec
 Acq. time 0.171 sec
 Width 10000.0 Hz
 2D Width 10000.0 Hz
 2D Resolution 0.091 Hz
 2 x 512 increments
 OBSERVE H1, 599.6560145 MHz
 DATA PROCESSING
 Gauss. apodization 0.047 sec
 F2 size 2048 x 2048
 FT size 2048 x 2048
 Total time 1 hr, 39 min, 39 sec

ROESY spectrum of Macropodumine D (**1**) in CDCl₃ (Enlarged)

HR-ESIMS spectrum of Macropodumine E (2)

HR-ESIMS spectrum of Macropodumine E (2) (Continued)

Elemental Composition Report

Page

Tolerance = 20.0 PPM / DBE: min = -1.5, max = 50.0

Isotope cluster parameters: Separation = 1.0 Abundance = 1.0%

Monoisotopic Mass, Odd and Even Electron Ions

40 formula(e) evaluated with 1 results within limits (up to 20 closest results for each mass)

DM-MIX

060710-9 101 (1.893) AM (Cen,2, 80.00, Ht,5000.0,566.89,1.00); Sm (Mn, 2x3.00); Cm (86;106)

Minimum: 30.00
Maximum: 100.00

Mass	RA	Calc. Mass	mDa	PPM	DBE	Score	Formula
524.2234	100.00	524.2260	-2.6	-5.0	10.5	1	C27 H35 N O8 Na

¹H NMR spectrum of Macropodumine E (**2**) in CDCl₃

^{13}C NMR spectrum of Macropodumine E (**2**) in CDCl_3

^{13}C NMR spectrum of Macropodumine E (**2**) in CDCl_3 (Enlarged)

^1H - ^1H COSY spectrum of Macropodumine E (**2**) in CDCl_3

^1H - ^1H COSY spectrum of Macropodumine E (**2**) in CDCl_3 (Enlarged)

HSQC spectrum of Macropodumine E (2) in CDCl₃

DM-30 CDCL3 HSQC

HSQC spectrum of Macropodumine E (**2**) in CDCl₃ (Enlargd)

DM-30 CDCl₃ HSQC

HMBC spectrum of Macropodumine E (**2**) in CDCl₃

DM-30 CDCl₃ HMBC

HMBC spectrum of Macropodumine E (**2**) in CDCl₃ (Enlarged 1)

HMBC spectrum of Macropodumine E (**2**) in CDCl₃ (Enlarged 2)

DM-30 CDCL3 HMBC

ROESY spectrum of Macropodumine E (**2**) in CDCl₃

ROESY spectrum of Macropodumine E (2) in CDCl₃ (Enlarged)