REVISED SUPPORTING INFORMATION

Manuscript No. jo0605142

Magnesium bistrifluoromethanesulfonimide as a new and efficient acylation catalyst

*Shivani and Asit K. Chakraborti,**

Department of Medicinal Chemistry, National Institute of Pharmaceutical Education and Research (NIPER), Sector 67, S. A. S. Nagar 160 062, Punjab, India. * Corresponding Author:
akchakraborti@niper.ac.in

Table of Contents

Spectral Data for known compounds:

Entry 1, Table 2: ¹ H NMR	S3
Entry 2, Table 2: ¹ H NMR	S4
Entry 3, Table 2: ¹ H NMR	S5
Entry 4, Table 2: ¹ H NMR	S6
Entry 5, Table 2: ¹ H NMR	S7
Entry 6, Table 2: ¹ H NMR	S8
Entry 7, Table 2: ¹ H NMR	S9
Entry 8, Table 2: ¹ H NMR	S10
Entry 9, Table 2: ¹ H NMR	S11
Entry 10, Table 2: ¹ H NMR	S12
Entry 11, Table 2: ¹ H NMR	S13
Entry 12, Table 2: ¹ H NMR	S14
Entry 13, Table 2: ¹ H NMR	S15
Entry 14, Table 2: ¹ H NMR	S16
Entry 15, Table 2: ¹ H NMR	S17
Entry 16, Table 2: ¹ H NMR	S18
Entry 17, Table 2: ¹ H NMR	S19
Entry 18 Table 2: ¹ H NMR	S20
Entry 18, Table 2: ¹³ C NMR	S21
Entry 18, Table 2: IR	S22
Entry 19, Table 2: ¹ H NMR	S23
Entry 19, Table 2: ¹³ C NMR	S24
Entry 20, Table 3: ¹³ C NMR	S25
Entry 1, Table 3: ¹ H NMR	S26
Entry 2, Table 3: ¹ H NMR	S27
Entry 3, Table 3: ¹ H NMR	S28
Entry 4, Table 3: ¹ H NMR	S29
Entry 5, Table 3: ¹ H NMR	S30
Entry 6, Table 3: ¹ H NMR	S31
Entry 7, Table 3: ¹ H NMR	S32
Entry 8, Table 3: ¹ H NMR	S33
Entry 9, Table 3: ¹ H NMR	S34
Entry 10, Table 3: ¹ H NMR	S35
Entry 11, Table 3: ¹³ C NMR	S36
Entry 12, Table 3: ¹ H NMR	S37
Entry 13, Table 3: ¹ H NMR	S38

Entry 2, Table 4:	¹ H NMR	S39
Entry 3, Table 4:	¹ H NMR	S40
Entry 4, Table 4:	¹ H NMR	S41
Entry 5, Table 4:	¹ H NMR	S42
Entry 6, Table 4:	¹ H NMR	S43
Entry 2, Table 5:	¹ H NMR	S44
Entry 3, Table 5:	¹ H NMR	S45
Entry 4, Table 5:	¹ H NMR	S46
Entry 5, Table 5:	¹ H NMR	S47
Entry 6, Table 5:	¹ H NMR	S48
Entry 7, Table 5:	¹ H NMR	S49
Entry 2, Table 6:	¹ H NMR	S50
Entry 2, Table 6:	¹³ C NMR	S51
Entry 2, Table 6:	IR	S52
Entry 3, Table 6:	¹ H NMR	S53
Entry 4, Table 6:	¹ H NMR	S54
Entry 5, Table 6:	¹ H NMR	S55
Entry 6, Table 6:	¹ H NMR	S56
Entry 7, Table 6:	¹ H NMR	S57

Entry 1, Table 2

Entry 2, Table 2

Entry 3, Table 2

Entry 4, Table 2

Entry 5, Table 2

Entry 6, Table 2

Entry 7, Table 2

Entry 8, Table 2

Entry 9, Table 2

Entry 10, Table 2

Entry 11, Table 2

Entry 12, Table 2

Entry 13, Table 2

Entry 14, Table 2

Entry 15, Table 2

Entry 16, Table 2

Entry 17, Table 2

Entry 18, Table 2

Entry 18, Table 2

Entry 18, Table 2

Entry 19, Table 2

Entry 19, Table 2

Entry 20, Table 2

Entry 1, Table 3

Entry 2, Table 3

Entry 3, Table 3

Entry 4, Table 3

Entry 5, Table 3

Entry 6, Table 3

Entry 7, Table 3

Entry 8, Table 3

Entry 9, Table 3

Entry 10, Table 3

Entry 11, Table 3

Entry 12, Table 3

Entry 13, Table 3

Entry 2, Table 4

Entry 3, Table 4

Entry 4, Table 4

Entry 5, Table 4

Entry 6, Table 4

Entry 2, Table 5

Entry 3, Table 5

Entry 4, Table 5

Entry 5, Table 5

Entry 6, Table 5

Entry 7, Table 5

Entry 2, Table 6

Entry 2, Table 6

Entry 2, Table 6

Entry 3, Table 6

Entry 4, Table 6

Entry 5, Table 6

Entry 6, Table 6

Entry 7, Table 6

Experimental:

General details

The ^1H and ^{13}C NMR spectra were recorded on Bruker Avance DPX 300 (300 MHz) spectrometer in CDCl_3 using TMS as internal standard. The IR spectra were recorded on Nicolet Impact 400 spectrometer as KBr pellets for solid and neat for liquid samples. The reactions were monitored by TLC (silica gel-G) and Shimadzu QP 5000 GCMS. The inert atmosphere for carrying out the reaction was maintained using dry nitrogen or argon using flame dried glasswares. Evaporation of solvents was performed at reduced pressure, using a Büchi rotary evaporator.

Typical procedure for 2-Acetoxy-naphthalene (entry 1, table 1)

A mixture of 2-naphthol (360 mg, 2.5mmol), Ac_2O (0.24mL, 3mmol) and $\text{Mg}(\text{NTf}_2)_2$ (1mol%) was stirred magnetically under neat condition at room temperature for 15 min. The reaction mixture was diluted with H_2O (10mL) and extracted with Et_2O (3 X15mL). The combined ethereal extracts were washed successively with 2% aqueous NaOH (15mL) and saturated brine (15mL), dried (Na_2SO_4) and concentrated under reduced pressure to afford the product, which was in full agreement with the m.pt. and spectral data (IR, ^1H and ^{13}C NMR) of an authentic sample of 2-acetoxy-naphthalene. All the remaining reactions were carried out following this general procedure except that the reaction mixture was heated at 80°C , wherever applicable. In each occasion the product obtained after the routine workup was of sufficient purity (spectral data) and did not require further purification.

Spectral Data of All Compounds

2-naphthyl acetate (entry 1, table 2)¹: ^1H NMR (300 MHz, CDCl_3): δ 2.36 (s, 3H), 7.24 (d, $J=8.85$ Hz, 1H), 7.47 (m, 2H); 7.56 (s, 1H), 7.78(m, 3H).

1-naphthyl acetate (entry 2, table 2)¹: ^1H NMR (300 MHz, CDCl_3): δ 2.45 (s, 3H), 7.23-7.25 (d, $J=6.59$ Hz, 1H), 7.43-7.55 (m, 3H), 7.72-7.75 (d, $J=8.29$ Hz, 1H), 7.84-7.88 (m, 2H).

4-Methoxyphenyl acetate (entry 3, table 2)²: ^1H NMR (300 MHz, CDCl_3): δ 2.27 (s, 3H), 3.79 (s, 3H), 6.88 (d, $J=8.8$ Hz, 2H), 7.00(d, $J=9.0$ Hz, 2H).

4-bromophenyl acetate (entry 4, table 2)³: ^1H NMR (300 MHz, CDCl_3): δ 2.29 (s, 3H), 6.97 (d, $J=8.7$ Hz, 2H), 7.48 (d, $J=8.1$ Hz, 2H).

4-chlorophenyl acetate (entry 5, table 2)⁴: ^1H NMR (300 MHz, CDCl_3): δ 2.25 (s, 3H), 6.62 (d, 2H, $J=8.9$ Hz), 7.31 (d, 2H, $J=8.9$ Hz)

4-acetoxyacetophenone (entry 6, table 2)⁵: ¹H NMR (300 MHz, CDCl₃): δ 2.32 (s, 3H), 2.52 (s, 3H), 7.10 (d, *J*= 8.0 Hz, 1H), 7.28 (t, *J*= 7.6 Hz, 1H), 7.50 (t, *J*= 7.7 Hz, 1H), 7.78 (d, *J*=7.8 Hz, 1H).

Ethyl-4-acetoxybenzoate (entry 7, table 2)⁶: ¹H NMR (300 MHz, CDCl₃): δ 1.38 (t, *J*= 7.1 Hz, 3H), 2.30 (s, 3H), 4.35 (q, *J*= 7.1 Hz, 2H), 7.15 (d, *J*= 8.6 Hz, 2H), 8.06 (d, *J*= 8.6 Hz, 2H).

4-nitrophenyl acetate (entry 8, table 2)¹: ¹H NMR (300 MHz, CDCl₃): δ 2.36 (s, 3H), 7.29 (d, *J*= 8.8 Hz, 2H), 8.28 (d, *J*=9.2 Hz, 2H).

4-cyanophenyl acetate (entry 9, table 2)⁷: ¹H NMR (300 MHz, CDCl₃): δ 2.33 (s, 3H), 7.25 (d, *J*= 8.5 Hz, 2H), 7.70 (d, *J*= 8.5 Hz, 2H).

2-nitrophenyl acetate (entry 10, table 2)⁸: ¹H NMR (300 MHz, CDCl₃): δ 2.35 (s, 3H), 7.23 (d, *J*= 8.10 Hz, 1H), 7.37 (t, *J*=7.75 Hz, 1H), 7.64 (t, *J*= 7.75 Hz, 1H), 8.06 (d, *J*= 8.10 Hz, 1H)

benzene-1, 2-diyl diacetate (entry 11, table 2)⁸: ¹H NMR (300 MHz, CDCl₃): δ 2.28 (s, 3H), 7.16-7.26 (m, 4H).

benzene-1, 3-diyl diacetate (entry 12, table 2)¹: ¹H NMR (300 MHz, CDCl₃): δ 2.27 (s, 3H), 6.92- 6.99 (m, 2H), 7.37(t, *J*=8.1Hz).

benzene-1, 4-diyl diacetate (entry 13, table 2)¹: ¹H NMR (300 MHz, CDCl₃): δ 2.28 (s, 3H), 7.09 (s, 3H).

benzene-1, 2,3-triyl triacetate (entry 14, table 2)⁵: ¹H NMR (300 MHz, CDCl₃): δ 2.28 (s, 3H), 7.10 (d, *J*= 8.03 Hz, 2H), 7.24 (d, *J*= 7.05 Hz, 2H).

S-phenyl thioacetate (entry 15, table 2)¹: ¹H NMR (300 MHz, CDCl₃): δ 2.4 (s, 3H), 7.40(m, 4H).

S-(4-methyl) phenyl thioacetate (entry 16, table 2)⁹: ¹H NMR (300 MHz, CDCl₃): δ 2.35 (s, 3H), 2.37(s, 3H), 7.20 (d, *J*= 8.0 Hz, 2H), 7.28 (d, *J*= 8.1 Hz, 2H).

S-(4-methoxy) phenyl thioacetate (entry 17, table 2)¹⁰: ¹H NMR (300 MHz, CDCl₃): δ 2.39 (s, 3H), 3.82 (s, 3H), 6.93 (d, *J*= 8.6 Hz, 2H), 7.31 (d, *J*= 8.60 Hz, 2H).

S-(4-Nitro) phenyl thioacetate (entry 18, table 2): M.P. IR (neat) 1709 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 2.49 (s, 3H), 7.59 (d, *J*= 9.2Hz, 2H), 8.24(d, *J*= 9.2 Hz, 2H). ¹³C NMR (300 MHz, CDCl₃): δ 30.48, 55.54, 123.87, 124.35, 126.20, 134.55, 191.24.

Furyl mercaptan (entry 19, table 2): IR (neat) 1693 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 2.34 (s, 3H), 4.14 9s, 2H), 6.20 (m,1H), 6.21 (m,1H), 7.31 (m, 1H). ¹³CNMR((300 MHz, CDCl₃): δ 23.15, 30.36, 106.27, 110.62, 142.22, 149.68, 194.54.

benzylthioacetae (entry 18, table 2)⁹: ¹H NMR (300 MHz, CDCl₃): δ 2.30 (s, 3H), 4.09 (s, 2H), 7.20-7.29(m, 5H).

benzylacetate (entry 1, table 3)¹: ¹H NMR (300 MHz, CDCl₃): δ 2.04 (s, 3H), 5.06 (s, 2H), 7.20-7.31 (m, 5H).

2-phenylethyl acetate (entry 2, table 3)¹¹: ¹H NMR (300 MHz, CDCl₃): δ 1.98 (s, 3H), 2.90 (t, *J*= 7.09 Hz, 2H), 4.25 (t, *J*= 7.09 Hz, 2H), 7.17-7.29 (m, 5H).

1-phenylethyl acetate (entry 3, table 3)¹²: ¹H NMR (300 MHz, CDCl₃): δ 1.52 (d, *J*= 6.6 Hz, 3H), 2.06 (s, 3H), 5.88 (m, 1H), 7.24-7.53 (m, 5H).

1-phenyl-2-ethyl acetate (entry 4, table 3)¹⁰: ¹H NMR (300 MHz, CDCl₃): δ 0.87 (t, *J*= 7.38 Hz, 3H), 1.73-1.99 (m, 2H), 2.00 (s, 3H), 5.66 (t, *J*= 6.86 Hz, 1H), 7.28 (m, 5H).

2-acetoxy-2-phenylacetophenone (entry 5, table 3)¹¹: ¹H NMR (300 MHz, CDCl₃): δ 2.15 (s, 3H), 6.86 (s, 1H), 7.23-7.44 (m, 8H), 7.90 (d, *J*= 7.2 Hz, 2H).

Cinnamyl acetate (entry 6, table 3)⁸: ¹H NMR (300 MHz, CDCl₃): δ 2.04 (s, 3H), 4.68 (d, *J*= 6.09 Hz, 2H), 6.19-6.29 (m, 1H), 6.58 (d, *J*= 16 Hz, 1H), 7.21-7.36 (m, 5H).

2,2-dimethyl-phenylacetate (entry 7, table 3)¹³: ¹H NMR (300 MHz, CDCl₃): δ 1.44 (6H), 1.97 (s, 3H), 3.05 (s, 2H), 7.17-7.31 (m, 5H).

Geranyl acetate (entry 8, table 3)¹⁴: ¹H NMR (300 MHz, CDCl₃): δ 1.60 (s, 3H), 1.68 (s, 3H), 1.70 (s, 3H), 2.05 (s, 3H), 2.09 (m, 4H), 4.59 (d, *J*= 7.1 Hz, 2H), 5.08 (m, 1H), 5.34 (t, *J*= 7.1 Hz, 1H).

Menthylacetate (entry 9, table 3)³: ¹H NMR (300 MHz, CDCl₃): δ 0.76 (d, *J*= 6.58 Hz, 3H), 0.90 (d, *J*= 6.32 Hz, 6H), 1.01 (m, 4H), 1.36 (m, 1H), 1.50 (m, 1H), 1.67 (m, 2H), 1.88 (m, 1H), 2.04 (s, 3H), 4.66 (m, 1H).

Isomenthylacetate (entry 10, table 3)⁶: ¹H NMR (300 MHz, CDCl₃): δ 0.85 (d, *J*= 7 Hz, 3H), 0.93 (m, 6H), 1.24-1.87 (m, 9H), 2.03 (s, 3H), 4.99-5.05 (m, 1H).

[(1*S*)-endo]-(-)-Bornyl acetate (entry 11, table 3)¹¹: ¹H NMR (300 MHz, CDCl₃): δ ¹³C NMR (75 MHz, CDCl₃): 13.42, 18.72, 19.66, 21.23, 27.02, 27.99, 36.71, 44.85, 47.72, 48.64, 79.81, 171.38.

1-ethynylcyclohexyl-1-yl acetate (entry 12, table 3)¹²: ¹H NMR (300 MHz, CDCl₃): δ 1.28-1.65 (m, 6H), 1.80-1.88 (m, 2H), 2.04 (s, 3H), 2.10-2.14 (m, 2H), 2.61 (m, 1H).

1-adamantanyl acetate (entry 13, table 3)¹¹: ¹H NMR (300 MHz, CDCl₃): δ 1.65 (m, 6H), 1.96 (m, 3H), 2.09 (m, 9H).

2-naphthyl propionate (entry 2, table 4)¹⁵: ¹H NMR (300 MHz, CDCl₃): δ 1.31 (t, *J*= 7.49 Hz, 3H), 2.65 (q, *J*= 7.53 Hz, 2H), 7.24 (d, *J*= 8.89 Hz, 1H), 7.47 (m, 2H), 7.55 (s, 1H), 7.78-7.86 (m, 3H).

2'-naphthyl 2, 2-dimethylacetate (entry 3, table 4)¹⁶: ¹H NMR (300 MHz, CDCl₃): δ 1.36 (d, *J* = 7.00Hz, 6H), 2.0-2.90(m,1H), 7.21(d, *J* = 8.90Hz, 1H), 7.47(m, 2H), 7.54(s, 1H), 7.77-7.84 (m, 3H).

2-naphthyl pivalate (entry 4, table 4)⁶: ¹H NMR (300 MHz, CDCl₃): δ 1.39(s, 9H), 7.20(d, *J* = 8.71 Hz, 1H), 7.46(m, 2H), 7.51(s, 1H), 7.75-7.83(m,3H).

2-naphthyl benzoate (entry 5, table 4)⁵: ¹H NMR (300 MHz, CDCl₃): δ 7.36 (d, *J* = 8.86Hz, 1H), 7.48 (m, 2H), 7.54(s, 1H), 7.63(m, 3H), 7.83(m,3H), 8.13(d, *J* = 8.86Hz, 1H), 8.27(d, *J* = 8.86, 1H).

2-naphthyl chloroacetate (entry 6, table 4)¹⁵: ¹H NMR (300 MHz, CDCl₃): δ 4.3 (s, 2H), 7.05 (m, 1H), 7.47-7.49(m, 2H), 7.60(s, 1H), 7.78-7.86 (m, 3H).

4-nitro benzoate (entry 2, table 5)¹⁷: ¹H NMR (300 MHz, CDCl₃): δ 7.43 (d, *J* = 9Hz, 2H), 7.56(t, *J* = 7.65Hz, 2H), 7.70(t, *J* = 7.35 Hz, 1H), 8.21(d, *J* = 7Hz, 2H), 8.34(d, *J* = 9Hz, 2H).

Adamantyl benzoate (entry 3, table 4)⁶: ¹H NMR (300 MHz, CDCl₃): δ 1.66 (s, 6H), 2.26 (s, 9H), 7.38-7.4 (m, 2H), 7.50-7.54 (m, 1H), 7.98 (d, *J* = 6.8 Hz, 2H).

Benzene-1,4-diyl dibenzoate (entry 4, table 5)⁸: ¹H NMR (300 MHz, CDCl₃): δ 7.17 (d, *J* = 9.1Hz, 4H), 7.45-7.66 (m, 6H), 8.20 (d, *J* = 7.7 Hz, 4H).

2-phenethyl benzoate (entry 5, table 5)¹⁸: ¹H NMR (300 MHz, CDCl₃): δ 3.02 (t, *J* = 6.7 Hz, 2H), 4.49 (t, *J* = 6.7 Hz, 2H), 7.14-7.49 9m, 8H), 8.00 (d, *J* = 6.7 Hz, 2H).

Menthyl benzoate (entry 6, table 5)¹⁸: ¹H NMR (300 MHz, CDCl₃): δ 0.80 (d, *J* = 7.0 Hz, 3H), 0.93 (dd, *J* = 7.0 Hz, 6H), 1.10 (m, 2H), 1.55 (m, 2H), 1.74 (m, 3H), 1.98 (m, 1H), 2.13 (m, 1H), 4.04 (m, 1H), 7.44 (t, *J* = 7.45 Hz, 2H), 7.55 (t, *J* = 7.35 Hz, 1H), 8.05 (d, *J* = 7.15 Hz, 2H)

2, 4, 6-trimethylphenyl benzoate (entry 7, table 5)^{12b}: ¹H NMR (300 MHz, CDCl₃): δ 2.15 (s, 6H), 2.28(s, 3H), 6.90(s, 2H), 7.51 (t, *J* = 7.7 Hz, 3H), 8.24 (d, *J* = 7.7 Hz).

4-nitro pivalate (entry 2, table 6): IR (neat): 1759 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 1.37 (s, 9H), 7.25(d, *J* = 8.6Hz, 2H), 8.27 (d, *J* = 8.6 Hz, 2H). ¹³C NMR(75 MHz, CDCl₃): δ 26.99, 39.29, 122.39, 125.13, 155.96, 176.13.

1-adamantyl pivalate (entry 3, table 6)⁸: ¹H NMR (300 MHz, CDCl₃): δ 1.14 (s, 9H), 1.64 (s, 9H), 2.12 (s, 6H).

benzyl pivalate (entry 4, table 6)¹³: ¹H NMR (300 MHz, CDCl₃): δ 1.22 (s, 9H), 5.10 (s, 2H), 7.33 (s, 5H).

Cinnamyl pivalate (entry 5, table 6)¹³: ¹H NMR (300 MHz, CDCl₃): δ 1.22 (s, 9H), 4.30 (d, *J* = 5.4 Hz, 1H), 4.71 (d, *J* = 6.2 Hz, 1H), 6.24- 6.37 (m, 1H), 6.60-6.66 (m, 1H), 7.22-7.39 (m, 5H)

2, 2-dimethyl—phenylpivalate (entry 7, table 3)¹³: ¹H NMR (300 MHz, CDCl₃): δ 1.12 (s, 1H), 1.43 (s, 6H), 3.20 (s, 2H), 7.17-7.28 (m, 5H).

bornyl pivalate (entry 8, table 6)¹⁸: ¹H NMR (300 MHz, CDCl₃): δ 0.83 (s, 3H), 0.88(s, 3H), 0.91 (s, 3H), 1.21 (s, 9H), 1.28 (m, 2H), 1.71 (m, 3H), 1.97 (m, 1H), 2.34 (m, 1H), 4.85 (d, *J* = 9.58 Hz, 1H).

Reference:

1. Pouchart, C. J.; Jacqlynn, B.; I ed. *The Aldrich Libraray of ¹³C and ¹H FT NMR Spectra*; Aldrich Chemical: Milwaukee, 1993; Vol. II.
2. Li, A.-X.; Li, T.-S.; Ding, T.-H. *Chem. Commun* **1997**, 1389-1390.
3. (a) Saravanam, P.; Singh, V. K. *Tetrahedron Lett.* **1999**, 40, 2611-2614. (b) Chandra, K. L.; Saravanam, P.; Singh, R. K.; Singh, V. K. *Tetrahedron* **2002**, 58, 1369-1374.
4. jkio
5. Cirovic, M. M. *Properties of Organic Compounds*; CRC, 1996; POC-personal edn, version 5.1.
6. Chakraborti, A. K.; Sharma, L.; Gulhane, R.; Shivani. *Tetrahedron* **2003**, 59, 7661-7668.
7. Alves, K. B.; Bastos, M. P.; Amaral, L. *J. Org. Chem.* **1978**, 43, 4032-4038.
8. Mohammadpoor-Baltrok, I.; Aliyan, H.; Khosropour, A. R. *Tetrahedron* **2001**, 57, 5851-5854.
9. (a) Iqbal, J.; Srivastava, R. *J. Org. Chem.* **1992**, 57, 2001-2007. (b) Ahmed, S.; Iqbal, J. *Tetrahedron Lett.* **1986**, 27, 3791-3794.
10. Chakraborti, A. K.; Gulhane, R.; Shivani. *Synlett* **2003**, 1805-1808.
11. Orita, A.; Tanahashi, C.; Kakuda, A.; Otera, J. *Angew. Chem. Int. Ed. Engl.* **2000**, 39, 2877-2879.
12. Ishihara, K.; Kubota, M.; Kurihara, H.; Yamamoto, H. *J. Am. Chem. Soc.* **1995**, 117, 4413-4414. (b) Ishihara, K.; Kubota, M.; Kurihara, H.; Yamamoto, H. *J. Org. Chem.* **1996**, 61, 4560-4567.
13. Orita, A.; Tanahashi, C.; Kakuda, A.; Otera, J. *J. Org. Chem.* **2001**, 66, 8926-8934.
14. Pouchart, C. J.; Jacqlynn, B.; I ed. *The Aldrich Libraray of ¹³C and ¹H FT NMR Spectra*; Aldrich Chemical: Milwaukee, 1993; Vol. I.
15. Prasad, M.; Jigajinni, V. B. *Indian J. Chem.* **1980**, 19B, 822-823.
16. Lowrance, W. W., Jr. *Tetrahedron Lett.* **1971**, 3453-3454.
17. Furniss, B. S.; Hannaford, A. J.; Smith, P. W. G.; Tatchell, A. R. *Vogel's Textbook of Practical Organic Chemistry*; 5th ed. Longman: Singapore, 1989.