Supporting Information

to the article “**Metal-Metal bonding in tetracyanometallates ($M = \text{Pt}^{\text{II}}$, Pd^{II} , Ni^{II}) of monovalent thallium. Crystallographic and spectroscopic characterisation of the compounds $\text{Tl}_2\text{Ni}(\text{CN})_4$ and $\text{Tl}_2\text{Pd}(\text{CN})_4$.**”

by Mikhail Maliarik, Jeffrey K. Nagle, Andrey Ilyuhin, Elena Murashova, János Mink, Mikhail Skripkin, Julius Glaser, Margit Kovacs, and Attila Horváth

Contents:

Figure S1. Experimental polycrystalline (blue) and calculated from the single-crystal data X-ray (red) diffractograms of the $\text{Tl}_2\text{M}(\text{CN})_4$ ($M = \text{Ni}$ (a) and Pt (b)) compounds.

Figure S2. Luminescence spectra of polycrystalline samples of:

- (a) $\text{Tl}_2\text{Ni}(\text{CN})_4$ (red) and $\text{K}_2\text{Ni}(\text{CN})_4 \times \text{H}_2\text{O}$ (black), excitation at 300 nm,
- (b) $\text{Tl}_2\text{Pd}(\text{CN})_4$ (red) and $\text{K}_2\text{Pd}(\text{CN})_4 \times 3\text{H}_2\text{O}$ (black); excitation at 400 nm

Figure S3. Excitation spectra of polycrystalline samples of $\text{Tl}_2\text{Ni}(\text{CN})_4$ (red) and $\text{K}_2\text{Ni}(\text{CN})_4 \times \text{H}_2\text{O}$ (black) with emission at 490 (a) and 620 (b) nm.

Figure S4. Excitation spectra of polycrystalline samples of $\text{Tl}_2\text{Pd}(\text{CN})_4$ (red) and $\text{K}_2\text{Pd}(\text{CN})_4 \times 3\text{H}_2\text{O}$ (black) with emission at 531 (a) and 606 (b) nm.

Figure S1.

a.

b.

Figure S2

a.

b.

Figure S3.

a.

b.

Figure S4.

a.

b.