S1 Supporting Information Contents Page

Title: Microcionamides A and B, Bioactive Peptides from the Philippine Sponge

Clathria (Thalysias) abietina

Authors: Rohan A. Davis, Gina C. Mangalindan, Zenaida P. Bojo, Rowena R. Antemano, Nell O. Rodriguez, Gisela P. Concepcion, Shiela C. Samson, Dennis de Guzman, Lourdes J. Cruz, Deniz Tasdemir, Mary Kay Harper, Xidong Feng, Guy T. Carter, and Chris M. Ireland

Contents:

S2 ¹H NMR Spectrum for Microcionamide A (**1**) in CD₃OD

S3 ¹³C NMR Spectrum for Microcionamide A (**1**) in CD₃OD

S4 ¹H NMR Spectrum for Microcionamide B (**2**) in CD₃OD

S5 ¹³C NMR Spectrum for Microcionamide B (**2**) in CD₃OD

S6 General Experimental Procedures

S7 Table of MS/MS studies and (+)-HRESIMS measurements for Microcionamide A.

S2 ^1H NMR Spectrum for Microcionamide A (**1**) in CD_3OD

S3 ^{13}C NMR Spectrum for Microcionamide A (**1**) in CD_3OD

S4 ^1H NMR Spectrum for Microcionamide B (**2**) in CD_3OD

S5 ^{13}C NMR Spectrum for Microcionamide B (**2**) in CD_3OD

S6 General Experimental Procedures

NMR spectra were recorded at 26 °C on a Varian Unity 500 spectrometer at 500.620 MHz for ^1H and 125.893 MHz for ^{13}C . The ^1H and ^{13}C chemical shifts are reported in parts per million relative to the reference solvent signals at δ 3.30 and 49.00 ppm for CD_3OD . FTIR and UV spectra were recorded on a Jasco FT/IR-420 spectrophotometer and a Hewlett Packard 8452A diode array spectrophotometer, respectively. Specific rotations were measured on a JASCO DIP-370 digital polarimeter. (+)-LRESIMS measurements were recorded on a ThermoFinnigan LCQ-classic mass spectrometer. MS/MS studies and (+)-HRESIMS measurements were recorded on a Bruker APEXII FTICR mass spectrometer. Flash chromatography was performed using a glass column (20 $\times$ 250 mm) packed with EM Science LiChroprep C_{18} 40-63 μm 100 Å. A Rainin Dynamax C_{18} 5 μm 100 Å column (10 $\times$ 250 mm) connected to a Shimazu LC-10AV/VP HPLC system was used for semipreparative HPLC. A Phenomenex Luna $\text{C}_{18}(2)$ 5 μm 100 Å column (4.6 $\times$ 250 mm) was used for analytical HPLC work. Automatic protein sequence analysis was carried out using an Applied Biosystems Procise sequencer. All solvents used for HPLC, $[\alpha]_{\text{D}}$, UV and MS were Fisher HPLC grade, and the water used was Barnstead E-pure 0.2 μm filtered. Synthetic reagents used were purchased from Sigma-Aldrich.

S7 Table of MS/MS studies and (+)-HRESIMS measurements for Microcionamide A

Experimental Mass	Theoretical Mass	Δ (mDa)	Δ (ppm)	Intensity	Relative Intensity (%)	Elemental Formula	Assignment
875.48843	875.48816	0.27	0.31	0.2625	26.24	C43H71N8O7S2	[M+H] ¹⁺
858.46174	858.46162	0.12	0.14	0.9249	92.46	C43H68N7O7S2	[M+H] ¹⁺ -NH ₃
840.45131	840.45105	0.26	0.31	0.0838	8.37	C43H66N7O6S2	[M+H] ¹⁺ -NH ₃ - H ₂ O
830.46661	830.4667	-0.09	-0.11	0.5904	59.03	C42H68N7O6S2	[M+H] ¹⁺ -NH ₃ -CO
813.43971	813.44015	-0.44	-0.54	0.1417	14.16	C42H65N6O6S2	[M+H] ¹⁺ -2NH ₃ -CO
757.46848				0.1291			
756.41473	756.41467	0.06	0.08	0.8662	86.59	C35H62N7O7S2	[M+H] ¹⁺ -NH -(CH) ₂ -Phe
745.37733	745.37755	-0.22	-0.30	0.1411	14.11	C37H57N6O6S2	[M+H] ¹⁺ -NH ₃ - I
728.42022				0.0367			
717.38256	717.38264	-0.08	-0.11	0.0850	8.49	C36H57N6O5S2	[M+H] ¹⁺ -NH ₃ -CO -I
711.39219				0.0280			
700.35601	700.35609	-0.08	-0.11	0.3401	34.00	C36H54N5O5S2	[M+H] ¹⁺ -2NH ₃ -CO - I
687.39312				0.0782			
676.35598				0.1523			
657.34602	657.34625	-0.23	-0.35	0.2262	22.62	C30H53N6O6S2	(NH)CIIIC
655.42114	655.42113	0.01	0.02	0.3208	32.07	C32H59N6O6S	[M+H] ¹⁺ -NH -(CH) ₂ -Phe -C
653.40531				0.0812			
644.38371				0.2053			
643.33046	643.3306	-0.14	-0.22	1.0003	100.00	C59H51N6O6S2	[M+H] ¹⁺ -NH -(CH) ₂ -Phe -I
637.41014				0.0377			
632.29348	632.29349	-0.01	-0.02	0.0820	8.20	C31H46N5O5S2	[M+H] ¹⁺ -NH ₃ - II
621.43212				0.0236			
610.39722				0.0189			
604.29794				0.0254			
598.30837				0.0252			
588.32735				0.0180			
587.2718	587.27202	-0.22	-0.37	0.1871	18.70	C30H43N4O4S2	[M+H] ¹⁺ -2NH ₃ -CO - II
563.27196				0.2424			
556.35269	556.35272	-0.03	-0.05	0.0273	2.73	C27H50N5O5S	IIIC
554.33798				0.0222			
552.41161				0.0447			

544.26198	544.26219	-0.21	-0.39	0.1402	14.02	C24H42N5O5S2	(NH)CIIC
542.33706	542.33707	-0.01	-0.02	0.1016	10.16	C26H48N5O5S	[M+H]1+ -NH -(CH)2-Phe -IC
540.32195				0.0270			
536.23618				0.0303			
535.27718				0.0407			
531.29988				0.1362			
530.24644	530.24654	-0.1	-0.19	0.4287	42.86	C23H40N5O5S2	[M+H]1+ -NH -(CH)2-Phe -II
522.36461				0.0853			
519.2097	519.20942	0.28	0.54	0.0432	4.32	C25H35N4O4S2	[M+H]1+ -NH3 - III
485.22576				0.0276			
474.18804	474.18796	0.08	0.17	0.0762	7.61	C24H32N3O3S2	[M+H]1+ -2NH3 -CO - III
453.34363	453.34353	0.1	0.22	0.0885	8.85	C24H45N4O4	IIII
450.18793				0.1496			
443.26812	443.26866	-0.54	-1.22	0.0276	2.76	C21H39N4O4S	IIIC
442.30608				0.0336			
439.32788				0.0550			
431.17796	431.17812	-0.16	-0.37	0.0677	6.77	C18H31N4O4S2	(NH)CIIC
425.34872				0.0363			
418.21598				0.0739			
417.16253	417.16247	0.06	0.14	0.2454	24.53	C17H29N4O4S2	[M+H]1+ -NH -(CH)2-Phe -III
409.28093				0.0881			
389.1673				0.0207			
386.15686				0.0311			
384.22812				0.0344			
378.22248							
372.14112				0.0637			
340.25948	340.25947	0.01	0.03	0.0795	7.95	C18H34N3O3	III
337.10398				0.0573			
330.18461	330.18459	0.02	0.06	0.0217	2.17	C15H28N3O3S	IIC
329.22236				0.0430			
318.09422	318.09406	0.16	0.50	0.0452	4.52	C12H20N3O3S2	(NH)CIC
317.13192				0.0295			
316.16847	316.16894	-0.47	-1.49	0.0181	1.81	C14H26N3O3S	[M+H]1+ -NH -(CH)2-Phe -IIIC
305.1318				0.0506			
304.07837	304.07841	-0.04	-0.13	0.0778	7.77	C11H18N3O3S2	[M+H]1+ -NH -(CH)2-Phe -IIII
303.11595				0.0323			
301.22748				0.0260			
296.19693				0.0569			
295.23803				0.0607			
286.17127				0.0436			

282.18144				0.0341				
273.07242				0.0268				
271.144				0.0468				
259.05695				0.0289				
252.15176				0.0409				
245.07763				0.0253				
231.06208				0.0421				
227.1754	227.1754	0	0.00	0.0656	6.56	C12H23N2O2	II	
199.18055				0.0558				
154.15913				0.0250				