Supporting Information

Microwave Assisted Preparation of Fused Bicyclic Heteroaryl Boronates: Application in One-Pot Suzuki Couplings

Erin F. DiMauro* and Jason R. Vitullo

*Department of Chemistry Research and Discovery, Amgen Inc., One Kendall Square,
Bldg. 1000, Cambridge, MA 02139*

Table of Contents

General Considerations	S2
Experimental Details and Analytical Data	S3
¹ H and ¹³ C NMR Spectra of boronates 5a , 6a , 7a	S14
HPLC of 1a-e , 2a-e , 3a-h , 7a , 12	S20
References	S40

General Considerations

Microwave reactions were performed in a Laboratory microwave using 10-20 mL sealed vials in an atmosphere of dry nitrogen. Reported reaction temperatures were preset and were measured and maintained within 1 °C by the instrument. Reaction times are reported as hold time at final temp (ramp time was 2 – 5 °/sec). Reactions were monitored using LCMS with UV detection at 254 nm and a low resonance electrospray mode (ESI). All compounds were purified by preparative medium pressure liquid chromatography (MPLC) with normal-phase silica gel (35 – 60 micron) columns and UV detection at 254 nm. Purity was measured using high performance liquid chromatography (HPLC) with UV detection at 254 nm (15 min; 1.5 mL/min flow rate; 0 to 100% 0.1% TFA in CH₃CN / 100 – 0% 0.1% TFA in H₂O). ¹H NMR spectra were recorded on a 400 MHz spectrometer at ambient temperature. Chemical shifts are reported in ppm from the solvent resonance (DMSO-d₆ 2.49 ppm). Data are reported as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants, and number of protons. Proton decoupled ¹³C NMR spectra were recorded on a AV-400 (100 MHz) spectrometer at ambient temperature. Chemical shifts are reported in ppm from tetramethylsilane (0 ppm) or from the solvent resonance. Mass spectra were obtained on a high resonance Electrospray Time-of-Flight mass spectrometer.

Experimental Details and Analytical Data

6-Bromoquinazolin-2-amine (4a): A stirred mixture of guanidine carbonate (281 g, 1.56 mol), NEt(*i*-Pr)₂ (540 mL, 3.12 mol) and NMP (2 L) was heated to 150-160 °C. A solution of 5-bromo-2-fluoro-benzaldehyde (250 g, 1.20 mol) in NMP (100 mL) was added dropwise via addition funnel over 1 h while maintaining reflux. Upon complete addition, the mixture was maintained at 150-160 °C for 1-2 h until complete consumption of the aldehyde as determined by LC analysis. Upon completion, the reaction was allowed to cool to below 100 °C, then quenched by the addition of ice (2 kg) and water (4 L). The resulting bronze solid was stirred for 30 min, then isolated by vacuum filtration, washing with water (1 L) then denatured EtOH (1 L). The solids were then transferred to a 5 L flask and stirred in denatured EtOH (2 L) for 2 h before refiltering. Subsequent washes in EtOH (0.5 L), a 1:1 mixture of toluene/EtOH (0.5 L), then toluene (0.5 L) and drying afforded the title compound (168 g, 0.750 mol, 48%) as a pale yellow solid. Analytical data is identical to that previously reported.¹

2-Fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzaldehyde (11): To a 100 mL round bottomed flask, was added 5-bromo-2-fluorobenzaldehyde **10** (8.26 g, 40.7 mmol), followed by bis(pinacolato)diboron (11.4 g, 44.8 mmol), KOAc (11.8 g, 120 mmol), and Pd(dppf)Cl₂ (0.89 g, 1.2 mmol). DMSO (60 mL) was added and the reaction was purged with N₂. The mixture was heated at 80 °C, under an atmosphere of N₂, overnight. The mixture was cooled to rt. Water (100 mL) was added and the mixture was extracted with EtOAc (3 x 200 mL). The organics were dried over MgSO₄, filtered and concentrated in vacuo. The crude product was purified by MPLC (SiO₂, eluent: 10%

EtOAc/hexanes) to afford the title compound (8.40 g, 98.5 mmol, 82%) as a white solid.

Analytical data is identical to that previously reported.²

2-Hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzaldehyde (12): To a 10-20 mL microwave reaction vessel was added 5-bromosalicylaldehyde **8** (1.00 g, 4.96 mmol), followed by bis(pinacolato)diboron (1.39 g, 5.45 mmol), KOAc (0.974 g, 9.92 mmol), Pd(dppf)Cl₂ (0.406 g, 4.96 mmol), and 1,4-dioxane (15.0 mL). The vessel was sealed, and the system was purged with N₂. The mixture was heated in the microwave for 45 min at 140 °C then allowed to cool to rt, transferred to a round bottom flask and concentrated to dryness. The crude product was purified by MPLC (SiO₂, eluent: 100% CH₂Cl₂) to afford the title compound (0.583 g, 2.35 mmol, 47%) as an off-white solid: HPLC: 94.9%; LCMS (ESI, pos. ion), *m/z* 249.1 [M+1]; HRMS: Calc'd for [C₁₃H₁₈BO₄]⁺: 249.12927, Found: 249.12949; ¹H NMR (400 MHz, DMSO-*d*₆): δ 11.18 (s, 1H), 10.37 (s, 1H), 8.10 (s, 1H), 7.86 (d, *J* = 8.0 Hz, 1H), 7.09 (d, *J* = 8.0 Hz, 1H), 1.37 (s, 12H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 191.7, 163.1, 142.0, 136.5, 121.9, 117.5, 116.9, 83.6, 24.6.

Ethyl 5-(2-aminoquinazolin-6-yl)benzo[b]thiophene-2-carboxylate (1a): General procedure A: 0.123 g (56.7 %), yellow solid. HPLC: 98.1 %; LCMS (ESI, pos. ion), *m/z* 350.0 [M+1]; HRMS: Calc'd for [C₁₉H₁₆N₃O₂S]⁺: 350.09577, Found: 350.09676; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.26 (s, 1H), 8.45 (s, 1H), 8.31 (s, 1H), 8.27-8.21 (m, 2H), 8.15 (dd, *J* = 8.8, 2.1 Hz, 1H), 7.99 (dd, *J* = 8.6, 1.8 Hz, 1H), 7.60 (d, *J* = 8.7 Hz, 1H), 7.0 (s, 2H), 4.43 (q, *J* = 7.1 Hz, 2H), 1.41 (t, *J* = 7.1 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.7, 161.9, 161.0, 151.4, 140.2, 139.3, 136.7, 133.8, 133.1, 133.0, 130.9, 126.3, 125.6, 123.6, 125.3, 123.4, 119.6, 61.5, 14.2.

Isopropyl 5-(2-aminoquinazolin-6-yl)benzo[*b*]thiophene-2-carboxylate (1b) General procedure A except with 2.0 equiv. K_2CO_3 in step 1. Isolated 0.145 g (59.6 %), yellow solid. HPLC: 94.9 %; LCMS (ESI, pos. ion), m/z 364.1 $[\text{M}+1]$; HRMS: Calc'd for $[\text{C}_{20}\text{H}_{18}\text{N}_3\text{O}_2\text{S}]^+$: 364.11142, Found: 364.11239; ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 9.16 (s, 1H), 8.34 (s, 1H), 8.18 (s, 1H), 8.16-8.09 (m, 2H), 8.05 (dd, $J = 8.8, 2.0$ Hz, 1H), 7.88 (dd, $J = 8.5, 1.5$ Hz, 1H), 7.50 (d, $J = 8.7$ Hz, 1H), 6.91 (s, 2H), 5.14 (sept, $J = 6.3$ Hz, 1H), 1.32 (d, $J = 6.3$ Hz, 6H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 162.7, 161.5, 161.0, 151.4, 140.2, 139.3, 136.7, 134.3, 133.1, 133.0, 130.7, 126.2, 125.5, 125.3, 123.6, 123.3, 119.6, 69.3, 21.6.

5-(2-Aminoquinazolin-6-yl)benzo[*b*]thiophene-2-carboxylic acid (1c): General procedure A with the following exceptions. Step 1: 1 h at 140 °C, 1:1 MeCN to EtOH (2 mL EtOH added after first 0.5 h). Step 2: 30 min, 90 °C, then 0.15 additional equiv. $\text{Pd}(\text{dppf})\text{Cl}_2$ added and 1 mL additional H_2O , then 95 °C for 30 minutes. Purified by washing with MeOH, CH_2Cl_2 , and 90:10:1 CH_2Cl_2 : CH_3OH : NH_4OH to isolate 0.061 g (23.8 %), yellow solid. HPLC: 89.0 %; LCMS (ESI, pos. ion), m/z 322.0 $[\text{M}+1]$; HRMS: Calc'd for $[\text{C}_{17}\text{H}_{12}\text{N}_3\text{O}_2\text{S}]^+$: 322.06447, Found: 322.06486; ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 9.24 (s, 1H), 8.25-8.11 (m, 3H), 7.97 (d, $J = 8.3$ Hz, 1H), 7.73 (d, $J = 8.3$ Hz, 1H), 7.57 (s, 2H), 7.57-7.53 (m, 1H), 6.94 (s, 1H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 163.8, 162.6, 160.8, 160.7, 151.1, 151.0, 140.9, 139.7, 135.1, 133.9, 133.2, 125.12, 125.07, 123.0, 122.9, 121.7, 119.7.

5-(2-Aminoquinazolin-6-yl)-N-phenylbenzo[b]thiophene-2-carboxamide (1d) General procedure A with following exceptions: Step 2: 1.5 h, 90 °C, 2.0 equiv K₂CO₃, 0.30 equiv. Pd(dppf)Cl₂, 1.5 equiv 6-bromoquinolin-2-amine (**4a**). Purified by washing with MeOH, CH₂Cl₂, and 90:10:1 CH₂Cl₂:CH₃OH:NH₄OH to isolate 0.109 g (41.3 %), tan solid. HPLC: 93.6 %; LCMS (ESI, pos. ion), *m/z* 397.1 [M+1]; HRMS: Calc'd for [C₂₃H₁₆N₄OS]⁺: 397.11176, Found: 397.11187; ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.6 (s, 1H), 9.18 (s, 1H), 8.4-6.8 (m, 12H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.7, 161.0, 160.3, 151.3, 141.2, 140.04, 139.99, 139.45, 139.0, 136.5, 133.2, 128.7, 125.9, 125.6, 125.4, 125.2, 123.8, 123.4, 122.8, 120.5, 119.7.

5-(2-Aminoquinazolin-6-yl)-N-methylbenzo[b]thiophene-2-carboxamide (1e)

General procedure A with the following exceptions: Step 2: 1.5 h, 95 °C, 2.0 equiv K₂CO₃, 0.30 equiv. Pd(dppf)Cl₂, 1.5 equiv 6-bromoquinolin-2-amine (**4a**). Isolated 0.038 g (17 %). Pale yellow solid. HPLC: 96.9 %; LCMS (ESI, pos. ion), *m/z* 335.0 [M+1]; HRMS: Calc'd for [C₁₈H₁₅N₄OS]⁺: 335.09611, Found: 335.09608; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.30 (s, 1H), 8.89 (d, *J* = 4.6 Hz, 1H), 8.39 (s, 1H), 8.30 (d, *J* = 1.9 Hz, 1H), 8.26 – 8.19 (m, 2H), 8.18 (s, 1H), 7.94 (dd, *J* = 8.5, 1.5 Hz, 1H), 7.63 (d, *J* = 8.7 Hz, 1H), 7.03 (s, 2H), 2.93 (d, *J* = 4.6 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 163.7, 161.8, 160.9, 151.3, 141.0, 140.0, 139.0, 136.4, 133.21, 133.19, 125.5, 125.2, 125.0, 124.5, 123.4, 122.6, 119.7, 26.2.

Ethyl 5-(2-aminoquinazolin-6-yl)benzofuran-2-carboxylate (2a) General procedure B. Isolated 0.074 g (33.2 %), yellow solid. HPLC: 95.9 %; LCMS (ESI, pos. ion), m/z 334.1 [M+1]; HRMS: Calc'd for $[C_{19}H_{16}N_3O_3]^+$: 334.11862, Found: 334.11853; 1H NMR (400 MHz, DMSO- d_6): δ 9.24 (s, 1H), 8.19 (d, J = 1.6 Hz, 1H), 8.18 (br s, 1H) 8.12 (dd, J = 8.9, 1.8 Hz, 1H), 7.95 (d, J = 8.0 Hz, 1H), 7.90 (d, J = 8.0 Hz, 1H), 7.86 (s, 1H), 7.58 (d, J = 8.8 Hz, 1H), 6.98 (s, 2H), 4.44 (q, J = 7.1 Hz, 2H), 1.41 (t, J = 7.1 Hz, 3H); ^{13}C NMR (100 MHz, DMSO- d_6): δ 162.7, 160.9, 158.6, 154.53, 154.52, 151.2, 145.8, 135.8, 133.3, 127.4, 127.0, 125.6, 125.2, 120.8, 119.6, 114.3, 112.6, 61.3, 14.1.

5-(2-Aminoquinazolin-6-yl)-N-phenylbenzofuran-2-carboxamide (2b)

General procedure B except initial temperature was 140 °C. Isolated 0.102 g (40.0 %), yellow solid. HPLC: 97.2 %; LCMS (ESI, pos. ion), m/z 381.1 [M+H]; 1H NMR (400 MHz, DMSO- d_6): δ 10.19 (s, 1H), 10.00 (s, 1H), 9.26 (s, 1H), 8.31 (d, J = 2.1 Hz, 1H), 8.19 (d, J = 1.8 Hz, 1H), 8.16 (dd, J = 8.8, 2.0 Hz, 1H), 8.00 (dd, J = 8.8, 2.1 Hz, 1H), 7.61 (d, J = 8.8 Hz, 1H), 7.52 (m, 3H), 7.45 (m, 2H), 7.26 (t, J = 7.3 Hz, 1H); ^{13}C NMR (100 MHz, DMSO- d_6): δ 194.9, 162.4, 160.8, 151.0, 145.5, 139.6, 133.8, 133.6, 132.3, 132.26, 132.27, 129.6, 128.9, 125.2, 123.90, 123.88, 123.86, 121.9, 120.3, 119.7, 114.0.

1-(5-(2-Aminoquinazolin-6-yl)benzofuran-2-yl)(morpholino)methanone (2c) General procedure B with following exceptions: 0.100 g scale. Step 1: 1.5 h, 140° C, 1.5 equiv K₂CO₃. Isolated 0.069 g (55.0 %), yellow solid. HPLC: 99.3 %; LCMS (ESI, pos. ion), *m/z* 375.1 [M+H]⁺; HRMS: Calc'd for [C₂₁H₁₉N₄O₃]⁺: 375.14517, Found: 375.14549; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.15 (s, 1H), 8.09 (d, *J* = 1.9 Hz, 1H), 8.06-7.99 (m, 2H), 7.76-7.82 (m, 1H) 7.74 (d, *J* = 8.8 Hz, 1H), 7.48 (d, *J* = 8.8 Hz, 1H), 7.45 (s, 1H), 6.88 (s, 2H), 3.68-3.60 (m, 4H), 2.51-2.43 (m, 4H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.6, 160.9, 158.8, 153.5, 151.2, 148.8, 135.5, 133.5, 133.3, 127.4, 125.6, 125.5, 125.2, 120.1, 119.7, 112.3, 111.2, 68.2, 41.0 (under DMSO).

1-(5-(2-Aminoquinazolin-6-yl)benzofuran-2-yl)(phenyl)methanone (2d) General procedure B. Isolated 0.180 g (73.5 %), yellow solid. HPLC: 99.3 %; LCMS (ESI, pos. ion), *m/z* 366.0 [M+H]⁺; HRMS: Calc'd for [C₂₃H₁₆N₃O₂]⁺: 366.12370, Found: 366.12371; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 8.23 (d, *J* = 1.4 Hz, 1H), 8.20 (d, *J* = 1.9 Hz, 1H), 8.13 (dd, *J* = 8.8 Hz, 2.1 Hz, 1H), 8.10 (s, 1H), 8.08 (s, 1H), 8.04-7.98 (m, 1H), 7.93-7.98 (m, 1H) 7.90 (s, 1H), 7.81 (t, *J* = 7.5 Hz, 1H), 7.69 (t, *J* = 7.6 Hz, 2H), 7.59 (d, *J* = 8.7 Hz, 1H), 6.99 (s, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 183.5, 162.3, 160.9, 154.8, 152.1, 151.2, 136.7, 135.9, 133.24, 133.23, 133.16, 129.2, 128.8, 127.8, 127.6, 125.6, 125.2, 121.3, 119.6, 117.2, 112.8.

1-(5-(2-Aminoquinazolin-6-yl)benzofuran-2-yl)-2,2-dimethylpropan-1-one (2e)

General procedure B. Isolated 0.166 g (71.6 %), pale yellow solid. HPLC: 99.6 %; LCMS (ESI, pos. ion), m/z 346.1 [M+H]⁺; HRMS: Calc'd for [C₂₁H₁₉N₃O₂]⁺: 346.15500, Found: 346.15589; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.24 (s, 1H), 8.18 (m, 2H), 8.11 (dd, *J* = 8.8, 2.0 Hz, 1H), 7.97-7.94 (m, 1H), 7.92 (s, 1H), 7.90-7.86 (m, 1H), 7.58 (d, *J* = 8.7 Hz, 1H), 6.98 (s, 2H), 1.43 (s, 9H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 195.5, 162.7, 160.9, 153.9, 151.9, 151.2, 135.7, 133.3, 133.2, 127.3, 127.2, 125.5, 125.2, 120.9, 119.6, 114.3, 112.6, 43.0, 26.5.

6-(2-PhenylH-imidazo[1,2-a]pyridin-6-yl)quinazolin-2-amine (3a): General procedure C. Isolated 0.066 g (25.6 %), yellow solid. HPLC: 98.5%; LCMS (ESI, pos. ion), m/z 338.1 [M+H]⁺; HRMS: Calc'd for [C₂₁H₁₆N₅]⁺: 338.14002, Found: 338.14093; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.19 (s, 1H), 8.96 (s, 1H), 8.42 (s, 1H), 8.17 (d, *J* = 1.9 Hz, 1H), 8.08 (dd, *J* = 8.8, 2.1 Hz, 1H), 8.01 (d, *J* = 7.2 Hz, 2H), 7.76-7.65 (m, 2H), 7.55 (d, *J* = 8.7 Hz, 1H), 7.47 (t, *J* = 7.6 Hz, 2H), 7.34 (t, *J* = 7.3 Hz, 1H), 6.97 (s, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.6, 161.0, 151.5, 144.9, 144.1, 134.0, 132.6, 129.9, 128.7, 127.8, 125.6, 125.3, 125.2, 124.8, 124.5, 123.8, 119.6, 116.7, 109.5.

6-(2-(2-Methoxyphenyl)*H*-imidazo[1,2-*a*]pyridin-6-yl)quinazolin-2-amine (3b)

General procedure C with 3.2 equiv. of K₂CO₃ in Step 2. Isolated 0.120 g (43.0 %), tan solid. HPLC: 96.5%; LCMS (ESI, pos. ion), *m/z* 368.1 [M+H]; HRMS: Calc'd for [C₂₂H₁₈N₅O]⁺: 368.15059, Found: 368.15074; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.25 (s, 1H), 9.11 (s, 1H), 8.52 (s, 1H), 8.40 (dd, *J* = 7.7, 1.5 Hz, 1H), 8.23 (d, *J* = 1.8 Hz, 1H), 8.15 (dd, *J* = 8.8, 2.0 Hz, 1H), 7.75 (s, 2H), 7.61 (d, *J* = 8.8 Hz, 1H), 7.41 (m, 1H), 7.15 (t, *J* = 7.5 Hz, 1H), 7.03 (s, 2H), 4.07 (s, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.5, 161.0, 156.5, 151.4, 142.7, 141.6, 132.5, 130.1, 128.5, 128.0, 125.3, 125.0, 124.6, 124.0, 123.9, 122.0, 120.6, 119.5, 116.4, 113.2, 111.5, 55.5.

6-(2-(3-Methoxyphenyl)*H*-imidazo[1,2-*a*]pyridin-6-yl)quinazolin-2-amine (3c)

General procedure C with 3.2 equiv of K₂CO₃ in Step 2. Isolated 0.112 g (42.2 %), peach solid. HPLC: 97.5%; LCMS (ESI, pos. ion), *m/z* 368.1 [M+H]; HRMS: Calc'd for [C₂₂H₁₈N₅O]⁺: 368.15059, Found: 368.15081; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.26 (s, 1H), 9.02 (s, 1H), 8.51 (s, 1H), 8.24 (d, *J* = 2.0 Hz, 1H), 8.15 (dd, *J* = 8.8, 2.0 Hz, 1H), 7.77 (m, 2H), 7.64 (m, 3H), 7.45 (t, *J* = 8.1 Hz, 1H), 7.04 (s, 2H), 6.99 (m, 1H), 3.92 (s, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.6, 161.0, 159.6, 151.4, 144.8, 144.0, 135.2, 132.6, 129.9, 129.8, 125.3, 125.2, 124.8, 124.6, 123.8, 119.6, 118.0, 116.8, 113.6, 110.7, 109.9, 55.1.

6-(2-(4-Methoxyphenyl)*H*-imidazo[1,2-*a*]pyridin-6-yl)quinazolin-2-amine (3d)

General procedure C. Isolated 0.101 g (36.3 %), peach solid. HPLC: 96.3 %; LCMS (ESI, pos. ion), m/z 368.1 [M+H]⁺; HRMS: Calc'd for [C₂₂H₁₈N₅O]⁺: 368.15059, Found: 368.15075; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.23 (s, 1H), 8.98 (s, 1H), 8.36 (s, 1H), 8.21 (d, *J* = 1.8 Hz, 1H), 8.13 (dd, *J* = 8.8, 2.0 Hz, 1H), 7.98 (d, *J* = 8.7 Hz, 2H), 7.72 (s, 2H), 7.59 (d, *J* = 8.7 Hz, 1H), 7.08 (d, *J* = 8.7 Hz, 2H), 7.01 (s, 2H), 3.86 (s, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.5, 161.0, 159.1, 151.4, 145.0, 144.0, 132.6, 130.0, 126.9, 126.4, 125.3, 125.1, 124.4, 124.3, 123.7, 119.6, 116.5, 114.1, 108.5, 55.1.

6-(2-(Thiophen-3-yl)*H*-imidazo[1,2-*a*]pyridin-6-yl)quinazolin-2-amine (3e)

General procedure C. Isolated 0.131 g (50.3 %), orange solid. HPLC: 97.3 %; LCMS (ESI, pos. ion), m/z 344.0 [M+H]⁺; HRMS: Calc'd for [C₁₉H₁₄N₅S]⁺: 344.09644, Found: 344.09690; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.23 (s, 1H), 9.00 (s, 1H), 8.33 (s, 1H), 8.22 (d, *J* = 1.9 Hz, 1H), 8.13 (dd, *J* = 8.7, 2.1 Hz, 1H), 8.01 (t, *J* = 2.0 Hz, 1H), 7.73 (s, 2H), 7.70 (d, *J* = 2.0 Hz, 2H), 7.59 (d, *J* = 8.8 Hz, 1H), 7.02 (s, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.5, 161.0, 159.1, 151.4, 145.0, 144.0, 132.6, 130.0, 126.9, 126.4, 125.3, 125.1, 124.4, 124.3, 123.7, 119.6, 116.5, 114.1, 108.5, 55.1.

6-(2-(Benzo[d][1,3]dioxol-5-yl)*H*-imidazo[1,2-*a*]pyridin-6-yl)quinazolin-2-amine (3f)

General procedure C. Isolated 0.077 g (26.6 %), peach solid. HPLC: 94.9 %; LCMS (ESI, pos. ion), m/z 382.1 [M+H]⁺; HRMS: Calc'd for [C₂₂H₁₆N₅O₂]⁺: 382.12985, Found: 382.13076; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.04 (s, 1H), 8.79 (s, 1H), 8.19 (s, 1H), 8.03 (d, *J* = 2.1 Hz, 1H), 7.95 (dd, *J* = 8.8, 2.2 Hz, 1H), 7.54 (d, *J* = 1.1 Hz, 2H), 7.44-7.37 (m, 3H), 6.87 (d, *J* = 8.5 Hz, 1H), 6.82 (s, 2H), 5.94 (s, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.5, 161.0, 151.4, 147.7, 147.0, 144.9, 143.9, 132.6, 130.0, 128.1, 125.3, 125.2, 124.6, 124.4, 123.7, 119.6, 119.3, 116.5, 108.9, 108.6, 105.9, 101.1.

6-(2-Ethyl*H*-imidazo[1,2-*a*]pyridin-6-yl)quinazolin-2-amine (3g)

General procedure C. Isolated 0.085 g (25.7 %), yellow solid. HPLC: 98.9 %; LCMS (ESI, pos. ion), m/z 290.1 [M+H]⁺; HRMS: Calc'd for [C₁₇H₁₆N₅]⁺: 290.14002, Found: 290.14007; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.23 (s, 1H), 8.96 (s, 1H), 8.19 (d, *J* = 1.8 Hz, 1H), 8.10 (dd, *J* = 8.8, 2.0 Hz, 1H), 7.78 (s, 1H), 7.71-7.56 (m, 3H), 7.01 (s, 2H), 2.79 (q, *J* = 7.5 Hz, 2H), 1.34 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.5, 160.9, 151.3, 149.1, 143.3, 132.6, 130.2, 125.3, 125.0, 123.74, 123.68, 123.5, 119.6, 116.1, 109.4, 21.7, 13.5.

6-(2-*tert*-Butyl*H*-imidazo[1,2-*a*]pyridin-6-yl)quinazolin-2-amine (3h) General procedure C, Isolated 0.193 g (53.5 %), pale yellow solid. HPLC: 99.4 %; LCMS (ESI, pos. ion), m/z 318.1 [M+H]; HRMS: Calc'd for [C₁₉H₂₀N₅]⁺: 318.17132, Found: 318.17162; ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.27 (s, 1H), 8.97 (s, 1H), 8.21 (d, J = 2.0 Hz, 1H), 8.12 (dd, J = 8.8, 2.1 Hz, 1H), 7.80 (s, 1H), 7.70 (s, 2H), 7.63 (d, J = 8.8 Hz, 1H), 7.04 (s, 2H), 1.45 (s, 9H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.5, 160.9, 156.7, 151.3, 143.2, 132.6, 130.3, 125.3, 125.0, 123.72, 123.69, 123.64, 119.6, 116.3, 107.6, 32.1, 30.1.

¹H and ¹³C NMR Spectra of boronates 5a, 6a, 7a

Ethyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[b]thiophene-2-carboxylate (5a)

No.	(ppm)	Height	No.	(ppm)	Value	Absolute Value	Non-Negative Value
1	1.37	1.0000	1	[1.30 .. 1.45]	14.384	1.46757e+10	14.384
2	1.38	0.1998	2	[4.34 .. 4.48]	2.000	2.04059e+9	2.000
3	1.40	0.0937	3	[7.75 .. 7.84]	0.936	9.55415e+8	0.936
4	4.39	0.0812	4	[8.04 .. 8.13]	0.969	9.88571e+8	0.969
5	4.41	0.0810	5	[8.27 .. 8.35]	0.968	9.87458e+8	0.968
6	7.79	0.0390	6	[8.37 .. 8.48]	0.976	9.96136e+8	0.976
7	7.81	0.0449					
8	8.08	0.0499					
9	8.10	0.0428					
10	8.30	0.1014					
11	8.43	0.0794					

**Ethyl
5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo[b]thiophene-2-carboxylate (5a)**

No.	(ppm)	Height
1	14.15	0.1859
2	24.69	0.5542
3	61.51	0.1457
4	83.87	0.2916
5	122.49	0.1520
6	131.08	0.1363
7	132.04	0.1355
8	132.78	0.1370
9	133.05	0.0981
10	138.25	0.1095
11	144.09	0.0945
12	161.91	0.0931

**Ethyl
5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzofuran-2-carboxylate
(6a)**

1H NMR (400 MHz, DMSO-d₆)

No.	(ppm)	Height	No.	(ppm)	Value	Absolute Value	Non-Negative Value
1	1.38	1.0000	1	[1.36 .. 1.44]	15.000	9.50247e+9	15.000
2	1.40	0.1872	2	[4.38 .. 4.48]	2.064	1.30756e+9	2.064
3	1.42	0.0905	3	[7.75 .. 7.80]	0.959	6.07247e+8	0.959
4	4.40	0.0255	4	[7.82 .. 7.87]	1.969	1.24715e+9	1.969
5	4.42	0.0785	5	[8.22 .. 8.26]	0.948	6.00589e+8	0.948
6	4.44	0.0778					
7	4.46	0.0249					
8	7.77	0.0301					
9	7.79	0.0553					
10	7.84	0.0532					
11	7.85	0.0995					
12	7.86	0.0343					
13	8.23	0.0758					

**Ethyl
5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzofuran-2-carboxylate
(6a)**

No.	(ppm)	Height
1	14.10	0.1696
2	24.66	0.5734
3	61.26	0.1285
4	83.82	0.2747
5	111.74	0.1463
6	114.21	0.1237
7	126.55	0.0933
8	130.48	0.1415
9	133.45	0.1333
10	145.25	0.0891
11	156.92	0.0914
12	158.57	0.0801

2-Phenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)H-imidazo[1,2-a]pyridine hydrobromide (7a)

¹H NMR (400 MHz, DMSO-d₆)

No.	(ppm)	Height	No.	(ppm)	Value	Absolute Value	Non-Negative Value
1	1.45	1.0000	1	[1.39 .. 1.52]	12.000	1.01591e+10	12.000
2	7.62	0.0141	2	[7.58 .. 7.75]	3.525	2.98427e+9	3.525
3	7.64	0.0437	3	[7.99 .. 8.07]	4.265	3.61071e+9	4.265
4	7.65	0.0380	4	[8.85 .. 8.90]	0.985	8.33653e+8	0.985
5	7.68	0.0571	5	[9.17 .. 9.22]	0.960	8.12371e+8	0.960
6	7.70	0.0824					
7	7.72	0.0314					
8	8.03	0.0825					
9	8.05	0.0777					
10	8.06	0.1658					
11	8.88	0.1159					
12	9.20	0.0840					

**2-Phenyl-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)H-imidazo[1,2-a]
pyridine hydrobromide (7a)**

No.	(ppm)	Height
1	24.67	0.3108
2	73.50	0.0162
3	84.84	0.2008
4	111.15	0.0633
5	112.02	0.0616
6	126.21	0.1692
7	126.56	0.0345
8	129.50	0.1647
9	130.41	0.0591
10	134.94	0.0631
11	135.91	0.0295
12	136.38	0.0404
13	141.22	0.0438

HPLC of 1a-e, 2a-e, 3a-h

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254.8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.678	PB	0.1240	275.72260	35.74591	98.1250
2	10.472	MM	0.0625	5.26859	1.40461	1.8750

Totals : 280.99118 37.15051

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.426	VV	0.0931	9.74244	1.43777	0.3899
2	6.690	VP	0.1477	11.91422	1.22228	0.4768
3	7.100	VB	0.1536	2448.59766	255.91881	97.9833
4	9.315	MM	0.0824	6.81434	1.37785	0.2727
5	10.472	MM	0.0613	21.92715	5.95753	0.8774

Totals : 2498.99580 265.91424

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	3.284	PB	0.0557	5.66309	1.58335	0.6398
2	5.086	BB	0.0751	787.90692	160.97371	89.0108
3	7.393	BB	0.0552	39.15524	11.09648	4.4234
4	8.350	BB	0.0555	29.38782	8.25367	3.3200
5	10.580	MM	0.0566	23.06849	6.78846	2.6061

Totals : 885.18156 188.69567

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.262	VV	0.0780	39.74881	7.73500	0.5822
2	6.663	VB	0.1083	6390.59668	928.89618	93.6059
3	9.842	VB	0.0705	396.78247	81.87730	5.8119

Totals : 6827.12796 1018.50848

1d

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.865	VB	0.0680	155.80443	34.97473	96.8705
2	10.471	MM	0.0720	5.03348	1.16572	3.1295

Totals : 160.83791 36.14045

1e

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254.8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	5.416	BV	0.0682	40.83327	9.48478	0.9645
2	5.684	VV	0.0902	37.27158	5.86219	0.8804
3	5.786	VP	0.0820	90.10445	16.42474	2.1283
4	6.139	BB	0.1005	4059.55225	635.95416	95.8866
5	6.905	VB	0.0772	5.93805	1.13278	0.1403

Totals : 4233.69959 668.85865

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

2b

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.937	BB	0.0589	28.77346	7.47882	0.7417
2	5.418	BP	0.0678	34.51199	8.08645	0.8896
3	6.324	PB	0.0891	46.62756	9.22359	1.2020
4	6.788	PB	0.1129	3769.38110	517.94885	97.1667

Totals : 3879.29412 541.73771

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.856	BB	0.0638	1784.20581	435.36487	99.2554
2	5.331	BB	0.0869	7.30997	1.16919	0.4067
3	6.901	PB	0.0569	6.07507	1.65255	0.3380

Totals : 1797.59085 438.18660

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254.8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	5.425	BP	0.0688	30.02914	6.89334	0.6771
2	6.595	FB	0.1164	4404.94629	594.88086	99.3229

Totals : 4434.97543 601.77420

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	5.412	BP	0.0671	17.72922	4.21602	0.3562
2	6.942	PB	0.1273	4959.49463	620.00189	99.6438

Totals : 4977.22385 624.21791

2e

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime (min)	Type	Width (min)	Area [mAU*s]	Height [mAU]	Area %
1	4.068	BB	0.0524	5.01093	1.52354	0.5727
2	4.393	PB	0.0530	861.55389	257.74030	98.4625
3	5.088	VP	0.0538	8.44258	2.47595	0.9649

Totals : 875.00741 261.73978

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

3b

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.362	BP	0.0509	43.42863	13.03276	2.9959
2	4.679	BV	0.0515	1398.23914	412.88303	96.4575
3	5.727	PB	0.0804	7.92325	1.58411	0.5466

Totals : 1449.59102 427.49990

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.678	PB	0.1240	275.72260	35.74591	98.1250
2	10.472	MM	0.0625	5.26859	1.40461	1.8750

Totals : 280.99118 37.15051

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254.8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.351	MM	0.0693	5.32416	1.28136	0.6508
2	4.643	PV	0.0535	797.49554	235.33218	97.4827
3	4.926	VV	0.0703	8.45757	1.69194	1.0338
4	5.048	VB	0.0870	6.81245	1.11934	0.8327

Totals : 818.08972 239.42482

3c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.306	BP	0.0560	18.54169	4.91304	1.1305
2	4.600	PV	0.0510	1579.34692	473.03860	96.2922
3	4.991	VB	0.0948	9.03824	1.33826	0.5511
4	5.457	FP	0.0538	14.91521	4.36746	0.9094
5	6.582	VB	0.0540	18.31887	5.34808	1.1169

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	3.910	BB	0.0485	15.67279	5.01620	1.7791
2	4.245	PB	0.0515	856.95929	252.91953	97.2759
3	4.722	BV	0.0640	8.32560	1.94160	0.9451

Totals : 880.95768 259.87732

3e

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	3.536	MM	0.0651	7.25140	1.85537	0.3219
2	4.180	PB	0.0512	82.41051	25.84863	3.6585
3	4.486	PB	0.0515	2138.44434	632.36823	94.9339
4	10.579	PB	0.0564	24.45484	6.73430	1.0856

Totals : 2252.56110 666.80652

3f

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	2.848	BB	0.0548	11.24249	3.21230	0.2889
2	3.525	BV	0.0505	3848.74634	1166.93652	98.8938
3	3.745	VB	0.0740	31.80764	5.98356	0.8173

Totals : 3891.79647 1176.13239

3g

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	3.812	BB	0.0516	6.64742	1.96022	0.1421
2	4.176	PB	0.0538	4651.64795	1363.46704	99.4183
3	4.830	VV	0.0772	11.52369	2.19826	0.2463
4	6.435	BB	0.0747	9.04462	1.86193	0.1933

Totals : 4678.86368 1369.48745

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

7a

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.480	PB	0.0644	5640.39600	1304.75793	99.3724
2	6.217	BB	0.0687	35.62379	7.88648	0.6276

Totals : 5676.01979 1312.64441

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,8 Ref=off

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	4.607	BV	0.0849	331.43057	59.62387	5.1139
2	4.859	VV	0.0752	6149.56885	1254.45825	94.8861

Totals : 6480.99942 1314.08213

12

References

- (1) Hynes, J. B.; Campbell, J. P. *J. Het. Chem.* **1997**, *34*, 385-387.
- (2) Holland, R.; Spencer, J.; Deadman, J. J. *Synthesis* **2002**, 2379-2382.