This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Mid- to late Holocene geomorphological and hydrological changes in the south Taihu area of the Yangtze delta plain, China

PLEASE CITE THE PUBLISHED VERSION

<https://doi.org/10.1016/j.palaeo.2018.03.012>

PUBLISHER

© Elsevier

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at:
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Chen, Ting, David Ryves, Zhanghua Wang, Jonathan Lewis, and Xuening Yu. 2019. "Mid- to Late Holocene Geomorphological and Hydrological Changes in the South Taihu Area of the Yangtze Delta Plain, China". figshare. <https://hdl.handle.net/2134/32548>.

1 **Mid- to late Holocene morphological and hydrological changes in the south**
2 **Taihu area of the Yangtze delta plain, China**

4 Ting Chen^{a, b}, David B. Ryves^c, Zhanghua Wang^{a*}, Jonathan P. Lewis^c, Xuening
5 Yu^a

^a *State Key Laboratory of Estuarine and Coastal Research, East China Normal University, Shanghai 200062, China*

^b Academy for Advanced Interdisciplinary Studies, Southern University of
Science and Technology, Shenzhen 518055, China

^c Centre for Hydrological and Ecosystem Science, Department of Geography,
Loughborough University, Loughborough LE11 3TU, UK

14 *Corresponding author:

15 *Zhanghua Wang, State Key Laboratory of Estuarine and Coastal Research, East*
16 *China Normal University, Shanghai 200062, China.*

17 *Tel: +86 13817202106*

18 *E-mail address: zhwang@geo.ecnu.edu.cn*

20 **Abstract**

The Taihu Plain of the Lower Yangtze valley, China was a centre of rice agriculture during the Neolithic period. Reasons for the rapid development of rice cultivation during this period, however, have not been fully understood for this coastal lowland, which is highly sensitive to sea-level change. To improve understanding of the morphological and hydrological context for evolution of prehistoric rice

agriculture, two sediment cores (DTX4 and DTX10) in the East Tiaoxi River Plain, south Taihu Plain, were collected, and analysed for radiocarbon dating, diatoms, organic carbon and nitrogen stable isotopes ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$), grain size and lithology. These multiproxy analyses revealed that prior to ca. 7500 cal. yr BP, the East Tiaoxi River Plain was a rapidly aggrading high-salinity estuary (the Palaeo-Taihu Estuary). After ca. 7500 cal. yr BP, low salinity conditions prevailed as a result of strong Yangtze freshwater discharge. Subsequently, seawater penetration occurred and saltmarsh developed between ca. 7000 and 6500 cal. yr BP due to accelerated relative sea-level rise. This transgression event influenced a large area of the Taihu Plain during the Holocene, as shown by multiple sediment records from previous studies. Persistent freshwater marsh (or subaerial land) formed due to dramatic shrinkage/closure of the Palaeo-Taihu Estuary after ca. 5600 cal. yr BP when sea level was relatively stable. We speculate that morphological and hydrological changes of the East Tiaoxi River Plain played an important role in agricultural development across the Taihu Plain during the Neolithic period. The closure of the Palaeo-Taihu Estuary and the formation of stable freshwater marsh (or subaerial land) after ca. 5600 cal. yr BP were critical preconditions encouraging the rapid rise of rice productivity in the Liangzhu period (5500-4500 cal. yr BP). This development changed the landscape and river systems, and thus provided adequate freshwater supply to the Taihu Plain.

Key words

Coastal wetland; Salinity; Sea level; Freshwater resource; Rice agriculture, Multiproxy analyses

1. Introduction

51 The lower Yangtze valley, East China is one of the centres where intense rice
52 growth started in the Neolithic period (Zong et al., 2007; Liu and Chen, 2012). Many
53 studies have been focused on the origin and domestication of rice farming in this
54 region during the Neolithic cultural period (Zong et al., 2007; Mo et al., 2011; Liu and
55 Chen, 2012). The first evidence of collection and consumption of wild rice (ca.
56 9000-10000 cal. yr BP) was found at Shangshan site, in Zhejiang Province (Liu and
57 Chen, 2012; Zuo et al., 2017). Evidence for rice cultivation also found at the
58 Kuahuqiao site at ca. 7700 cal. yr BP (Zong et al., 2007). Although marine inundation
59 caused by sea-level rise brought rice cultivation to an end at the Kuahuqiao site at ca.
60 7500 cal. yr BP (Zong et al., 2007), rice farming subsequently expanded to the south
61 of the Hangzhou Bay and in the Taihu Plain, north of Hangzhou Bay (Chen et al.,
62 2008; Zong et al., 2012a; Zheng et al., 2012). Rice domestication started during the
63 period of the Majiabang culture (7000-5800 cal. yr BP), the first Neolithic culture that
64 appeared on the Taihu Plain, though this society still relied mostly on hunting, fishing
65 and gathering (Cao et al., 2006; Fuller et al., 2007; Mo et al., 2011; Zong et al., 2012a;
66 Xu, 2015). More rice cultivation was practiced during the Songze period (5800-5500
67 cal. yr BP), but hunting, fishing and gathering were still important at that time (Cao et
68 al., 2006; Mo et al., 2011). Subsequently, rice farming intensified and its yield
69 increased dramatically during the Liangzhu culture (5500-4500 cal. yr BP) (Fan, 2011;
70 Mo et al., 2011; Zhuang et al., 2014; Zhang et al., 2015). For example, substantial
71 quantities of carbonised rice were found at several Liangzhu sites (Fan, 2011), one of
72 which even reached to tens of thousands of kilograms; numerous specialised fine
73 stone tools used for rice farming, such as ploughs and sickles, were also found (Mo et
74 al., 2011). Relying on rapid advances in rice farming, the Liangzhu culture developed
75 into a sophisticated and complex society and was considered as one of the most

advanced Neolithic societies in the world (Jiang and Liu, 2006; Zhu, 2006; Lawler, 2009; Mo et al., 2011; Zong et al., 2012a; Zhuang et al., 2014).

Several studies suggested that a warm/humid climate promoted the development of rice farming in the lower Yangtze delta (Yu et al., 2000; Chen et al., 2005; Innes et al., 2009, 2014; Patalano et al., 2015), together with ancient people's successful water and landscape management of rice paddies (Zong et al., 2007; Zhuang et al., 2014). Recently, more attention has been paid to the role of the hydrological environment, which is the result of a complex of sea-level, climate and geomorphological conditions (Zong et al., 2007; Qin et al., 2011; Zheng et al., 2012; Long et al. 2014; Patalano et al., 2015). For example, Zong et al. (2007) found that freshwater low-land swamps at the Kuahuqiao site were selected by Neolithic people for rice cultivation. Qin et al. (2011) reported that rice-based agriculture occurred during two intervals of lower salinity, between ca. 7850-7210 cal. yr BP and ca. 3000-2290 cal. yr BP. Although these studies of hydrological background based on individual cores provided insights into our understanding of the development of rice farming in the lower Yangtze valley, new studies on key sites allowing an integrated analysis of previous studies are still required to provide a more detailed environmental context on a regional scale.

We focus here on the hydrological changes during the mid- to late-Holocene in the East Tiaoxi River Plain, part of the southern Taihu Plain (Fig. 1A and 1B). The East Tiaoxi River Plain is a critical part of the Taihu plain, because it was a palaeo-incised valley during the Last Glacial Maximum (LGM) and was occupied by the Palaeo-Taihu Estuary during the early- to mid-Holocene (Fig. 1C; Hong, 1991). Through this estuary, sea water could have reached the centre of the Taihu Plain, and freshwater from the west uplands, which flows into the Taihu lake at present, instead was discharged into Hangzhou Bay (Hong, 1991). Such hydrologic conditions had

restricted the freshwater resource for Neolithic people in the Taihu Plain. Morphological and hydrological changes within the East Tiaoxi River Plain from the Mid-Holocene are therefore potentially critical to understanding the evolution of rice farming in the Taihu Plain over the Neolithic period.

Fig. 1 Maps of the Taihu Plain and location of study sites. (A) Map of the present Taihu Plain (after Song et al., 2013), showing geomorphology, hydrology and locations of core DTX4 and DTX10, and cores collected from previous studies (red solid circles; references in supplementary Table 1). (B) Map of the East Tiaoxi River Plain, including West and East Tiaoxi River and locations of cores DTX4 and DTX10. (C) Palaeotopographical map of the Taihu Plain during the Last Glacial Maximum (after Wang et al., 2012), showing the Palaeo-Taihu valley (blue solid line) and Palaeo-Taihu Estuary (green solid line). (D) and (E) Remote sensing image in 2016, around Luosheyang Lake and Qianshanyang Lake from Google Earth, showing part of East Tiaoxi River and locations of core DTX4 and DTX10, respectively.

A complete understanding of the evolution of morphological and hydrological environments of the East Tiaoxi River Plain during the mid- to late-Holocene, however, has been hampered by a lack of high-quality sediment archives with reliable dating allowing detailed sedimentological analysis. To address this gap and fulfill the research aim mentioned above, two sediment cores (DTX 4 and DTX10), spanning the last 7100 and 7600 years respectively, were collected beside the Luosheyang Lake and the Qianshanyang Lake in the East Tiaoxi River Plain in 2014 (Fig. 1). Using these two cores, we studied the past morphological and hydrological environments of the East Tiaoxi River Plain over the last 7600 years, by using a multi-proxy approach including lithology, grain size, diatom analysis and organic geochemistry (including stable isotopes of carbon and nitrogen: $\delta^{13}\text{C}$, $\delta^{15}\text{N}$). Additionally, we compared these new results with the hydrological environment inferred from other parts of the Taihu Plain from previously studied cores (Itzstein-Davey et al., 2007; Atahan et al., 2008; Zong et al., 2011, 2012b; Wang et al., 2012; Innes et al., 2014; Liu et al., 2015). Based on these results, we discuss how changing morphological and hydrological conditions

in the East Tiaoxi River Plain influenced the expansion of rice cultivation in the Taihu Plain over the Neolithic period.

2. Study area and site description

During the LGM, the landscape of the Taihu Plain was characterised by a series of river terraces (T1, T2 and T3) and incised valleys (Fig. 1C) cutting through these terraces (Yan and Huang, 1987; Li et al., 2000, 2002; Wang et al., 2012). The terraces T1, T2 and T3 were regions where the thickness of Holocene sediments were at 20-30 m, 5-15 m, and < 5 m, respectively (Yan and Huang, 1987; Li et al., 2000, 2002; Wang et al., 2012). The largest two of these palaeo valleys were the Yangtze valley in the north and the Qiantang valley in the south. The Palaeo-Taihu valley lay along the western highlands, and freshwater discharge from western mountains flowed through it southwards into the Palaeo-Qiantang valley (Yan and Huang, 1987; Hong, 1991; Wang et al., 2012). When sea level rose from -18 m to -4 m between ca. 8600 cal. yr BP and ca. 7300 cal. yr BP, the south part of the Palaeo-Taihu valley became an estuary, allowing sea water from Hangzhou Bay to reach the central Taihu Plain (Hong, 1991; Wang et al., 2012).

Present-day geomorphological and hydrological conditions of the Taihu Plain, however, are dramatically different. Today, the Taihu Plain is characterised by a saucer-like depression, in the centre of which lies Taihu Lake (Fig. 1A), the third largest freshwater lake in China. Taihu Lake receives freshwater inflows (e.g. the East and West Tiaoxi Rivers; Fig. 1A) from the mountainous areas to the west of the Taihu Plain and provides an important freshwater resource for the human population in the Taihu Plain.

The East Tiaoxi River Plain is the southern part of the Taihu Plain and lies between Taihu Lake in the north and the Qiantang River/Hangzhou Bay in the south

(Fig. 1A and 1B). Relief is slightly higher (2-5 m) in the south and east, and lower (< 2 m) in the depression near Taihu Lake. The East Tiaoxi River, the largest river in the East Tiaoxi River Plain, flows northward into Taihu Lake after joining with the West Tiaoxi River in the city of Huzhou.

Luosheyang Lake and Qianshanyang Lake are located near the East Tiaoxi River (Fig. 1B, 1D and 1E), and are about 140 km west of Shanghai. They are both naturally open, shallow (< 2 m maximum depth) and flat-bottomed freshwater lakes, through which the East Tiaoxi River and its tributaries flow. Over recent decades, the area of the Luosheyang Lake has been reduced due to extensive marginal development for agriculture (e.g. rice paddies) to about 1.56 km², while the Qianshanyang Lake has been almost completely reclaimed.

3. Material and methods

3.1 Coring, sampling, AMS ¹⁴C dating and age-depth model

Both cores DTX4 and DTX10, 5.7 and 4.6 m long respectively, were obtained in May 2014 using a gouge corer (Eijkelpkamp Company, the Netherlands), with a diameter of 3 cm. Core DTX4 (30°38'16.465" N, 120°05'28.546" E) was collected from a reclaimed agricultural field which was previously the edge of the Luosheyang Lake, and core DTX10 from a rice paddy near present Qianshanyang Lake (Fig. 1B, 1D and 1E). Lithology of both cores was examined carefully during the drilling, including particle composition, structure, colour, and presence of plant macrofossils. Recent cultural sediments at the top of each core were removed (the uppermost 80 cm at DTX4 and 20 cm at DTX10). Ten plant macrofossils (excluding root material) and organic-rich samples from the two cores were dated via AMS ¹⁴C dating by Beta Analytic, USA (Table 1). All conventional dates were calibrated using the INTCAL 13

database (Talma and Vogel, 1993; Reimer et al., 2009) and calibrated dates with two sigma were presented as ‘medial point ages $\pm$ standard deviation’ (Table 1). A linear interpolation method was used to construct the age-depth model, using the program ‘Clam’ with 10,000 iterations (Blaauw, 2010; Fig. 2). The sedimentation rate (SR), the best estimated age and the minima and maxima of age confidence intervals were calculated every 2 cm (core DTX4) and 4 cm (DTX10) simultaneously, using the ‘Clam’ program. In addition, , a sedimentation hiatus was identified in core DTX10 at a sediment depth of 1.0 m due to a leaching structure indicative of pedogenesis (0.88-1.0 m).

Table 1 AMS ^{14}C dating results of core DTX4 and DTX10

Core name	Depth (m)	Dated material	$\delta^{13}\text{C}$ (‰)	Conventional age (yr BP)	Calibrated age (2 sigma, cal. yr BP)	Median Calibrated age (cal. yr BP)	Lab. code
Core DTX4	1.21	Plant fragments	-27.2	1610 $\pm$ 30	1560–1410	1485 $\pm$ 75	Beta-383485
	1.45	Plant fragments	-28.8	940 $\pm$ 30	930–785	858 $\pm$ 73	Beta-406458
	1.81	Plant fragments	-27.8	2860 $\pm$ 30	3065–2920	2993 $\pm$ 73	Beta-383486
	1.91	Plant fragments	-26.4	4680 $\pm$ 30	5575–5550	5562 $\pm$ 13	Beta-406459
	2.5	Organic-rich mud	-25.7	5680 $\pm$ 30	6500–6405	6453 $\pm$ 48	Beta-382369
	5.33	Plant fragments	NA	6150 $\pm$ 30	7160–6950	7055 $\pm$ 105	Beta-383487
Core DTX10	0.88	Plant fragments	NA	2720 $\pm$ 30	2870–2760	2815 $\pm$ 55	Beta-382371
	2.3	Plant fragments	-27.6	6650 $\pm$ 30	7580–7480	7530 $\pm$ 50	Beta-382366
	2.91	Organic-rich mud	-25.1	8380 $\pm$ 30	9475–9400	9438 $\pm$ 38	Beta-385582
	4.25	Plant fragments	-26.1	6780 $\pm$ 40	7680–7575	7628 $\pm$ 53	Beta-385583

196

197 **Fig. 2** Age-depth model, sedimentation rate (SR) and density distribution of AMS ^{14}C dates (shaded in
 198 yellow) basing on the ‘Clam’ program (Blaauw, 2010), and lithology profile for core DTX4 and
 199 DTX10. A sedimentation hiatus was identified at 1.0 m in core DTX10 because of clear indications of
 200 pedogenesis (leaching) apparent in the lithology between 0.88-1.0 m.

Sediment subsamples were selected at 2 to 10 cm intervals, using 2 to 5 cm thick slices, with thickness chosen according to sedimentation rate: in core DTX4, every 4 cm interval (4 cm thick) from 0.8-1.2 m, 5 cm (5 cm thick) from 1.2-1.8 m, 2 cm (2 cm thick) from 1.8-2.14 m, 3 cm (3 cm thick) from 2.14-2.5 m, and 10 cm (5 cm thick) from 2.5-5.7 m; in core DTX10, every 4 cm (4 cm thick) from 0.2-2.3 m, and 10 cm (5 cm thick) from 2.3-4.6 m. A total of 84 and 77 subsamples were obtained for core DTX4 and DTX10 respectively. All subsamples were stored in a fridge at 4 °C prior to analyses.

3.2 Diatom analysis

Diatom concentrations were low in most samples, with a total of only 29 samples in core DTX4 and 10 in core DTX10 containing diatoms at high enough concentration to permit using the standard water bath method (Renberg, 1990; Battarbee et al., 2001). Due to extremely low diatom concentrations, 8 samples in core DTX4 and 19 in core DTX10, were treated with heavy liquid separation (sodium polytungstate, SPT) using the following procedures (Battarbee and Kneen, 1982; Battarbee et al., 2001). After organic matter and carbonate were removed using hydrogen peroxide (H₂O₂) and HCl, clay grains were removed by adding a few drops of weak NH₃ solution (Battarbee et al., 2001). Then, the non-toxic heavy liquid SPT (3Na₂WO₄·9H₂O) with a density of 2.26 g ml⁻¹ was used twice to separate diatoms from mineral grains with a density above this. Diatom slides were made in the regular way using the diatom-enriched supernatant taken from the top of SPT solutions, after washing with distilled water. A known number of microspheres were added to all samples to assess diatom concentration (Battarbee and Kneen, 1982). The average number of diatoms counted for samples prepared with regular water-bath and SPT method is 266 and 269 valves, respectively, in core DTX4, with the exception of one sample at 5.675 m, for

which only 111 valves were counted due to extremely low diatom concentration. The number of diatoms counted for regularly treated samples was lower in core DTX10 than that in core DTX4 due to lower diatom concentration, but counts of at least 174 valves were obtained with an average of 192, while for SPT treated samples, over 270 valves with an average of 331 were counted. Valves were identified to species level where possible using general (Krammer and Lange-Bertalot, 1986, 1988, 1991a, 1991b; <http://westerndiatoms.colorado.edu/>) and more specialised coastal floras (Witkowski et al., 2000). Diatoms were also classified into their salinity (freshwater, brackish and marine) and habitat preferences (planktonic, benthic), based on the literature and website resources (e.g. van der Wuff and Huls, 1976; Vos and de Wolf, 1993; Ryves et al., 2004; <http://craticula.ncl.ac.uk/Molten/jsp/>; <http://www.marinespecies.org/>) and ecological knowledge.

In most samples some signs of diatom dissolution were apparent: e.g. thinner valve walls in the sub-fossil material. Therefore, the state of dissolution for each valve was recorded and the F-index, the ratio of pristine valves to the sum of pristine and dissolved valves, was calculated for each sample (Ryves et al., 2001). Even where the standard water-bath method was suitable, and diatom concentrations were higher (generally in the fresh water phases), preservation was judged not consistently good enough to support applying a salinity model (which remain robust; Juggins, 2013), as poor preservation is known to cause bias in model outputs (e.g. Ryves et al. 2006, 2009). Percentage abundances of diatom data were calculated and plotted using the stratigraphic software package C2 (Juggins, 1991-2009). Diatom zones were determined based on cluster analysis using the CONISS function in the programme Tilia 2.0.41.

3.3 Grain size analysis

Grain size analyses were conducted on each subsample. These samples were firstly pretreated with 10 % H₂O₂ and then 10 % HCl to remove organic matter and carbonates respectively, and then washed in distilled water to remove residual HCl. Following this, 5 ml of 5 % Calgon® (sodium hexametaphosphate) was added to each sample before shaking in an ultrasonic bath for 15 minutes to prevent flocculation of fine-grained particles (Beuselinck et al., 1998). Measurements were performed with a Beckman Coulter Laser Diffraction Particle Size Analyzer (LS13320).

3.4 Measurement of carbon and nitrogen element and their stable isotopes

Samples for total organic carbon (TOC), total nitrogen (TN), $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis were firstly freeze-dried, milled and sieved with a 74 μm (200 mesh) sieve. The fine fraction was collected and a subsample treated with 1M HCl to remove carbonate followed by washing for four to five times with distilled water before drying in an oven at 45 °C. Samples without additional treatment of HCl were used to measure TC and TN, whilst carbonate-free samples were analysed for TOC and isotopic carbon and nitrogen (Zhang et al., 2007).

TC, TOC and TN were measured using an organic element analyzer (Carlo-Erba model EA1110, Italy) in the State Key Laboratory of Marine Geology at Tongji University, China. TOC concentration was calibrated with equation (1) (Yang et al., 2011):

$$\text{TOC (\%)} = \text{TOC}_{\text{measured}} (\%) \times (12 - \text{TC (\%)}) / (12 - \text{TOC}_{\text{measured}} (\%)) \quad (1)$$

TOC/TN refers to the weight ratio of TOC (%) to TN (%) in this paper. Stable isotopic carbon and nitrogen were measured using a Thermo Deltaplus XL mass spectrometer (continuous flow mode) at the Third Institute of Oceanography, State Oceanic Administration, China. The stable isotopic ratios were expressed as $\delta^{13}\text{C}$ and

$\delta^{15}\text{N}$, in standard units per mil (‰), with respect to PeeDee Belemnite (PDB) and atmospheric nitrogen, respectively. The standard samples used for carbon and nitrogen isotope measuring referred to Urea#2 and Acetanilide#1 from Biogeochemistry Laboratories Indiana University. The precision for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were < 0.2 ‰ and < 0.3 ‰, respectively, based on replicated measurements ($n = 5$).

TOC/TN ratio and their isotopes are widely used to trace the source of organic matter in lakes and river-estuary-marine systems (Müller and Mathesius, 1999; Meyers, 2003; Wilson et al., 2005; Lamb et al., 2007; Leng and Lewis, 2017). Previous studies have found that TOC/TN ratios are > 12 , 4-6, and < 10 for terrestrial vegetation, bacteria, and algae respectively (in Müller and Mathesius, 1999; Lamb et al., 2007; Leng and Lewis, 2017 and references). $\delta^{13}\text{C}$ values are in a range of -21 to -31 ‰, -20 to -30 ‰, -17 to -22 ‰ for typical terrestrial C_3 plants, freshwater plankton, and marine plankton respectively (Müller and Voss, 1999 and references in; Lamb et al., 2007; Leng and Lewis, 2017 and references in). Terrestrial plants usually have relative low $\delta^{15}\text{N}$ values compared with aquatic plankton (Thornton and McManus, 1994; Middelburg and Nieuwenhuize, 1998; Müller and Voss, 1999). These values are the underlying principles on which we base our interpretation of TOC/TN, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ profiles in cores DTX4 and DTX10.

3.5 Collection of previous archive cores and palaeogeographic reconstruction of the Taihu Plain

To correlate morphological and hydrological conditions in the East Tiaoxi River Plain to other parts of the Taihu Plain and reconstruct palaeohydrological conditions of the whole Taihu Plain, previously published borehole records were compiled, selecting only those which have good chronological (radiocarbon age) control. In total, 23 sediment cores were selected and subdivided into 3 sections A-A', B-B' and C-C'

(Fig. 1A and 1C) to make stratigraphic comparisons (core details including location, dating depth, dating material, dating results and references are provided in Supplementary Table 1). Conventional ages in these cores were calibrated in software Calib 7.0 (Talma and Vogel, 1993) using the INTCAL 13 database where dating materials were plant fragments, organic rich mud, seeds, wood and pollen residues, and the Marine 13 database and ΔR value of 135 ± 42 when dating materials were marine molluscs (Yoneda et al., 2007; Reimer et al., 2013). A linear interpolation method was used to construct ages for key depths, based on the program 'Clam' with 10,000 iterations (Blaauw, 2010).

Using sedimentary, morphological and hydrological changes obtained in our new cores (DTX10 and DTX4 presented here) together with those inferred in the 23 selected existing cores, stratigraphic transections for the sections A-A', B-B' and C-C' were reconstructed. Accordingly, sedimentary morphological and hydrological conditions across the whole Taihu Plain were reconstructed for several periods between ca. 7500 and ca. 5500 cal. yr BP, associated with palaeotopography contexts during the LGM (Fig. 1C; Hong, 1991; Li et al., 2002; Wang et al., 2012) and relative sea-level changes in the Yangtze delta during the Holocene (Zong, 2004; Wang et al., 2012, 2013).

4. Results and interpretation

4.1 Lithology, AMS ^{14}C dating and sedimentary accumulation rates in cores DTX4 and DTX10

4.1.1 Core DTX4

The bottom 2.7 m of the sequence (i.e. 5.7-3.0 m depth) consisted of grey homogeneous mud with abundant plant fragments (Fig. 2). This changed to dark,

organic-rich mud over 3.0-2.0 m, followed by a peaty mud layer (2.0-1.8 m). Grey homogeneous mud with abundant plant fragments reoccurred at 1.8-1.45 m, with peaty mud at 1.75 and 1.5-1.45 m. Plant fragments declined from 1.45 m to the top. In total, six AMS ^{14}C dates were obtained over the sequence (Table 1). These ages are in reasonable sequential order with the exception of the date obtained at 1.21 m, which is older than the age obtained from the peaty mud at 1.45 m. We reject this reversed age because it could be derived from reworked plant macrofossils. By contrast, the age from peaty mud is more reliable because the peaty mud was formed by *in situ* deposition of local marsh plants. Stanley (2000) reported that it is quite common that radiocarbon dates of Holocene sediments do not become progressively older with depth in cores from the Yangtze delta, due to the introduction of old carbon during sediment transport and storage. This ‘old carbon’ phenomenon has also been discussed by Wang et al. (2012). The remaining five dates were used to generate the linear interpolation age-depth model and to calculate sedimentation rate (SR) with the program ‘Clam’ (Blaauw, 2010; Fig. 2). Relatively high SR of 0.47 cm per year (cm yr^{-1}) from 7050 to 6450 cal. yr BP (5.33-2.5 m) was followed by a reduced SR of 0.056 cm yr^{-1} from 6450-5560 cal. yr BP (2.5-1.91 m) and 0.004 cm yr^{-1} from 5560-2990 cal. yr BP (1.91-1.81 m). After 2990 cal. yr BP, SR increased to 0.017 from 1.81-1.45 m and to 0.072 from 1.45-0.8 m (Fig. 2).

4.1.2 Core DTX10

From the base of the sequence (4.7 m) to 1.0 m, lithology was grey homogeneous mud with plant fragments occurring occasionally (Fig. 2). This changed to light grey homogeneous, stiff mud with bluish leaching structures at 1.0-0.88 m, indicating subaerial pedogenesis processes and hence a sedimentation hiatus. At 0.88 m, it became dark, homogeneous mud with abundant plant fragments to the top. Four

AMS ^{14}C dates were obtained throughout the sequence (Table 1). The reversed date at 2.91 m (9475-9400 cal. yr BP) was rejected, as we argue that it was also derived from reworked plant material as discussed earlier (Stanley, 2000; Wang et al., 2012). The remaining three dates were used to construct the age-depth model and to calculate SR based on ‘Clam’ (Fig. 2). When running the age-depth model in ‘Clam’, a sedimentation hiatus was set at 1.0 m, as reflected by the pedogenesis at 1.0-0.88 m. Core DTX10 showed similar SR pattern to core DTX4. Before 7460 cal. yr BP (below 1 m), SR was high (up to 2 cm yr^{-1}), thereafter pedogenesis and a sedimentation hiatus occurred from 7460 to 2810 cal. yr BP (1.0-0.88 m). Following this, SR rose to 0.03 cm yr^{-1} after 2810 cal. yr BP (0.88-0 m).

4.2 *Diatom assemblages*

4.2.1 *Core DTX4*

A total of 236 species were identified and four habitat groups were distinguished including freshwater, salt-tolerant freshwater, brackish and marine species. No diatoms were found from 2.0-2.75 m and 1.2-1.45 m. All diatom species $> 5 \%$ were plotted as percentages of the total assemblage in Figure 3.

367

368 **Fig. 3** Diatom taxa in core DTX4 (> 5 % relative abundance) separated into freshwater (including both freshwater and salt-tolerant freshwater groups), brackish, and marine species,
 369 percentages of planktonic species, F-index and diatom concentration. Calibrated AMS ¹⁴C dates (in dark) and calculated age regions (in grey) for important boundaries are presented as
 370 ‘medial point ages ± standard deviation’, as in Figures 4, 5 and 7.

In Zone I (5.7-5.0 m, 7140-6990 cal. yr BP), diatom concentration was the highest over the whole sequence (avg. 6.84×10^3 valve g^{-1} wet weight). On the contrary, F-index was the highest, with 29.1 % of valves on average remaining in pristine state (Fig. 3). The two most abundant species were *Aulacoseira granulata* (avg. 13.8 %) and *Aulacoseira ambigua* (avg. 12.6 %). These two salt-tolerant freshwater planktonic species with low salt tolerance favour shallow fresh water (low salinity tolerant) and well-mixed hydrological conditions (Kilham, 1990; Owen and Crossley, 1991). Freshwater benthic species such as *Pseudostaurosira brevistriata* (avg. 9.2 %), *Staurosira construens* (avg. 4.9 %) and its varieties of *Staurosira construens f. venter* (avg. 4.3 %) and *Staurosira construens f. construens* (avg. 2.5 %) followed. Brackish benthic (e.g. *Diploneis smithii* var. *dilatata*, *Amphora copulata*) and freshwater benthic species (e.g. *Gomphonema gracile/parvulum*, *Eunotia minor*) were also found, but at low percentages (less than 3 % for each). The average percentage of freshwater species (including both freshwater and freshwater with low salt tolerance groups) in this section was 88.5 % while brackish species accounted for only 9.9 %, and the average percentage of planktonic species was 39.5 %.

In Zone II (5.0-4.0 m, 6990-6780 cal. yr BP), the diatom concentration decreased to 3.23×10^3 valve g^{-1} wet weight, while the F-index declined slightly to 13.9 %. Relative concentration of freshwater planktonic (e.g. *A. ambigua*) and freshwater benthic with low salt tolerance species (e.g. *P. brevistriata*, *S. construens* and its varieties of *S. construens f. venter* and *S. construens f. construens*) decreased slightly, with the exception of *A. granulata* which remained fairly constant. Correspondingly, percentages of brackish and freshwater benthic species (including *D. smithii* var. *dilatata*, *A. copulata*, *Gyrosigma acuminatum*, *G. gracile*) rose slightly. A brief recovery of the low salinity planktonic (*A. ambigua/granulata*) and benthic (*P. brevistriata*, *S. construens* and its varieties) group occurred at the top of this unit

(4.0-4.4 m). The average percentages of freshwater species dropped to 78.5 % while brackish species rose to 19.3 %, and planktonic species decreased to 29.5 %.

In Zone III (4.0-2.7 m, 6780-6500 cal. yr BP), a further drop of diatom concentration (avg. 2.23×10^3 valve g^{-1} wet weight) occurred, accompanied by increasing dissolution (F-index dropped to 7.9 %). *A. granulata* remained relatively abundant (avg. 10.8 %), despite a further decrease in the low salinity assemblages, giving way to brackish benthic (e.g., *D. smithii* var. *dilatata*, *A. copulata*, *Gyrosigma strigilis* and *Navicula menisculus*) and freshwater benthic (e.g. *G. accuminatum*, *G. gracile*, *Cymbella tumida*) species. The average percentage of freshwater species declined to 66.7 % and brackish species rose to 31.6 %, with the planktonic group falling to 20.4 %.

A remarkable shift in the diatom assemblage occurred in Zone IV (2.0 m from the top, 5570 cal. yr BP to the present), associated with a slight increase in diatom concentration (avg. 2.85×10^3 valve g^{-1} wet weight) and increase in F-index (avg. 9.7 %). This unit was dominated by freshwater benthic and epiphytic species of *G. parvulum/gracile*, *E. minor*, *Fragilaria vaucheriae* and *Cymbella silesiaca*, with very few brackish benthic (e.g. *D. smithii* var. *dilatata*, *A. copulata*, *N. menisculus*, *G. strigilis*) and freshwater to low salinity tolerant planktonic species (e.g. *A. granulata/ambigua*). In the uppermost samples (1.1-1.0 m), *Diadesmis confervacea* became frequent (up to 45 %), an indicator of aerial/terrestrial or very shallow-water conditions (Gell et al., 2007). Freshwater species accounted for 95.1 % and brackish species were < 5 % on average, while planktonic species accounted for only 3.9 % of the assemblage.

420 4.4.2 Core DTX10

421 Diatom concentration was low with an average value of 0.46×10^3 valve g^{-1} wet
422 weight for the whole core (Fig. 4), probably resulting from strong dissolution of
423 diatoms reflected by low average F-index value ($\sim 7.4\%$) and dilution caused by high
424 sediment accumulation rates or low diatom productivity. No diatoms were found from
425 2.7-3.0 m and 0.84-2.0 m and only a few broken diatom fragments were found on
426 slides for samples from the top to 0.84 m. Preservation in one sample at 0.66 m was
427 slightly better, but most remaining diatoms were the dissolved centres of large benthic
428 genera *Pinnularia* or *Cymbella*, but which cannot be identified to species level. For
429 other samples from 0.3 to 0.84 m, broken fragments of freshwater genera, such as
430 *Eunotia* species, were seen. A total of 154 species were identified for the whole
431 sequence and diatom species $> 5\%$ plotted as relative abundance in Figure 4.

432 The diatom assemblage from the base to 2.0 m (7650-7520 cal. yr BP; except
433 2.7-3.0 m where no diatoms were found) was dominated by *Cyclotella striata* (avg.
434 22.5 %), which is usually abundant in river-mouth/estuarine environments (Ryu et al.,
435 2005), *Actinoptychus senarius* (avg. 19.8 %), the relatively low-salinity coastal
436 planktonic species (Grönlund, 1993; Hasle and Syvertsen, 1996), *Paralia sulcata* (avg.
437 9.3 %) which is used as a marker for the coastal East China Sea (Tada et al., 1999.
438 Ryu et al., 2005), and *Chaetoceros* spp. resting spores, typical marine species (avg.
439 9.2 %). Other species found but in relatively low percentage were coastal species of
440 *Thalassiosira nanolineata* (Grönlund, 1993; Hasle and Syvertsen, 1996) and brackish
441 benthic species such as *Diploneis smithii* var. *dilatata/smithii*, *Nitzschia granulata*,
442 *Cocconeis costata* var. *pacifica*. In total, percentages of marine and brackish species
443 were 56.3 % and 37.9 % on average respectively, and planktonic species 77.4 %
444 (including tychoplanktonic taxa).

Fig. 4 Diatom taxa in core DTX10 (> 5 % relative abundance) with separated into freshwater, brackish and marine species, percentages of planktonic species, F-index and diatom concentration.

Although preservation was very poor in samples from the top to 0.84 m, it was clear that the original diatom assemblage was completely different. Almost 99% of diatom valves (as represented by the sample at 0.66 m) belonged to freshwater benthic species (including *Gomphonema gracile*, *Cymbella aspera*, *Eunotia minor*, *E. formica* and unidentified *Pinnularia* and *Cymbella* species). We argue that the switch from a brackish/marine to a freshwater assemblage from Zone I to Zone II reflects a real change to the salinity and hydrology of the system as the vestigial fragments of diatoms in Zone II are resistant and have distinctive morphologies. Had these taxa

been part of the assemblage of earlier sections, they would have been significant aspects of the assemblages there also. By the same token, most of the taxa in the earlier, more marine parts of Zone I are also very resistant to dissolution (e.g. *Chaetoceros* cysts, *Paralia sulcata*, *Cyclotella striata*) with distinctive taphonomic end-members, and would have appeared in Zone I had they been present in the original assemblage in that section. Thus we are confident that Zone I was deposited under brackish-marine conditions while Zone II represents a freshwater system.

4.3 Grain size, TOC/TN ratio and C-N stable isotopes

Based on grain size composition, TOC, TN, TOC/TN and C-N stable isotope ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) variations, cores DTX4 and DTX10 were divided into five and three zones, denoted as Zone I-V, respectively (Figs 5-7). Zone II and V in core DTX4 (Fig. 5) were both divided into three sub-zones based on variation of TOC/TN ratio, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values. Zone I and III in core DTX10 were also divided into three and two sub-zones respectively.

Fig. 5 Lithology and sedimentary parameters of core DTX4, including AMS ^{14}C age, grain size distribution, TOC, TN, TOC/TN, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and interpretation of sedimentary environment.

4.3.1 Core DTX4

In Zone I (5.7-5.5 m, 7140-7100 cal. yr BP), clay content ($< 4 \mu\text{m}$) was around 24.8 %, silt (4-63 μm) 67.7 %, and sand (63-2000 μm) 7.3 % on average. In Zone II (5.5-2.5 m, 7100-6450 cal. yr BP), clay content (avg. 19.6 %) decreased and silt content (avg. 72.7 %) increased slightly, particularly for grains between 16 and 63 μm . No change occurred in sand content. Particles between 8-16 μm increased slightly while particles between 16-63 μm fell in II_3 (4.0-2.5 m, 6780-6450 cal. yr BP) compared to II_1 (5.5-5.0 m, 7100-6990 cal. yr BP) and II_2 (5.0-4.0 m, 6990-6780 cal.

yr BP). Grain size assemblages in Zone III (2.5-2.0 m, 6450-5570 cal. yr BP) remained similar to those of Zone II, followed by a sudden increase in sand content in Zone IV (2.0-1.8 m, 5570-2990 cal. yr BP) and clay content in Zone V (1.8 m to the top, 2990 cal. yr BP to the present) with an opposite trend in silt content.

The linear correlation of TOC to TN (Fig. 6A) implied that TN content was controlled mostly by organic matter, while the positive TN intercept of 0.07 % indicated a slight contribution from inorganic nitrogen. Plots of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ against TOC/TN revealed a mixture of freshwater and marine algae and terrestrial C_3 plants (Fig. 6B and 6C). In Zone I, TOC (avg. 0.47 %), TN (avg. 0.06 %) and TOC/TN (avg. 7.48) minima coincided with relatively high $\delta^{13}\text{C}$ (avg. -24.04 ‰) and $\delta^{15}\text{N}$ (avg. $+3.16$ ‰), indicating contribution from marine plankton and bacteria (Müller and Mathesius, 1999; Müller and Voss, 1999; Lamb et al., 2007). The bi-plot of $\delta^{13}\text{C}$ to TOC/TN showed that Zone I samples lay in the domain of marine algae or marine particulate organic carbon (POC; Fig. 6B). The bi-plot of $\delta^{15}\text{N}$ against TOC/TN reflected a predominant contribution from algae as well (Fig. 6C).

An abrupt increase in TOC (1.07-2.12 %, avg. 1.47 %), TN (0.10-0.16 %, avg. 0.13 %) and TOC/TN (9.07-13.50, avg. 11.53) characterised zone II with lower $\delta^{13}\text{C}$ (from -25.58 ‰ to -26.61 ‰, avg. -26.74 ‰) and $\delta^{15}\text{N}$ (from $+1$ ‰ to $+4.39$ ‰, avg. $+2.25$ ‰), signifying the sudden increase in importance of C_3 terrestrial plants and reflecting the formation of a marsh environment. A shift on the bi-plots of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ vs. TOC/TN implied a mixture of freshwater POC and C_3 plants for samples in Zone II with one exception of marine POC in Zone II₂. The gradual increase in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ from Zone II₁ to II₂, and to II₃, coincided with changes in position in bi-plots of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ against TOC/TN which was interpreted as a slight, but persistent, increase in the importance of marine algae/POC from the bottom to the top of Zone II.

Fig. 6 Correlation between TOC and TN (A and D), biplot of $\delta^{13}C$ values and TOC/TN (B and E), correlation between $\delta^{15}N$ values and TOC/TN (C and F) for core DTX4 (A to C) and DTX10 (D to F), respectively. Regions for different type of organic matter are after [Lamb et al. \(2007\)](#).

TOC, TN, TOC/TN values increased gradually to maxima of 10.83 %, 0.51 % and 21.09, respectively, from Zone III to the top of IV where lithology changed to peaty mud, while $\delta^{13}C$ and $\delta^{15}N$ values dropped consistently to minima of -27.76 ‰ and -0.55 ‰, respectively. Integrating patterns on the two bi-plots ([Fig. 6B and 6C](#)),

Zone III was dominated mostly by terrestrial C₃ plants and secondarily by freshwater algae/POC, while typical terrestrial C₃ plants dominated the organic matter composition of Zone IV. In Zone V, a rapid drop in TOC (avg. 1.34 %), TN (avg. 0.14 %) and the TOC/TN ratio (avg. 9.13) occurred, with lowest values in subzone V₂ followed by an increase in subzone V₃. The $\delta^{13}\text{C}$ values remained depleted except for subzone V₂ (1.25-1.45 m) where they showed peaks, while the $\delta^{15}\text{N}$ values increased suddenly and remained high from the bottom to the top. The pattern on the two bi-plots (Fig. 6B and 6C) suggested an increase in freshwater algae/POC and bacteria in Zone V, especially in Zone V₂ where algae and bacteria dominated.

4.3.2 Core DTX10

In Zone I (from the base to 1.0 m, 7650-7460 cal. yr BP), silt content was 50-74 %, the highest values in the profile, with lesser amounts of clay and sand (Fig. 7). Zone II (1.0-0.88 m, 7460-2810 cal. yr BP) was a transitional zone for all proxies. There was a slight and gradual increase in silt content from the bottom to the top, while sand content remained constant. In Zone III (0.88 m to the top, 2810 cal. yr BP to present) both silt and clay content dropped while sand content rose dramatically to 26 % on average, with a slight increase in clay and silt content at the top of Zone III₂.

Fig. 7. Lithology and sedimentary parameters of core DTX10, including AMS ^{14}C age, grain size distribution, TOC, TN, TOC/TN, $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and sedimentary environments interpretation.

A linear correlation between TOC and TN was found, but two discrete groups were identified (Fig. 6D): the samples in Zone I formed one group and samples in Zone II and III a second. This showed that TN values were mostly controlled by organic matter, but also inorganic nitrogen contributed slightly as shown by the positive TN intercept of 0.035 % and 0.09 % respectively. Plotting $\delta^{13}\text{C}$ against TOC/TN revealed a mixture of two end members of POC/algae/bacteria and C_3 terrestrial plants (Fig. 6E). This explanation was supported by the strongly negative linear correlation between $\delta^{15}\text{N}$ and TOC/TN ($r = -0.85$; Fig. 6F). TOC values were 0.42-0.89 % (avg. 0.59 %) in zone I and TN 0.08-0.14 % (avg. 0.10 %) (Fig. 7), while TOC/TN ratios were 4.35-7.41 (avg. 5.80). These minimum values were accompanied

by the highest $\delta^{13}\text{C}$ (from -23.82 ‰ to -26.32 ‰ , avg. -24.73 ‰) and $\delta^{15}\text{N}$ values (from $+3.44\text{ ‰}$ to $+6.12\text{ ‰}$, avg. 4.30 ‰). Very small increases in TOC, TN and TOC/TN coincided with minor decreases of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ from Zone I₁ to I₃. Integrating patterns on the plots of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ against TOC/TN, a mixture of freshwater and marine algae/POC and bacteria composed the organic matter source for samples in Zone I, with the freshwater algal/POC contribution increasing from Zone I₁ to I₃. In Zone II, TOC and TN values increased rapidly up to 1.79 ‰ and 0.14 ‰ on average and TOC/TN up to 12.37. On the contrary, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ declined to -27.89 ‰ and $+2.39\text{ ‰}$, respectively. Changes in all these proxies reflected an abrupt addition of terrestrial organic matter from C₃ plants, supported by the bi-plots of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ against TOC/TN. In Zone III, this pattern of consistently high TOC, TN and TOC/TN and lower $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ continued, with small decreases in these parameters in Zone III₂. On bi-plots of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ vs. TOC/TN, samples of Zone III₁ fell in the region typical of C₃ terrestrial plants and samples of Zone III₂ in the region between C₃ plants and freshwater POC/algae.

5. Discussion

5.1 Sedimentary, morphological and hydrological changes in the East Tiaoxi River Plain

The dominance of marine diatoms and marine algae/POC contribution for the TOC in core DTX10 confirms that there was a high salinity estuary (the Palaeo-Taihu Estuary) in the East Tiaoxi River Plain before ca. 7500 cal. yr BP (Figs 4, 7; Hong, 1991), in response to rapid early to mid-Holocene sea-level rise (Chappell and Polach, 1991; Bard et al., 1996; Bird et al., 2007; Wang et al., 2012). The high sedimentation accumulation rate up to 2 cm yr^{-1} in core DTX10 during this period signified rapid

infilling of this estuary. From ca. 7500 cal. yr BP, freshening occurred at site DTX10 indicated by the increase in organic source from freshwater algae/POC (Zone I₂-I₃ in Fig. 7). At some time after ca. 7500 cal. yr BP, the area around core DTX10 was subaerially exposed until ca. 2800 cal. yr BP, inferred from the sedimentation hiatus and observed pedogenesis (1.0-0.88 m in Fig. 2 and 7). In contrast, the area around core DTX4 changed from estuary to a low-salinity marsh during ca. 7100-7000 cal. yr BP, indicated by dominant organic matter changing from marine algae/POC to freshwater algae/POC and C₃ terrestrial plants (Fig. 5). The estuary at 7100 cal. yr BP was also characterised by low salinity, with the diatom assemblages dominated by salt-tolerant freshwater group of *A. granulata/ambigua*, *P. brevistriata*, *S. construens* and its varieties of *S. construens f. venter* and *S. construens f. construens* (Fig. 3). These diatom taxa are often found in isolation basins (e.g. freshwater lakes formed as sea level falls) (Stabell, 1985), and therefore, their high concentrations indicate that the site DTX4 was in the stage of isolation from seawater during ca. 7100-7000 cal. yr BP. Consequently, we infer that the freshening of the East Tiaoxi River Plain likely started between 7500-7100 cal. yr BP, possibly due to coastal development and freshwater discharge from the Yangtze River, benefitting from a warmer and, more importantly, wetter climate during the Holocene Thermal Maximum (Wang et al., 2005). Particularly, *A. granulata* is often found in rivers/lakes in the flood and delta plains in eastern China and has been considered as an indicator of freshwater discharge to estuaries (Chen et al., 2011; Dong et al., 2008; Liu et al., 2012; Wang et al., 2009). We thus suppose high and constant percentages of *A. granulata* during 7100-6500 cal. yr BP in core DTX4 implied the strong freshwater influence of Yangtze River runoff.

At some time after ca. 7000 cal. yr BP, although the marsh was dominated by C₃ plants and strongly influenced by the Yangtze freshwater discharge, penetration of salt

water occurred at core DTX4 as percentages of brackish benthic diatom and influence from marine POC/algae increased (Zone II₂ and II₃ in Fig. 5; Fig. 6). During this period, the infilling of the Palaeo-Taihu Estuary continued, but at a lower SR of 0.48 cm yr⁻¹. The return of sea water was likely caused by the sudden sea-level rise at ca. 6.8 ± 0.2 ka BP due to the disappearance of the west section of the Laurentide ice sheet (Blanchon and Shaw, 1995; Carlson et al., 2007). From ca. 6500 to 5600 cal. yr BP, sea water retreated slowly again and the sedimentary environment transitioned gradually from salt marsh to freshwater marsh in the area near core DTX4, reflected by the composition of organic matter over this period (Fig. 5), while site DTX10 was subaerially exposed (1.0-0.88 m in Fig. 7). Correspondingly, the infilling rate of the Palaeo-Taihu Estuary fell dramatically, likely signifying a process of shrinking due to the stable or slightly declining sea level from ~6300 cal. yr BP (Fairbanks, 1989; Bard et al., 1996; Bird et al., 2007; Wang et al., accepted). After ca. 5600 cal. yr BP, the East Tiaoxi River Plain transitioned through a range of environments from stable freshwater marsh (core DTX4) or dry land (core DTX10), characterised by freshwater benthic diatoms and C₃ terrestrial plants respectively (Fig. 3-7). In other words, no sea water penetration occurred and entirely freshwater environments persisted from ca. 5600 cal. yr BP throughout to the present. In terms of the Palaeo-Taihu Estuary, it likely shrank dramatically as it was filled up with sediment.

This study also demonstrates that viable and informative diatom counts can be made using the sodium polytungstate (SPT) density separation technique even on material which has very low diatom concentrations (here, largely as a result of dissolution). There is some indication that there is some preferential recovery of certain taxa (compare adjacent samples using the standard water-bath and SPT method in core DTX10 in Figure 4, for example for *Diploneis smithii* in Zone I) but there is clearly little systematic difference between the two methods, thus confirming

the utility of the approach in such sediments. Results here support the use of SPT as a very useful technique which may have wider application in estuarine and marine sediments, where the option of working on material with well-preserved or abundant diatoms (or other siliceous remains) may simply not be available, and yet may deliver much information of great palaeoecological value, even with poorly preserved assemblages (cf. [Ryves et al. 2006, 2009](#)).

5.2 Environmental changes in the whole Taihu Plain

We recognise six stages for the morphological and hydrologic evolution of the Palaeo-Taihu Estuary according to the multi-proxy analyses in cores DTX4 and DTX10. Before 7500 cal. yr BP, it was a high salinity estuary; 7500-7100 cal. yr BP, a low salinity estuary influenced strongly by the Yangtze freshwater discharge; 7100-7000 cal. yr BP, it developed into a low salinity to freshwater marsh; 7000-6500 cal. yr BP, salt water penetrated into the marsh; 6500-5600 cal. yr BP, gradual retreat of marine influence; and from 5600 cal. yr BP, stable freshwater conditions. A similar history of hydrologic change can be inferred from the stratigraphic transections across the Taihu Plain ([Fig. 8](#)).

Fig. 8. Stratigraphic transections of collected cores, including section A-A' (A) and B-B' (B) and C-C' (C). In section A-A', core 97C is after Ding (2004), core 97A after Zhou and Zheng (2000), and core DS, JLQ, GT and SL are after Yan and Huang (1987) and Hong (1991), and core XL is after

Liu et al. (2015). In section B-B', site LTD is after Li et al. (2008), core W1 after Chang et al. (1994) and Wang et al. (2001), core E2 after Wang et al. (2001), core YJD and PW after Zong et al. (2011, 2012b) and Innes et al. (2014), core WJB after Qin et al. (2011) and MJB after Long et al. (2014). In section C-C', core CXS, CD and TYL are after Zong et al. (2012b), core SQ, TCM and TL after Zong et al. (2011), core QP after Atahan et al. (2008) and Itzstein-Davey et al. (2007), ZX-1 after Chen et al. (2005) and Tao et al. (2006) and core GEL after Wang et al. (2012). 0 m is the start depth for each core. Ages in black are from references for each core and ages in grey black are linearly interpolated based on "Clam" (Blaauw, 2010).

Estuarine facies dominated before 7500 cal. yr BP in multiple cores including 97C, 97A, DS, JLQ, GT and XL in the palaeo-Taihu Estuary (Fig. 8A; references for these cores are in Supplementary Table 1), due to rapid sea-level rise in the early Holocene (Wang et al., 2012). The sedimentary facies then turned into tidal flat in these cores around 7500 cal. yr BP owing to the infilling of the estuary (Fig. 8A). Brackish tidal flat conditions also occurred in other cores like ZX-1, GFL and TL in the east Taihu Plain (Fig. 8C). As the relative sea level reached approximately -6 m at 7500 cal. yr BP (Wang et al., 2012), the late Pleistocene interfluvial terrace T1 region (including around Shanghai City) was a shallow sea environment due to this transgressive phase (Li et al., 2001; Zong et al., 2004; Wang et al., 2012, 2013). No sea water penetration occurred throughout the Holocene in the region of the late Pleistocene interfluvial terrace T3 (Fig. 1C) where palaeo-altitude was the highest during the LGM (as recorded in core E2, YJD, CD, CXS and SQ; Fig. 8B and 8C). The central Taihu Plain was isolated from western uplands due to sea water inundation through the Palaeo-Taihu Estuary at ca. 7500 cal. yr BP (Fig. 9A).

Fig. 9. Palaeogeographic map for the Taihu Plain before ca. 7500 cal. yr BP (A), during ca. 7500-7000 cal. yr BP (B), during ca. 7000-6500 cal. yr BP (C), and at ca. 5500 cal. yr BP (D). Rivers in (D) are based on [Hong \(1991\)](#), and the possible formation of Taihu Lake is based on [Wang et al. \(2001\)](#). The number of Liangzhu (5500-4500 cal. yr BP) cultural sites grew rapidly to 461, while it is only 78 and 93 during the Majiabang (7000-5800 cal. yr BP) and Songze (5800-5500 cal. yr BP) period, respectively (from [Zheng, 2002](#); [Chen, 2002](#); [Xu, 2015](#)).

The marine regression at ca. 7500-7000 cal. yr BP recorded in core DTX4 and DTX10 was also seen in cores in the Palaeo-Taihu Estuary (e.g. cores JLQ, GT and WJB; Fig. 8A). In the central Taihu Plain, low salinity to freshwater marshes developed at ca. 7600-7120 cal. yr BP in core WJB (Fig. 8B), indicated by reduced concentration of marine dinoflagellate cysts and *Chenopodiaceae* pollen (Qin et al., 2011). Consequently, this sea water regression event promoted the rapid expansion of low salinity to freshwater marshes, especially in the area of the Palaeo-Taihu valley (Fig. 9B).

Evidence for sea water penetration after 7000 cal. yr BP was found in several sediment cores (e.g. PW, WJB, MJB) in the central Taihu Plain (Fig. 8B) in addition to those in the Palaeo-Taihu Estuary (Fig. 8A). Moreover, this marine transgression reached areas where no sea water influence was observed in the records before. For example, the foraminifera *Ammonia compressiuscula* and *Ammonia* cff. *sobrina*, which prefer brackish tidal flat environments, occurred for the first time in the peat layer dated at ca. 7000 cal. yr BP at site LTD, located at the head of the Palaeo-Taihu valley, northwest of the Taihu Plain (Li et al., 2008; Fig. 8B). Marine and brackish diatom species, marine species of dinoflagellate cysts and foraminifera were only found at 50-90 cm (dated at ca. 7000 cal. yr BP) in core W1 in the north part of Taihu Lake (Fig. 8B and 9C; Chang et al., 1994, Wang et al., 2001). Sea water also inundated the low salinity marsh around core PW in the south east Taihu Plain at some time after 7200 cal. yr BP (Fig. 8B; Zong et al., 2011; Innes et al., 2014). Therefore, the transgression between ca. 7000-6500 cal. yr BP influenced a large area, including the innermost Taihu Plain, through low-lying area such as the Palaeo-Taihu valley (Fig. 9C). Low salinity marsh then returned from ca. 6200 to 5600 cal. yr BP at sites LTD, W1, PW and DTX4 (Fig. 8), and thereafter stable freshwater conditions in

the central Taihu Plain developed completely after ca. 5600 cal. yr BP, likely indicating the closure of the Palaeo-Taihu Estuary (Fig. 9D).

5.3 Role of hydrological environments on the development of rice farming

Rice growth is susceptible to salinity conditions and demands suitable water depth, in addition to warm and humid climate (Zeng et al., 2003; Yu et al., 2000; Chen et al., 2005; Innes et al., 2009, 2014; Patalano et al., 2015). The Taihu Plain was semi-encircled by sea water before ca. 7500 cal. yr BP (Fig. 9A), corresponding with no sedentary Neolithic settlements and no remains of rice cultivation (Mo et al., 2011). Concurrent with withdrawal of sea water and expansion of low salinity and freshwater marshes between ca. 7500 cal. yr BP and ca. 7000 cal. yr BP, rice cultivation began (Cao et al., 2006; Fuller et al., 2007; Mo et al., 2011; Zong et al., 2011) and the Majiabang culture developed in the Taihu Plain, in addition to benefits of heat and precipitation provided by the optimum climate (Chen et al., 2005; Wang et al., 2005). However, no rapid advance in rice cultivation or productivity, or in the number of Neolithic sites, occurred during the late Majiabang and early Songze period. This is likely due to the lack of adequate freshwater supply, because the central Taihu plain was isolated from the western uplands by the Taihu-Palaeo Estuary until at around 5600 cal. yr BP (Fig. 9), and hence no river discharge from the western uplands entered the central Taihu plain.

We further suggest that the freshening of the East Tiaoxi River Plain or the shrinkage/closure of the Palaeo-Taihu Estuary at ca. 5600 cal. yr BP, was critical for the rapid expansion of rice agriculture across the whole Taihu Plain during the Liangzhu period (Fig. 9D), in the context of deteriorating climate conditions compared with the Majiabang and Songze period (Wang et al., 2005; Innes et al., 2014). Firstly, the shrinkage/closure of the Palaeo-Taihu Estuary, which is supported

by extensive distribution of Liangzhu cultural sites in the Tiaoxi River Plain (Zheng, 2002), prevented the intrusion of sea water from Hangzhou Bay into the Taihu Plain, allowing freshwater conditions, particularly in the southern and western parts. Secondly, it restricted the discharge of freshwater from the western uplands (e.g. the west Tiaoxi river) into the Hangzhou Bay, and instead, forced this freshwater to flow eastwards, likely forming three previously existing rivers: the Loujiang River, the Wusongjiang River and the Dongjiang River (Fig. 9D; Hong, 1991). This increasing density of the river network would provide an increasing area of freshwater wetlands available to be transformed into rice paddies and greater quantity of fresh water for the east Taihu Plain. This conjecture is supported by the fact that the area of rice paddy fields excavated in several archeological sites recently was several times larger during the Liangzhu period than during the Majiabang and Songze period, and were connected to natural and artificial creeks, instead of wells and ponds, which had stronger capability of water storage (Cao et al., 2007; Hu et al., 2013; Zheng et al., 2014; Zhuang et al., 2014). Thirdly, new or enlarged freshwater marsh environments of the East Tiaoxi River Plain would have supplied additional freshwater wetland resources. Lastly, water level rise, in response to sea-level rise after the Taihu Plain was separated from the Hangzhou Bay and the East China Sea, would also encourage formation of freshwater wetlands (Hong, 1991; Chen et al., 1997). All these advantages together would have increased opportunity for domesticated rice cultivation in the whole Taihu Plain and rice productivity, supporting the continuous and dramatic advancement of the Liangzhu culture.

We also speculate that the terrestrialisation and freshening of the East Tiaoxi River Plain may have facilitated communication between the north and east Taihu Plain, with the capital city of Liangzhu at Mojiaoshan, southwest of Taihu Plain (Fig. 9D) and even with areas south of Hangzhou Bay (Chen, 2015; Xu, 2015), given the

barrier that the Palaeo-Taihu Estuary would have represented. Such easier communication would support the development and flourishing of the Liangzhu capital city. In return, the capital city would be act as a hub for advanced technology and cultural innovation, promoting the further expansion of rice farming and cultural development in the region over the Liangzhu period.

6. Conclusions

A multiproxy sedimentological analysis combining chronological, lithological, geochemical and biological analyses of two sediment cores (DTX4 and DTX10) collected from the East Tiaoxi River Plain, southern Yangtze delta Plain, have shed light on changes in landscape and hydrology in this region over the last 7500 years. Before ca. 7500 cal. yr BP, the Palaeo-Taihu Estuary existed along the present day East Tiaoxi River Plain. It infilled rapidly by ca. 7500 cal. yr BP and was characterised by low salinity conditions, because of the large supply of freshwater from the Yangtze. Sea water, however, again penetrated the East Tiaoxi River Plain after ca. 7000 cal. yr BP due to an abrupt sea-level rise, and a salt marsh environment developed. This transgression was also recorded in other parts of the Taihu Plain where no sea water influence occurred before, hence, it was possibly the largest sea-level transgression during the Holocene, based on the synthesis of a large number of hydrological and environmental records from the region. After ca. 6500 cal. yr BP, sea-water penetration gradually declined and infilling rate of the Palaeo-Taihu Estuary fell owing to slowing sea-level rise. The dramatic shrinkage/closure of the Palaeo-Taihu Estuary occurred after ca. 5600 cal. yr BP, corresponding to the formation of stable freshwater marsh (or subaerial land) conditions in the East Tiaoxi River Plain.

Geomorphological and hydrological changes in the East Tiaoxi River Plain played an important role in the rise of rice cultivation and development of Neolithic cultures across the Taihu Plain. Especially, the freshening of the East Tiaoxi River Plain or its terrestrialisation after ca. 5600 cal. yr BP, was likely one critical precondition encouraging rapid increase of rice productivity (and so cultural development) during the Liangzhu period.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (Grant No. 41576042) and the China Scholarship Council Postgraduate Scholarship Program, which allowed TC spending a year at Loughborough University. We are grateful to two anonymous reviewers for their helpful comments.

References

Atahan, P., Itzstein-Davey, F., Taylor, D., Dodson, J., Qin, J., Zheng, H., Brooks, A., 2008. [Holocene-aged sedimentary records of environmental changes and early agriculture in the lower Yangtze, China. Quaternary Science Reviews 27, 556-570.](#)

Bard, E., Hamelin, B., Arnold, M., Montaggioni, L., Cabioch, G., Faure, G., Rougerie, F., 1996. [Deglacial sea-level record from Tahiti corals and the timing of global meltwater discharge. Nature 382, 241-244.](#)

Battarbee, R.W., Jones, V.J., Flower, R.J., Cameron, N.G., Bennion, H., Carvalho, L., Juggins, S., 2001. [Diatoms, in: Smol, J.P., Birks, H.J.B., Last, W.M.,](#)

802 Bradley, R.S., Alverson, K. (Eds.), Tracking Environmental Change Using Lake
 803 Sediments: Terrestrial, Algal, and Siliceous Indicators. Springer Netherlands,
 804 Dordrecht, pp. 155-202.

805 Battarbee, R.W., Kneen, M.J., 1982. The use of electronically counted
 806 microsphere sin absolute diatom analysis. *Limnology & Oceanography* 27, 184-188.

807 Beuselinck, L., Govers, G., Poesen, J., Degraer, G., Froyen, L., 1998. Grain-size
 808 analysis by laser diffractometry: comparison with the sieve-pipette method. *Catena* 32,
 809 193-208.

810 Bird, M., Fifield, L., Teh, T., Chang, C., Shirlaw, N., Lambeck, K., 2007. An
 811 inflection in the rate of early mid-Holocene eustatic sea-level rise: A new sea-level
 812 curve from Singapore. *Estuarine, Coastal and Shelf Science* 71, 523-536.

813 Blaauw, M., 2010. Methods and code for ‘classical’ age-modelling of
 814 radiocarbon sequences. *Quaternary Geochronology* 5, 512-518.

815 Blanchon, P., Shaw, J., 1995. Reef drowning during the last deglaciation:
 816 Evidence for catastrophic sea-level rise and ice-sheet collapse. *Geology* 23, 4-8.

817 Cao, Z.H., Ding, J.L., Hu, Z.Y., Knicker, H., Kögel-Knabner, I., Yang, L.Z., Yin,
 818 R., Lin, X.G., Dong, Y.H., 2006. Ancient paddy soils from the Neolithic age in
 819 China’s Yangtze River Delta. *Naturwissenschaften* 93, 232-236.

820 Cao, Z.H., Yang, L.Z., Lin, X.G., Hu, Z.Y., Dong, Y. H., Zhang, G.Y., Lu, Y.C.,
 821 Yin, R., Wu, Y.L., Ding, J.L., Zheng, Y.F., 2007. Morphological characteristics of
 822 paddy fields, paddy soil profile, phytolith and fossil rice grain of the Neolithic age in
 823 Yangtze river delta. *Acta Pedologica Sinica* 44, 838-847 (In Chinese, with English
 824 abstract).

825 Carlson, A.E., Clark, P.U., Raisbeck, G.M., Brook, E.J., 2007. Rapid Holocene
 826 Deglaciation of the Labrador Sector of the Laurentide Ice Sheet. *Journal of Climate*
 827 20, 5126-5133.

828 Chang, W.Y.B., Xu, X.M., Yang, J.R., Liu, J.L., 1994. Evolution in Taihu Lake
 829 ecosystem as evidence of changes in sediment profiles. *Journal of Lake Science* 6,
 830 217-226.

831 Chappell, J., Polach, H., 1991. Post-glacial sea-level rise from a coral record at
 832 Huon Peninsula, Papua New Guinea. *Nature* 349, 147-149.

833 Chen, J., 2002. Neolithic cultures in the Yangtze delta, China and their
 834 environments. PhD Thesis, East China Normal University, China (in Chinese, with
 835 English abstract).

836 Chen, J., 2015. The Formation of the Songze Culture. *Relics from South* 1, 57-65
 837 (in Chinese, with English abstract).

838 Chen, X., Yang, X., Dong, X., Liu, Q., 2011. Nutrient dynamics linked to
 839 hydrological condition and anthropogenic nutrient loading in Chaohu Lake (southeast
 840 China). *Hydrobiologia* 661, 223-234.

841 Chen, Z., Hong, X., Li, S., Wang, L., Shi, X., 1997. Study of archaeology-related
 842 environment evolution of Taihu Lake in southern Chang Jiang Delta Plain. *Acta*
 843 *Geographica Sinica* 52, 131-137 (in Chinese, with English abstract).

844 Chen, Z., Wang, Z., Schneiderman, J., Taol, J., Cail, Y., 2005. Holocene climate
 845 fluctuations in the Yangtze delta of eastern China and the Neolithic response. *The*
 846 *Holocene* 15, 915-924.

847 Chen, Z.Y., Zong, Y.Q., Wang, Z.H., Wang, H., Chen, J., 2008. Migration
848 patterns of Neolithic settlements on the abandoned Yellow and Yangtze River deltas
849 of China. *Quaternary Research* 70, 301-314.

850 Ding, Y. F., 2004. Deposit record of climate and environmental changes of Taihu
851 Lake since 10,000 a. Master Thesis, East China Normal University, China (in Chinese,
852 with English abstract).

853 Dong, X., Bennion, H., Battarbee, R., Yang, X., Yang, H., Liu, E., 2008.
854 Tracking eutrophication in Taihu Lake using the diatom record: potential and
855 problems. *Journal of Paleolimnology* 40, 413-429.

856 Fairbanks, R.G., 1989. A 17, 000-year glacio-eustatic sea level record: influence
857 of glacial melting rates on the Younger Dryas event and deep-ocean circulation.
858 *Nature* 342, 637-642.

859 Fan, Y. P., 2011. On the development of rice farming in Tailake area during the
860 Neolithic age of China. Master Thesis, Nanjing Agriculture University, China (in
861 Chinese, with English abstract).

862 Fuller, D.Q., Harvey, E., LING, Q., 2007. Presumed domestication? Evidence
863 for wild rice cultivation and domestication in the fifth millennium BC of the Lower
864 Yangtze region. *Antiquity* 81, 316-331.

865 Gell, P., Tibby, J., Little, F., Baldwin, D., Hancock, G., 2007. The impact of
866 regulation and salinisation on floodplain lakes: The lower River Murray, Australia.
867 *Hydrobiologia* 591, 135-146.

868 Grönlund, T., 1993. Diatoms in surface sediments of the Gotland Basin in the
869 Baltic Sea. *Hydrobiologia* 269-270, 235-242.

- 870 Hasle, G.R., Syvertsen, E.E., 1996. Chapter 2-Marine Diatoms. Identifying
871 Marine Diatoms & Dinoflagellates, 5-385.
- 872 Hong, X., 1991. Origin and evolution of the Taihu Lake. Marine Geology and
873 Quaternary Geology 11, 87-99 (in Chinese, with English abstract).
- 874 Hu, L., Chao, Z., Gu, M., Li, F., Chen, L., Liu, B., Li, X., Huang, Z., Li, Y.,
875 Xing, B., 2013. Evidence for a Neolithic Age fire-irrigation paddy cultivation system
876 in the lower Yangtze River Delta, China. Journal of Archaeological Science 40,
877 72-78.
- 878 Innes, J.B., Zong, Y., Chen, Z., Chen, C., Wang, Z., Wang, H., 2009.
879 Environmental history, palaeoecology and human activity at the early Neolithic
880 forager/cultivator site at Kuahuqiao, Hangzhou, eastern China. Quaternary Science
881 Reviews 28, 2277-2294.
- 882 Innes, J.B., Zong, Y., Wang, Z., Chen, Z., 2014. Climatic and palaeoecological
883 changes during the mid-to Late Holocene transition in eastern China: high-resolution
884 pollen and non-pollen palynomorph analysis at Pingwang, Yangtze coastal lowlands.
885 Quaternary Science Reviews 99, 164-175.
- 886 Itzstein-Davey, F., Atahan, P., Dodson, J., Taylor, D., Zheng, H., 2007.
887 Environmental and cultural changes during the terminal Neolithic: Qingpu, Yangtze
888 delta, eastern China. The Holocene 17, 875-887.
- 889 Jiang, L., Liu, L., 2006. New evidence for the origins of sedentism and rice
890 domestication in the Lower Yangzi River, China. Antiquity 80, 355-361.
- 891 Juggins, S., 1991-2009. C2 Data Analysis. Newcastle University, Newcastle.
892 Available at: <http://www.staff.ncl.ac.uk/staff/stephen.juggins/software/C2Home.htm>.

- 893 Juggins, S., 2013. Quantitative reconstructions in palaeolimnology: new
894 paradigm or sick science? *Quaternary Science Reviews* 64, 20-32.
- 895 Kilham, P., 1990. Ecology of *Melosira* Species in the Great Lakes of Africa, in:
896 Tilzer, M.M., Serruya, C. (Eds.), *Large Lakes: Ecological Structure and Function*.
897 Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 414-427.
- 898 Krammer, K., Lange-Bertalot, H., 1986. Bacillariophyceae. 1: Teil: Naviculaceae.
899 In: Ettl, H., Gärtner, G., Gerloff, J., Heynig, H., Mollenhauer, D. (Eds.),
900 Süßwasserflora von Mitteleuropa, Band 2/1. Gustav Fischer Verlag, Stuttgart, New
901 York.
- 902 Krammer, K., Lange-Bertalot, H., 1988. Bacillariophyceae. 2: Teil:
903 Bacillariaceae, Epithemiaceae, Surirellaceae. In: Ettl, H., Gärtner, G., Gerloff, J.,
904 Heynig, H., Mollenhauer, D. (Eds.), Süßwasserflora von Mitteleuropa, Band 2/2.
905 Gustav Fischer Verlag, Stuttgart, New York.
- 906 Krammer, K., Lange-Bertalot, H., 1991a. Bacillariophyceae. 3: Teil: Centrales,
907 Fragilariaceae, Eunotiaceae. In: Ettl, H., Gärtner, G., Gerloff, J., Heynig, H.,
908 Mollenhauer, D. (Eds.), Süßwasserflora von Mitteleuropa, Band 2/3. Gustav Fischer
909 Verlag, Stuttgart, Jena.
- 910 Krammer, K., Lange-Bertalot, H., 1991b. Bacillariophyceae. 4: Teil:
911 achnanthaceae. In: Ettl, H., Gärtner, G., Gerloff, J., Heynig, H., Mollenhauer, D.
912 (Eds.), Süßwasserflora von Mitteleuropa, Band 2/4. Gustav Fischer Verlag, Stuttgart,
913 Jena.
- 914 Lamb, A.L., Vane, C.H., Wilson, G.P., Rees, J.G., Moss-Hayes, V.L., 2007.
915 Assessing $\delta^{13}\text{C}$ and C/N ratios from organic material in archived cores as Holocene

916 sea level and palaeoenvironmental indicators in the Humber Estuary, UK. *Marine*
 917 *geology* 244, 109-128.

918 Lawler, A., 2009. *Beyond the Yellow River: How China Became China*. *Science*
 919 325, 930-935.

920 Leng, M.J., Lewis, J.P., 2017. C/N ratios and Carbon Isotope Composition of
 921 Organic Matter in Estuarine Environments, in: Weckström, K., Saunders, K.M., Gell,
 922 P.A., Skilbeck, C.G. (Eds.), *Applications of Palaeoenvironmental Techniques in*
 923 *Estuarine Studies*. Springer Netherlands, Dordrecht, pp. 213-237.

924 Li, C.X., Chen, Q.Q., Zhang, J.Q., Yang, S., Fan, D., 2000. Stratigraphy and
 925 paleoenvironmental changes in the Yangtze Delta during the Late Quaternary. *Journal*
 926 *of Asian Earth Sciences* 18, 453-469.

927 Li, C.X., Wang, P.X., Sun, H.P., Zhang, J.Q., Fan, D., Deng, B., 2002. Late
 928 Quaternary incised-valley fill of the Yangtze delta (China): its stratigraphic
 929 framework and evolution. *Sedimentary Geology* 152, 133-158.

930 Li, L., Zhu, C., Lin, L., Zhao, Q., Shi, G., Zhu, H., 2008. Transgression records
 931 between 7500-5400 BC on the stratum of the Luotudun site in Yixing, Jiangsu
 932 province. *Acta Geographica Sinica* 63, 1189-1197.

933 Liu, L., Chen, X., 2012. *The archaeology of China: From the late paleolithic to*
 934 *the early bronze age*. Cambridge World Archaeology, Cambridge.

935 Liu, Q., Yang, X., Anderson, N.J., Liu, E., Dong, X., 2012. Diatom ecological
 936 response to altered hydrological forcing of a shallow lake on the Yangtze floodplain,
 937 SE China. *Ecohydrology* 5, 316-325.

938 Liu, Y., Sun, Q., Thomas, I., Zhang, L., Finlayson, B., Zhang, W., Chen, J., Chen,
939 Z., 2015. Middle Holocene coastal environment and the rise of the Liangzhu City
940 complex on the Yangtze delta, China. *Quaternary Research* 84, 326-334.

941 Long, T., Qin, J., Atahan, P., Mooney, S., Taylor, D., 2014. Rising waters: New
942 geoarchaeological evidence of inundation and early agriculture from former
943 settlement sites on the southern Yangtze Delta, China. *Holocene* 24, 546-558.

944 Meyers, P.A., 2003. Applications of organic geochemistry to paleolimnological
945 reconstructions: a summary of examples from the Laurentian Great Lakes. *Organic*
946 *Geochemistry* 34, 261-289.

947 Middelburg, J.J., Nieuwenhuize, J., 1998. Carbon and nitrogen stable isotopes in
948 suspended matter and sediments from the Schelde Estuary. *Marine chemistry* 60,
949 217-225.

950 Mo, D., Zhao, Z., Xu, J., Li, M., 2011. Holocene Environmental Changes and the
951 Evolution of the Neolithic Cultures in China, in: Martini, P.I., Chesworth, W. (Eds.),
952 *Landscapes and Societies: Selected Cases*. Springer Netherlands, Dordrecht, pp,
953 299-319.

954 Müller, A., Mathesius, U., 1999. The palaeoenvironments of coastal lagoons in
955 the southern Baltic Sea, I. The application of sedimentary C_{org}/N ratios as source
956 indicators of organic matter. *Palaeogeography, Palaeoclimatology, Palaeoecology* 145,
957 1-16.

958 Müller, A., Voss, M., 1999. The palaeoenvironments of coastal lagoons in the
959 southern Baltic Sea, II. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ ratios of organic matter—sources and sediments.
960 *Palaeogeography, Palaeoclimatology, Palaeoecology* 145, 17-32.

- 961 Owen, R.B., Crossley, R., 1992. Spatial and temporal distribution of diatoms in
962 sediments of Lake Malawi, Central Africa, and ecological implications. *Journal of*
963 *Paleolimnology* 7, 55-71.
- 964 Patalano, R., Wang, Z., Leng, Q., Liu, W., Zheng, Y., Sun, G., Yang, H., 2015.
965 Hydrological changes facilitated early rice farming in the lower Yangtze River Valley
966 in China: A molecular isotope analysis. *Geology* 43.
- 967 Qin, J., Taylor, D., Atahan, P., Zhang, X., Wu, G., Dodson, J., Zheng, H.,
968 Itzstein-Davey, F., 2011. Neolithic agriculture, freshwater resources and rapid
969 environmental changes on the lower Yangtze, China. *Quaternary Research* 75, 55-65.
- 970 Reimer, P.J., Baillie, M.G.L., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G.,
971 Ramsey, C.B., Buck, C.E., Burr, G.S., Edwards, R.L., 2009. *IntCal09 and Marine09*
972 *Radiocarbon Age Calibration Curves, 0-50,000 Years cal. BP.* *Radiocarbon* 51,
973 1111-1150.
- 974 Renberg, I., 1990. A procedure for preparing large sets of diatom slides from
975 sediment cores. *Journal of Paleolimnology* 4, 87-90.
- 976 Ryu, E., Yi, S., Lee, S.-J., 2005. Late Pleistocene-Holocene paleoenvironmental
977 changes inferred from the diatom record of the Ulleung Basin, East Sea (Sea of Japan).
978 *Marine Micropaleontology* 55, 157-182.
- 979 Ryves, D.B., Juggins, S., Fritz, S.C., Battarbee, R.W., 2001. Experimental
980 diatom dissolution and the quantification of microfossil preservation in sediments.
981 *Palaeogeography Palaeoclimatology Palaeoecology* 172, 99-113.
- 982 Ryves, D.B., Clarke, A.L., Appleby, P.G., Amsinck, S.L., Jeppesen, E.,
983 Landkildehus, F., Anderson, N.J., 2004. Reconstructing the salinity and environment

984 of the Limfjord and Vejler. Canadian Journal of Fisheries & Aquatic Sciences 61,
985 1988-2006 (1919).

986 Ryves, D.B., Battarbee, R.W., Juggins, S., Fritz, S.C., Anderson, N.J., 2006.
987 Physical and chemical predictors of diatom dissolution in freshwater and saline lake
988 sediments in North America and West Greenland. Limnology and Oceanography 51,
989 1355-1368.

990 Ryves, D.B., Battarbee, R.W., Fritz, S.C., 2009. The dilemma of disappearing
991 diatoms: Incorporating diatom dissolution data into palaeoenvironmental modelling
992 and reconstruction. Quaternary Science Reviews 28, 120-136.

993 Song, B., Li, Z., Saito, Y., Okuno, J.I., Li, Z., Lu, A., Hua, D., Li, J., Li, Y.,
994 Nakashima, R., 2013. Initiation of the Changjiang (Yangtze) delta and its response to
995 the mid-Holocene sea level change. Palaeogeography Palaeoclimatology
996 Palaeoecology 388, 81-97.

997 Stabell, B., 1985. The development and succession of taxa within the diatom
998 genus *Fragilaria* Lyngbye as a response to basin isolation from the sea. Boreas 14,
999 273-286.

1000 Stanley, D.J., 2000. Radiocarbon Dates in China's Holocene Yangtze Delta:
1001 Record of Sediment Storage and Reworking, Not Timing of Deposition. Journal of
1002 Coastal Research 16, 1126-1132.

1003 Tada, R., Irino, T., Koizumi, I., 1999. Land-ocean linkages over orbital and
1004 millennial timescales recorded in Late Quaternary sediments of the Japan Sea.
1005 Paleoceanography 14, 236-247.

1006 Talma, A.S., Vogel, J.C., 1993. A Simplified Approach to Calibrating C14 Dates.
 1007 Radiocarbon 35 317-322.

1008 Tao, J., Chen, M.T., Xu, S., 2006. A Holocene environmental record from the
 1009 southern Yangtze River delta, eastern China. Palaeogeography, Palaeoclimatology,
 1010 Palaeoecology 230, 204-229.

1011 Thornton, S.F., McManus, J., 1994. Application of Organic Carbon and Nitrogen
 1012 Stable Isotope and C/N Ratios as Source Indicators of Organic Matter Provenance in
 1013 Estuarine Systems: Evidence from the Tay Estuary, Scotland. Estuarine Coastal &
 1014 Shelf Science 38, 219-233.

1015 van der Werff, A., Huls, H., 1957-1974, Reprinted 1976. Diatomeënfloora van
 1016 Nederland. Koeltz Science Publishers, Koenigstein.

1017 Voß, M., Struck, U., 1997. Stable nitrogen and carbon isotopes as indicator of
 1018 eutrophication of the Oder river (Baltic sea). Marine chemistry 59, 35-49.

1019 Vos, P.C., de Wolf, H., 1993. Diatoms as a tool for reconstructing sedimentary
 1020 environments in coastal wetlands; methodological aspects. Hydrobiologia 269,
 1021 285-296.

1022 Wang, C., Li, X., Lai, Z., Tan, X., Pang, S., Yang, W., 2009. Seasonal variations
 1023 of *Aulacoseira granulata* population abundance in the Pearl River Estuary. Estuarine
 1024 Coastal & Shelf Science 85, 585-592.

1025 Wang, J., Chen, X., Zhu, X.H., Liu, J.L., Chang, W.Y.B., 2001. Taihu Lake,
 1026 lower Yangtze drainage basin: evolution, sedimentation rate and the sea level.
 1027 Geomorphology 41, 183-193.

- 1028 Wang, Y., Cheng, H., Edwards, R.L., He, Y., Kong, X., An, Z., Wu, J., Kelly,
1029 M.J., Dykoski, C.A., Li, X., 2005. The Holocene Asian monsoon: links to solar
1030 changes and North Atlantic climate. *Science* 308, 854-857.
- 1031 Wang, Z., Zhuang, C., Saito, Y., Chen, J., Zhan, Q., Wang, X., 2012. Early
1032 mid-Holocene sea-level change and coastal environmental response on the southern
1033 Yangtze delta plain, China: implications for the rise of Neolithic culture. *Quaternary*
1034 *Science Reviews* 35, 51-62.
- 1035 Wang, Z., Zhan, Q., Long, H., Saito, Y., Gao, X., Wu, X., Li, L., Zhao, Y., 2013.
1036 Early to mid-Holocene rapid sea-level rise and coastal response on the southern
1037 Yangtze delta plain, China. *Journal of Quaternary Science* 28, 659-672.
- 1038 Wang, Z., Ryves, D.V., Lei, S., Nian, X., Lv, Y., Tang, L., Wang, L., Wang, J.,
1039 Chen, J., (accepted). Middle Holocene marine flooding and human response in the
1040 south Yangtze coastal plain, East China. *Quaternary Science reviews*.
- 1041 Wilson, G.P., Lamb, A.L., Leng, M.J., Gonzalez, S., Huddart, D., 2005. $\delta^{13}\text{C}$ and
1042 C/N as potential coastal palaeoenvironmental indicators in the Mersey Estuary, UK.
1043 *Quaternary Science Reviews* 24, 2015-2029.
- 1044 Witkowski, A., Dr, Ichtiologii, Lange-Bertalot, H., Metzeltin, D., 2000. *Diatom*
1045 *flora of marine coasts I*. A.R.G. Gantner Verlag K.G.
- 1046 Xu, P., 2015. Research on the archaeological cultures of Neolithic age in
1047 Ningzhen area and Huantaihu area. PhD Thesis, Jilin University, China (in Chinese,
1048 with English abstract).

- 1049 Yan, Q., Huang, S., 1987. Evolution of Holocene sedimentary environment in
1050 the Hangzhou-Jiaxing-Huzhou Plain. *Acta Geographica Sinica* 42, 1-15 (in Chinese,
1051 with English abstract).
- 1052 Yang, S., Tang, M., Yim, W.W.S., Zong, Y., Huang, G., Switzer, A.D., Saito, Y.,
1053 2011. Burial of organic carbon in Holocene sediments of the Zhujiang (Pearl River)
1054 and Changjiang (Yangtze River) estuaries. *Marine chemistry* 123, 1-10.
- 1055 Yu, S., Zhu, C., Song, J., Qu, W., 2000. Role of climate in the rise and fall of
1056 Neolithic cultures on the Yangtze Delta. *Boreas* 29, 157-165.
- 1057 Zeng, L., Lesch, S.M., Grieve, C.M., 2003. Rice growth and yield respond to
1058 changes in water depth and salinity stress. *Agricultural Water Management* 59, 67-75.
- 1059 Zhan, Q., Wang, Z.H., Xie, Y., Xie, J., He, Z., 2012. Assessing C/N and $\delta^{13}\text{C}$ as
1060 indicators of Holocene sea level and freshwater discharge changes in the subaqueous
1061 Yangtze delta, China. *The Holocene* 22, 697-704.
- 1062 Zhang, J., Wu, Y., Jennerjahn, T.C., Ittekkot, V., He, Q., 2007. Distribution of
1063 organic matter in the Changjiang (Yangtze River) Estuary and their stable carbon and
1064 nitrogen isotopic ratios: Implications for source discrimination and sedimentary
1065 dynamics. *Marine chemistry* 106, 111-126.
- 1066 Zhang, X., Huang, D., Deng, H., Snape, C., Meredith, W., Zhao, Y., Du, Y.,
1067 Chen, X., Sun, Y., 2015. Radiocarbon dating of charcoal from the Bianjiashan site in
1068 Hangzhou: New evidence for the lower age limit of the Liangzhu Culture. *Quaternary*
1069 *Geochronology* 30, Part A, 9-17.

- 1070 Zheng, C.G., 2002. Environmental archaeology on the Temporal-Spatial
1071 distribution of Cultures sites in Taihu Lake area during 7 ka BP - 4 ka BP. PhD Thesis,
1072 Nanjing University, China (in Chinese, with English abstract).
- 1073 Zheng, Y. F., Chen, X.G., Ding, P., 2014. Studies on the archaeological paddy
1074 fields at Maoshan site in Zhejiang. *Quaternary Sciences*, 34, 85-96.
- 1075 Zheng, Y., Sun, G., Chen, X., 2012. Response of rice cultivation to fluctuating
1076 sea level during the Mid-Holocene. *Chinese Science Bulletin* 57, 370-378.
- 1077 Zhou, H., Zheng, X., 2000. The Impact of Environmental Changes on the
1078 Development of Prehistoric Civilization: The Decline of the Ancient Liang zhu
1079 Culture in the Southern Plain of Yangtze River Delta. *Journal of East China Normal*
1080 *University (Natural Science)* 4, 71-77 (in Chinese, with English abstract).
- 1081 Zhu, N., 2006. Impacts of the rice farming on cultural development in the Taihu
1082 and Hangzhou Bay area, In: Shanghai Museum (Eds.), Symposium on the civilization
1083 course of the lower reach region of the Yangzte Rvier. Shanghai Book/paint publishers,
1084 Shanghai, pp, 69-88 (in Chinese).
- 1085 Zhuang, Y., Ding, P., French, C., 2014. Water management and agricultural
1086 intensification of rice farming at the late-Neolithic site of Maoshan, Lower Yangtze
1087 River, China. *Holocene* 24, 531-545.
- 1088 Zong, Y.Q., 2004. Mid-Holocene sea-level highstand along the southeast coast
1089 of China. *Quaternary International* 117, 55-67.
- 1090 Zong, Y., Chen, Z., Innes, J.B., Chen, C., Wang, Z., Wang, H., 2007. Fire and
1091 flood management of coastal swamp enabled first rice paddy cultivation in east China.
1092 *Nature* 449, 459-462.

1093 Zong, Y.Q., Innes, J.B., Wang, Z.H., Chen, Z.Y., 2011. Mid-Holocene coastal
1094 hydrology and salinity changes in the east Taihu area of the lower Yangtze wetlands,
1095 China. *Quaternary Research* 76, 69-82.

1096 Zong, Y., Wang, Z., Innes, J., Chen, Z., 2012a. Holocene environmental change
1097 and Neolithic rice agriculture in the lower Yangtze region of China: A review.
1098 *Holocene* 22, 623-635.

1099 Zong, Y., Innes, J.B., Wang, Z., Chen, Z., 2012b. Environmental change and
1100 Neolithic settlement movement in the lower Yangtze wetlands of China. *Holocene* 22,
1101 659-673.

1102 Zuo, X., Lu, H., Jiang, L., Zhang, J., Yang, X., Huan, X., He, K., Wang, C., Wu,
1103 N., 2017. Dating rice remains through phytolith carbon-14 study reveals
1104 domestication at the beginning of the Holocene. *Proceedings of the National*
1105 *Academy of Sciences of the United States of America* 114, 6486.

1
2 **Supplementary Table 1**
3 Dating results for collected cores from previous studies

Core name	Location	Depth (m)	Dating material	Conventional age (yr BP)	Calibrated age (2 δ , cal. yr BP)	Reference
97A	/	2.1	Peat	/	2305-2000 (1 δ)	Zhou and Zheng, 2000
		4.11	Plant fragment	/	7660-75809 (1 δ)	
		5.78	Plant fragment	/	7630-7580 (1 δ)	
		10.2	Shell	/	8330-8220 (1 δ)	
		13.8	Plant fragment	/	8425-8365 (1 δ)	
JLQ	/	2.5	Bulk organic	4975 $\pm$ 70	5597-5797	Hong, 1991; Zong et al., 2011
		11.2	Bulk organic	7400 $\pm$ 80	8039-8372	
DS, SL, GT	/	/	/	/	/	Hong, 1991; Yan and Huang, 1987
XL	30 °24'05" N, 120 °01'38" E	4.87	Wood	3940 $\pm$ 40	4420-4220	Liu et al., 2015
		5.9	Seed	5700 $\pm$ 40	7580-7440	
		7.6	Plant material	6420 $\pm$ 40	7560-7530	
LTD	31 °21' N, 121°19 '42' E	2.16	Mud	5281 $\pm$ 175	6400-5660	Li et al., 2008
		2.45	Peat	5464 $\pm$ 165	6570-5910	
		2.64	Peat	6016 $\pm$ 200	7310-6410	
		2.88	Peat	8036 $\pm$ 190	9430-8510	
W1	/	0.3	Pollen	5020 $\pm$ 75	5638-5910	Chang et al., 1994; Wang et al., 2001
		0.60-0.70	Mud	6145 $\pm$ 370	6206-7703	

		1.60-1.70	Mud	8575 $\pm$ 410	8544-10593	
E2	/	0.2	Mud	2575 $\pm$ 110	2354-2868	Chang et al., 1994; Wang et al., 2001
		0.32	Mud	4765 $\pm$ 70	5433-5607	
		0.45	Mud	5495 $\pm$ 75	6175-6448	
		1.14-1.25	Mud	5970 $\pm$ 170	6441-7184	
		2.26	Mud	6575 $\pm$ 75	7410-7585	
YJD	30 °59'45" N, 120 °35'31" E	0.7-0.72	Pollen residue	3570 $\pm$ 40	3816-3977	Zong et al., 2012b
		1.0-1.02	Pollen residue	5600 $\pm$ 40	6299-6453	
PW	30 °57'30" N, 120 °38'25" E	1.59-1.61	Pollen residue	2700 $\pm$ 40	2750-2868	Innes et al., 2014
		1.85-1.87	Pollen residue	4430 $\pm$ 40	4871-5075	
		2.25-2.27	Pollen residue	4720 $\pm$ 40	5323-5417	
		3.75-3.77	Peat	6290 $\pm$ 50	7153-7320	
WJB	30 °49'04" N, 120 °36'56" E	3.5-3.52	Charcoal	6410 $\pm$ 60	7245-7434	Qin et al., 2011
		3.5-3.52	Pollen residue	6250 $\pm$ 70	6970-7315	
		3.74-3.76	Peat	6600 $\pm$ 50	7431-7569	
		3.74-3.76	Peat	6630 $\pm$ 60	7429-7590	
		6-6.02	Clay	7350 $\pm$ 60	8024-8318	

ZX-1	/	1.1	Organic-rich mud	2580 ± 60	2462-2794	Chen et al., 2005; Tao et al., 2006
		4.5	Shell	4160 ± 40	3875-4221	
		8.8	Organic-rich mud	6850 ± 80	7570-7853	
CXS	31 °22'44" N, 120 °47'30" E	0.5-0.52	Pollen residue	2960 ± 40	2985-3234	Zong et al., 2012b
		0.72-0.74	Pollen residue	4190 ± 40	4609-4768	
CD	31 °24'16" N, 120 °50'37" E	0.9-0.92	Pollen residue	2000 ± 40	1867-2060	Zong et al., 2012b
		1.2-1.22	Plant fragment	3160 ± 40	3324-3459	
SQ	31 °11'50" N, 121 °06'25" E	0.75-0.77	Pollen residue	5410 ± 40	6175-6296	Zong et al., 2011
		1.40-1.42	Pollen residue	7310 ± 50	8005-8204	
QP	31 °07'44" N, 120 °54'39" E	0.62-0.64	Pollen residue	1827 ± 35	1695-1865	Atahan et al., 2008; Itzstein-Davey et al., 2007
		1.20-1.22	Pollen residue	2152 ± 35	2038-2185	
		1.82-1.84	Pollen residue	2386 ± 35	2342-2493	
		2.10-2.12	Pollen residue	3853 ± 40	4215-4409	
		2.38-2.40	Pollen residue	5780 ± 30	6498-6656	
		2.42-2.44	Pollen residue	5600 ± 40	6299-6453	
		2.5-2.52	Pollen residue	5114 ± 35	5749-5830	
		2.58-2.60	Pollen residue	4920 ± 35	5594-5718	

TYL	31 °11'54" N, 121 °06'40" E	1.2-1.22	Pollen residue	2330 ± 40	2304-2469	Zong et al., 2012b
		1.6-1.62	Pollen residue	2390 ± 40	2338-2505	
		2.0-2.02	Pollen residue	3680 ± 40	3897-4096	
TCM	31 °01'48" N, 121 °05'30" E	1.72-1.74	Pollen residue	4140 ± 40	4567-4825	Zong et al., 2011
		2.12-2.14	Pollen residue	5230 ± 40	5912-6029	
GFL	31 °03'52" N, 121 °11'30" E	0.28	/	1070 ± 50	909-1088	Wang et al., 2012
		0.62-0.64	Wood	945 ± 30	794-925	
		0.70-0.72	Pollen residue	2057 ± 30	1945-2118	
		0.75	/	2710 ± 40	2752-2879	
		0.88-0.90	Charcoal	2453 ± 30	2362-2544	
		1.00	/	4110 ± 40	4521-4730	
		1.05	/	4600 ± 40	5346-5466	
		1.74-1.76	Charcoal	5517 ± 55	6276-6403	
TL	30 °53'12" N, 121 °18'42" E	2.1-2.12	Pollen residue	5800 ± 40	6493-6678	Zong et al., 2011
		2.3-2.32	Pollen residue	7390 ± 50	8152-8344	

References

- Atahan, P., Itzstein-Davey, F., Taylor, D., Dodson, J., Qin, J., Zheng, H., Brooks, A., 2008. Holocene-aged sedimentary records of environmental changes and early agriculture in the lower Yangtze, China. *Quaternary Science Reviews* 27, 556-570.
- Chang, W.Y.B., Xu, X.M., Yang, J.R., Liu, J.L., 1994. Evolution in Taihu Lake ecosystem as evidence of changes in sediment profiles. *Journal of Lake Science* 6, 217-226.
- Chen, Z., Wang, Z., Schneiderman, J., Taol, J., Cail, Y., 2005. Holocene climate fluctuations in the Yangtze delta of eastern China and the Neolithic response. *The Holocene* 15, 915-924.

- Hong, X., 1991. Origin and evolution of the Taihu Lake. *Marine Geology and Quaternary Geology* 11, 87-99 (in Chinese, with English abstract).
- Innes, J.B., Zong, Y., Wang, Z., Chen, Z., 2014. Climatic and palaeoecological changes during the mid-to Late Holocene transition in eastern China: high-resolution pollen and non-pollen palynomorph analysis at Pingwang, Yangtze coastal lowlands. *Quaternary Science Reviews* 99, 164-175.
- Itzstein-Davey, F., Atahan, P., Dodson, J., Taylor, D., Zheng, H., 2007. Environmental and cultural changes during the terminal Neolithic: Qingpu, Yangtze delta, eastern China. *The Holocene* 17, 875-887.
- Liu, Y., Sun, Q., Thomas, I., Zhang, L., Finlayson, B., Zhang, W., Chen, J., Chen, Z., 2015. Middle Holocene coastal environment and the rise of the Liangzhu City complex on the Yangtze delta, China. *Quaternary Research* 84, 326-334.
- Li, L., Zhu, C., Lin, L., Zhao, Q., Shi, G., Zhu, H., 2008. Transgression records between 7500-5400 BC on the stratum of the Luotuodun site in Yixing, Jiangsu province. *Acta Geographica Sinica* 63, 1189-1197.
- Qin, J., Taylor, D., Atahan, P., Zhang, X., Wu, G., Dodson, J., Zheng, H., Itzstein-Davey, F., 2011. Neolithic agriculture, freshwater resources and rapid environmental changes on the lower Yangtze, China. *Quaternary Research* 75, 55-65.
- Tao, J., Chen, M.T., Xu, S., 2006. A Holocene environmental record from the southern Yangtze River delta, eastern China. *Palaeogeography, Palaeoclimatology, Palaeoecology* 230, 204-229.
- Yan, Q., Huang, S., 1987. Evolution of Holocene sedimentary environment in the Hangzhou-Jiaxing-Huzhou Plain. *Acta Geographica Sinica* 42, 1-15 (in Chinese, with English abstract).
- Wang, J., Chen, X., Zhu, X.H., Liu, J.L., Chang, W.Y.B., 2001. Taihu Lake, lower Yangtze drainage basin: evolution, sedimentation rate and the sea level. *Geomorphology* 41, 183-193.
- Wang, Z., Zhuang, C., Saito, Y., Chen, J., Zhan, Q., Wang, X., 2012. Early mid-Holocene sea-level change and coastal environmental response on the southern Yangtze delta plain, China: implications for the rise of Neolithic culture. *Quaternary Science Reviews* 35, 51-62.
- Zhou, H., Zheng, X., 2000. The Impact of Environmental Changes on the Development of Prehistoric Civilization: The Decline of the Ancient Liang zhu Culture in the Southern Plain of Yangtze River Delta. *Journal of East China Normal University (Natural Science)* 4, 71-77 (in Chinese, with English abstract).
- Zong, Y.Q., Innes, J.B., Wang, Z.H., Chen, Z.Y., 2011. Mid-Holocene coastal hydrology and salinity changes in the east Taihu area of the lower Yangtze wetlands, China. *Quaternary Research* 76, 69-82.
- Zong, Y., Innes, J.B., Wang, Z., Chen, Z., 2012b. Environmental change and Neolithic settlement movement in the lower Yangtze wetlands of China. *Holocene* 22, 659-673.