Supporting Information

**Mild Cleavage of Aryl Mesylates: Methanesulfonate as
Convenient Protecting Group for Phenols**

*Tobias Ritter, Kyrill Stanek, Igor Larrosa, and Erick M. Carreira**

Laboratorium für Organische Chemie, ETH-Zürich,

ETH Hönggerberg, HCI, LOC

Materials and Methods

All reagents and reaction solvents were obtained and purified before use when necessary. Reactions were performed in oven-dried glassware under a slight positive pressure of dry argon. Air- and moisture-sensitive liquids and solutions were transferred under inert atmosphere by syringe. Syringes employed were all polypropylene disposable. Organic solutions were concentrated by rotary evaporation at 40 °C. Diisopropylamine was distilled from KOH. THF and CH₂Cl₂ were dried and purified through activated alumina columns as described by Grubbs *et al.*¹

Flash chromatography: silica gel 60 (230 – 400 mesh, 0.04 – 0.063 mm) from *Merck* at rt and 0.3 – 0.5 mbar air pressure.

Thin layer chromatography (TLC): *Merck* 0.25 mm silica gel 60 *F₂₅₄* TLC glass plates. Visualization of the developed chromatogram was performed by either UV fluorescence at 254 nm or oxidative stain by ceric ammonium molybdate solution or KMnO₄ / NaHCO₃ water solution.

Melting points: *Büchi* 510 apparatus. All melting points were measured in open capillaries and are uncorrected.

IR spectra: *Perkin-Elmer Paragon 1000 Fourier Transform single beam spectrophotometer*. The samples were prepared as thin films on NaCl salt plates and are reported as absorption maxima in cm⁻¹.

NMR spectra: ¹H, ¹³C NMR spectra were recorded on a *Varian Mercury-300* spectrometer operating at 300 MHz for ¹H-NMR and 75 MHz for ¹³C-NMR. Spectra were referenced internally to residual proton solvent signals. Data for ¹H are reported as follows: chemical shift (δ , ppm), multiplicity *s*(singlet), *d*(doublet), *t*(triplet), *q*(quartet), and *m*(multiplet), coupling constant (*J*, Hz), integration. Data for ¹³C are reported in terms of chemical shift (δ , ppm), coupling constant (*J*, Hz) (where appropriate).

Mass spectra: EI mass spectra were performed by the MS service at ETH Zürich.

Elemental analyses: Mikrolabor für Organische Chemie at ETH-Zürich.

Lithium diisopropylamid is prepared by the following procedure: To THF at -78 °C (15.0 ml) is added diisopropylamine (1.97 ml, 1.20 equiv) followed by n-BuLi (1.61 M in hexane, 7.76 ml, 1.00 equiv). The solution is stirred at -78 °C for 30 min.

Supporting Information

General Procedure for the Preparation of Arylmethanesulfonates.

To a solution of phenol (1.00 equiv) in CH_2Cl_2 (conc= 1 M) at 0 °C is added NEt_3 (1.50 equiv). To this solution or suspension is added MsCl (1.25 equiv) over a period of 15 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water and the phases are separated. The aqueous phase is extracted with CH_2Cl_2 . The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is purified by chromatography on silica gel or recrystallized.

General Procedure for the Deprotection of Arylmethanesulfonates.

To a solution of the methanesulfonate aryl ester (1.00 equiv) in THF (conc= 0.1 M- 1.0 M) at -78 °C to 23 °C is added a freshly prepared solution of LDA (1.60-1.80 equiv per mesylate group) in THF. After 1 -30 min 5% aqueous HCl or aqueous NH_4Cl is added followed by ethyl acetate. The phases are separated and the aqueous phase is extracted with ethyl acetate or CH_2Cl_2 . The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is purified by chromatography on silica gel or recrystallized to afford the pure phenols.

To a solution of 4-fluorophenol (4.00 g, 35.7 mmol, 1.00 equiv) in CH_2Cl_2 (35 ml) at 0 °C is added NEt_3 (7.46 ml, 53.6 mmol, 1.50 equiv). To this solution is added MsCl (3.45 ml, 44.6 mmol, 1.25 equiv) over a period of 15 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water (10 ml). The phases are separated and the aqueous phase is extracted with CH_2Cl_2 (2* 5 ml). The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1 to 1/1 gradient) to afford the title compound as a clear liquid in 97% yield.

R_f = 0.49 (hexane/ ethyl acetate 1/1).

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.28-7.24 (m, 2 H), 7.13-7.07 (m, 2 H), 3.14 (s, 3 H).

$^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 161.4, (J = 247 Hz), 145.1, 124.0, (J = 9 Hz), 117.0 (J = 24 Hz), 37.5.

IR (thin film): 3034, 2942, 1500, 1369, 1195, 1156.

HRMS (EI): Calcd' for $(\text{C}_7\text{H}_7\text{FO}_3\text{S}^+)$, 190.0848; found, 190.0848.

Anal. Calcd. for $\text{C}_7\text{H}_7\text{FO}_3\text{S}$: C, 44.21; H, 3.71; F, 9.99. Found: C, 44.38; H, 4.00.

Supporting Information

To a solution of **methanesulfonic acid 4-fluoro-phenyl ester** (382 mg, 2.00 mmol, 1.00 equiv) in THF (4.0 ml) at 0 °C is added a freshly prepared solution of LDA in THF (3.20 mmol, 1.60 equiv). After 1 min 5% aqueous HCl is added followed by CH₂Cl₂ (20 ml). The phases are separated and the aqueous phase is extracted with CH₂Cl₂. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1) to afford the title compound as a clear liquid in 92% yield.

¹H-NMR (300 MHz, CDCl₃): δ 6.96-6.87 (*m*, 2 H), 6.81-6.49 (*m*, 2 H), 4.58 (*s*, 1 H).

This analytical data corresponds to:

Fukuhara, T.; Akiyama, Y.; Yoneda, N.; Tada, T.; Hara, S. *Tetrahedron Lett.* **2002**, 43, 6583-6585.

To a solution of 3-bromophenol (4.08 g, 23.6 mmol, 1.00 equiv) in CH₂Cl₂ (25 ml) at 0 °C is added NEt₃ (4.83 ml, 34.7 mmol, 1.50 equiv). To this solution is added MsCl (2.23 ml, 28.9 mmol, 1.25 equiv) over a period of 15 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water (10 ml). The phases are separated and the aqueous phase is extracted with CH₂Cl₂ (2* 5 ml). The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1 to 1/1 gradient) to afford the title compound as a clear liquid in 97% yield.

R_f = 0.50 (hexane/ ethyl acetate 1/1).

¹H-NMR (300 MHz, CDCl₃): δ 7.49-7.45 (*m*, 2 H), 7.33-7.22 (*m*, 2 H), 3.16 (*s*, 3 H).

¹³C-NMR (75 MHz, CDCl₃): δ 149.6, 131.3, 130.9, 125.7, 123.1, 121.1, 38.9.

IR (thin film): 3094, 3032, 2940, 1581, 1468, 1371, 1197, 1156, 890.

HRMS (EI): Calcd^a for (C₇H₇BrO₃S)⁺, 249.9299; found, 249.9276.

Anal. Calcd. for C₇H₇BrO₃S: C, 33.48; H, 2.81; found: C, 33.53; H, 2.87.

Supporting Information

To a solution of **methanesulfonic acid 3-bromo-phenyl ester** (502 mg, 2.00 mmol, 1.00 equiv) in THF (2.0 ml) at 0 °C is added a freshly prepared solution of LDA in THF (3.00 mmol, 1.50 equiv). After 5 min 5% aqueous HCl (10 ml) is added followed by ethyl acetate (20 ml). The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1) to afford the title compound in 95% yield.

¹H-NMR (300 MHz, CDCl₃): δ 7.14-7.01 (*m*, 3 H), 6.79-6.74 (*m*, 1 H), 4.99 (*s*, 1 H).

Fischer, A.; Henderson, G.; Narayanan, C. *Can. J. Chem.* **1983**, 63, 1045-1052

Proof of purity: **Anal.** Calcd. for C₆H₅BrO: C, 41.65; H, 2.91; found: C, 41.93; H, 3.13.

For further analytical data, see: Brossi, A.; Hsu, F.-L.; Rice, K. C.; Rozwadowska, M. D.; Schmidhammer, H. *Helv. Chim. Acta* **1981**, 64, 1672-1681.

To a solution of 2-iodophenol (2.00 g, 9.10 mmol, 1.00 equiv) in CH₂Cl₂ (10 ml) at 0 °C is added NEt₃ (1.90 ml, 13.6 mmol, 1.50 equiv). To this solution is added MsCl (0.920 ml, 11.4 mmol, 1.25 equiv) over a period of 20 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water (10 ml). The phases are separated and the aqueous phase is extracted with CH₂Cl₂. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1) to afford the title compound in 94% yield.

R_f = 0.42 (hexane/ ethyl acetate 2/1).

¹H-NMR (300 MHz, CDCl₃): δ 7.84 (*dd*, *J* = 7.8 Hz, 1.6 Hz, 1 H), 7.47-7.35 (*m*, 2 H), 7.03 (*m*, 1 H), 3.28 (*s*, 3 H).

¹³C-NMR (75 MHz, CDCl₃): δ 149.7, 140.4, 130.2, 129.0, 123.4, 89.9, 39.5.

IR (thin film): 3030, 2937, 1573, 1463, 1440, 1371, 1204, 1169, 969, 861.

HRMS (EI): Calcd' for (C₇H₇IO₃S⁺), 297.9161; found, 297.9156.

Anal. Calcd. for C₇H₇IO₃S: C, 28.20; H, 2.37; found: C, 28.32; H, 2.41.

Supporting Information

To a solution of **methanesulfonic acid 2-iodo-phenyl ester** (596 mg, 2.00 mmol, 1.00 equiv) in THF (2.0 ml) at 0 °C is added a freshly prepared solution of LDA in THF (3.20 mmol, 1.60 equiv). After 1 min 5% aqueous HCl (10 ml) is added followed by ethyl acetate (20 ml). The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1) to afford the title compound in 93% yield.

¹H-NMR (300 MHz, CDCl₃): δ 7.66 (*dd*, *J* = 7.9 Hz, 1.5 Hz, 1 H), 7.24 (*m*, 1 H), 6.99 (*dd*, *J* = 8.2 Hz, 1.5 Hz, 1 H), 6.68 (*m*, 1 H), 5.29 (*s*, 1 H).

This analytical data corresponds to: Kitagawa, H.; Shibata, T.; Matsuo, J.; Mukaiyama, T. *Bull. Chem. Soc. Jpn.* **2002**, 75, 339 - 346.

Proof of purity: **Anal.** Calcd. for C₆H₅IO: C, 32.76; H, 2.29; found: C, 33.06; H, 2.35.

To a solution of 2,6-dimethoxyphenol (1.54 g, 10.0 mmol, 1.00 equiv) in CH₂Cl₂ (20 ml) at 0 °C is added NEt₃ (2.09 ml, 15.0 mmol, 1.50 equiv). To this solution is added MsCl (0.970 ml, 12.5 mmol, 1.25 equiv) over a period of 5 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added aqueous NH₄Cl (8 ml). The phases are separated and the aqueous phase is extracted with CH₂Cl₂. The combined organic phases are washed with 2 N HCl (20 ml), water (20 ml), and brine (20 ml), dried (Na₂SO₄) and concentrated in vacuo to afford the title compound in 100% yield.

mp: 98°C.

R_f = 0.42 (hexane/ ethyl acetate 2/1).

¹H-NMR (300 MHz, CDCl₃): δ 7.16 (*t*, *J* = 8.4 Hz, 1 H), 6.61 (*d*, *J* = 8.4 Hz, 2 H), 3.88 (*s*, 6 Hz), 3.29 (*s*, 3 H).

¹³C-NMR (75 MHz, CDCl₃): δ 153.1, 128.1, 127.3, 104.9, 56.3, 39.9.

IR (thin film): 3037, 2977, 1608, 1508, 1493, 1448, 1365, 1307, 1111, 870, 779.

Anal. Calcd. for C₉H₁₂O₅S: C, 46.54; H, 5.21; found: C, 46.60; H, 5.38.

Supporting Information

To a solution of **methanesulfonic acid 2,6-dimethoxy-phenyl ester** (112 mg, 0.500 mmol, 1.00 equiv) in THF (5.0 ml) at 0 °C is added a freshly prepared solution of LDA in THF (0.901 mmol, 1.80 equiv). After 1 min 5% aqueous HCl (10 ml) is added followed by ethyl acetate (20 ml). The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with CH₂Cl₂ to afford the title compound in 95% yield.

¹H-NMR (300 MHz, CDCl₃): δ 6.87 (*t*, *J* = 8.1 Hz, 1 H), 6.57 (*d*, *J* = 8.1 Hz, 1 H), 5.55 (*s*, 1 H), 3.86 (*s*, 6H).

This analytical data corresponds to:

Jing, X.; Gu, W.; Bie, P.; Ren, X.; Pan, X. *Synth. Commun.*, **2001**, *31*, 861-868.

To a solution of 3-dimethylaminophenol (2.00 g, 14.6 mmol, 1.00 equiv) in CH₂Cl₂ (15 ml) at 0 °C is added NEt₃ (6.10 ml, 21.9 mmol, 1.50 equiv). To this solution is added MsCl (1.47 ml, 18.2 mmol, 1.25 equiv) over a period of 20 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water (10 ml). The phases are separated and the aqueous phase is extracted with CH₂Cl₂. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1) to afford the title compound in 92% yield.

mp: 50 °C

R_f = 0.27 (hexane/ ethyl acetate 2/1).

¹H-NMR (300 MHz, CDCl₃): δ 7.24-7.18 (*m*, 1 H), 6.65-6.57 (*m*, 3 H), 3.11 (*s*, 3 H), 2.96 (*s*, 6 H).

¹³C-NMR (75 MHz, CDCl₃): δ 152.1, 150.9, 130.3, 111.2, 108.9, 105.7, 40.6, 37.4.

IR (thin film): 2938, 2812, 1611, 1570, 1506, 1363, 1179, 1130.

MS (ESI): Calcd⁺ for (C₇H₇IO₃SH⁺), 216.1; found, 216.2.

Anal. Calcd. for C₉H₁₃O₃S: C, 50.21; H, 6.09; N, 6.51; found: C, 50.36; H, 6.15; N, 6.53.

Supporting Information

To a solution of **methanesulfonic acid 3-dimethylamino-phenyl ester** (432 mg, 2.00 mmol, 1.00 equiv) in THF (2.0 ml) at 0 °C is added a freshly prepared solution of LDA in THF (3.20 mmol, 1.60 equiv). After 1 min NH_4Cl (0.5 g) is added and the suspension is concentrated. To the residue is added CH_2Cl_2 and water. The phases are separated and the aqueous phase is extracted with CH_2Cl_2 . The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1) to afford the title compound in 89% yield.

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.13-7.07 (*m*, 1 H), 6.37-6.32 (*m*, 1 H), 6.24-6.18 (*m*, 2H), 4.94 (*s*, 1 H), 2.92 (*s*, 6H).

This analytical data corresponds to: Jacquesy, J.-C.; Jouannetaud, M.-P.; Morellet, G.; Vidal, Y. *Bull. Soc. Chim. Fr.* **1986**, 625-629.

Proof of purity: **Anal.** Calcd. for $\text{C}_8\text{H}_{11}\text{NO}$: C, 70.04; H, 8.08; found: C, 69.81; H, 8.22.

To a solution of 5-fluoro-2-nitrophenol (1.57 g, 10.0 mmol, 1.00 equiv) in acetone (20 ml) at 23 °C is added K_2CO_3 (1.66 g, 12.0 mmol, 1.20 equiv). To this suspension is added MsCl (1.48 g, 11.0 mmol, 1.10 equiv). The suspension is stirred at 23 °C until all starting material was consumed as assayed by TLC. The suspension is concentrated and ethyl acetate (40 ml) and brine (10 ml) are added. The phases are separated and the organic phase is concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ CH_2Cl_2 (2/3) to afford the title compound in 96% yield.

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 8.14 (*dd*, $J = 9.3$ Hz, 5.6 Hz, 1 H), 7.31 (*dd*, $J = 8.4$ Hz, 2.8 Hz, 1 H), 7.18 (*ddd*, $J = 9.3$ Hz, 6.84 Hz, 2.5 Hz, 1 H), 3.39 (*s*, 3 H).

$^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 164.6 (*d*, $J = 258$ Hz), 114.9 (*d*, $J = 22$ Hz), 113.3 (*d*, $J = 22$ Hz), 39.4.

IR (thin film): 3584, 3117, 2943, 1560, 1534, 1486, 1354, 1274, 1186, 1146, 1082, 997, 808, 614.

MS (EI): Calcd' for $(\text{C}_7\text{H}_6\text{FNO}_5\text{S}^+)$, 235.1; found, 235.0.

Anal. Calcd. for $\text{C}_7\text{H}_6\text{FNO}_5\text{S}$: C, 35.75; H, 2.57, N, 5.96; found: C, 35.70; H, 2.75; N, 6.17.

Supporting Information

To a solution of **methanesulfonic acid 5-fluoro-2-nitro-phenyl ester** (112 mg, 0.500 mmol, 1.00 equiv) in THF (5.0 ml) at -78 °C is added a freshly prepared solution of LDA in THF (0.900 mmol, 1.80 equiv). After 1 min aqueous NH₄Cl (4 ml) is added followed by CH₂Cl₂. The phases are separated and the aqueous phase is extracted with CH₂Cl₂. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with CH₂Cl₂ to afford the title compound in 67% yield.

mp: 33°C-35°C

This analytical data corresponds to:

Nicholson, H. *J. Chem. Soc.* **1939**, 1405-1408.

For further analytical data, see: Gerson, H.; Clarke, D. D.; Gershon, M. *Monatsh. Chem.* **1995**, 126, 1161-1166.

To a solution of 2-methoxy-4-nitrophenol (3.00 g, 17.7 mmol, 1.00 equiv) in CH₂Cl₂ (20 ml) at 0 °C is added NEt₃ (3.70 ml, 26.6 mmol, 1.50 equiv). To this solution is added MsCl (1.79 ml, 22.2 mmol, 1.25 equiv) over a period of 30 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water (10 ml). The phases are separated and the aqueous phase is extracted with CH₂Cl₂. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is recrystallized from ethyl acetate to afford the title compound in 97% yield.

mp: 108 °C

R_f = 0.36 (hexane/ ethyl acetate 3/2).

¹H-NMR (300 MHz, CDCl₃): δ 7.89-7.85 (m, 2 H), 7.46-7.42 (m, 1 H), 4.00 (s, 3 H), 3.26 (s, 3 H).

¹³C-NMR (75 MHz, CDCl₃): δ 152.2, 147.2, 143.0, 125.0, 116.8, 108.4, 56.9, 39.2.

IR (thin film): 3093, 3045, 2944, 1534, 1497, 1360, 1348, 1326, 1268, 1162, 1020.

HRMS (EI): Calcd. for (C₈H₉NO₆S⁺), 247.0151; found, 247.0150.

Anal. Calcd. for C₈H₉NO₆S: C, 38.87; H, 3.67, N, 5.67; found: C, 38.83; H, 3.76; N, 5.58.

Supporting Information

To a solution of **methanesulfonic acid 2-methoxy-4-nitro-phenyl ester** (494 mg, 2.00 mmol, 1.00 equiv) in THF (4.0 ml) at 23 °C is added a freshly prepared solution of LDA in THF (3.20 mmol, 1.60 equiv). After 30 min 5% aqueous HCl (10 ml) is added followed by ethyl acetate. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (2/1) to afford the title compound in 65% yield.

¹H-NMR (300 MHz, CDCl₃): δ 7.89 (*dd*, *J* = 8.7 H, 2.5 Hz, 1 H), 7.77 (*d*, *J* = 2.6 Hz, 1 H), 6.98 (*d*, *J* = 8.7 Hz, 1 H), 6.22 (*s*, 1 H), 4.00 (*s*, 3 H).

Proof of purity: **Anal.** Calcd. for C₇H₇NO₄: C, 49.71; H, 4.17, N, 8.28; found: C, 49.84; H, 4.29; N, 8.16.

This analytical data corresponds to:

Harvison, P.; Forte, A. J.; Nelson, S; Sidney, D. *J. Med. Chem.* **1986**, 29, 1737-1743.

To a solution of 2-5-di-*tert*-butylhydroquinone (4.00 g, 18.0 mmol, 1.00 equiv) in CH₂Cl₂ (20 ml) at 0 °C is added NEt₃ (7.52 ml, 54.0 mmol, 3.00 equiv). To this solution is added MsCl (3.63 ml, 45.0 mmol, 2.50 equiv) over a period of 15 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water (10 ml). The phases are separated and the aqueous phase is extracted with CH₂Cl₂. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is recrystallized from ethyl acetate to afford the title compound in 77% yield.

mp: 173 °C

R_f = 0.15 (hexane/ ethyl acetate 5/1).

¹H-NMR (300 MHz, CDCl₃): δ 7.58 (*s*, 2 H), 3.25 (*s*, 6 H), 1.39 (*s*, 18 H).

¹³C-NMR (75 MHz, CDCl₃): δ 146.2, 140.5, 120.9, 39.4, 34.8, 30.4.

IR (thin film): 3019, 2942, 2873, 1501, 1347, 1193, 1157, 1110, 871.

HRMS (EI): Calcd' for (C₁₆H₂₆O₆S₂⁺), 378.1171; found, 378.1171.

Anal. Calcd. for C₁₆H₂₆O₆S₂: C, 75.63; H, 9.97; found: C, 75.36; H, 9.91.

Supporting Information

To a solution of **methanesulfonic acid 2,5-di-tert-butyl-4-methanesulfonyloxy-phenyl ester** (757 mg, 2.00 mmol, 1.00 equiv) in THF (15.0 ml) at 0 °C is added a freshly prepared solution of LDA in THF (6.40 mmol, 3.20 equiv). After 30 min 5% aqueous HCl (9 ml) is added followed by ethyl acetate. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (5/1) to afford the title compound in 70% yield.

mp: 216 °C.

This analytical data corresponds to:

Fujita, S.; Sano, K. *J. Org. Chem.* 1979, 44, 2647-2651.

Proof of purity: **Anal.** Calcd. for $\text{C}_{14}\text{H}_{22}\text{O}_2$: C, 75.63; H, 9.97; found: C, 75.36; H, 9.91.

For further analytical data, see: Shulgin, A. T.; Kerlinger, H. O. *Tetrahedron Lett.*, **1965**, 6, 3355-3359.

To a solution of 4-acetamidophenol (4.00 g, 26.5 mmol, 1.00 equiv) in CH_2Cl_2 (50 ml) at 0 °C is added NEt_3 (5.53 ml, 39.7 mmol, 1.50 equiv). To this solution is added MsCl (2.67 ml, 33.0 mmol, 1.25 equiv) over a period of 30 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water (25 ml) and ethyl acetate (600 ml). The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is recrystallized from ethyl acetate to afford the title compound in 78% yield.

mp: 177 °C

R_f = 0.41 (MeOH/ CH_2Cl_2 8/92).

¹H-NMR (300 MHz, acetone d_6): δ 9.31 (s, 1 H), 7.72 (m, 2 H), 7.25 (m, 2 H), 3.24 (s, 3 H), 2.08 (s, 3 H).

¹³C-NMR (75 MHz, acetone d_6): δ 168.7, 145.4, 139.2, 123.1, 120.7, 37.2, 24.1.

IR (thin film): 3300, 1664, 1538, 1365, 1194, 1156, 834.

MS (EI): Calcd' for $(\text{C}_9\text{H}_{11}\text{NO}_4\text{S}^+)$, 229.0; found, 229.0.

Anal. Calcd. for $\text{C}_9\text{H}_{11}\text{NO}_4\text{S}$: C, 47.15; H, 4.84, N, 6.11; found: C, 47.33; H, 4.94, N, 6.19.

Supporting Information

To a solution of **methanesulfonic acid 4-acetamido-phenyl ester** (459 mg, 2.00 mmol, 1.00 equiv) in THF (20 ml) at 23 °C is added a freshly prepared solution of LDA in THF (12.0 mmol, 6.0 equiv). After 30 min 5% aqueous HCl (10 ml) is added followed by ethyl acetate. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with CH_2Cl_2 / MeOH (95/5 to 90/10 gradient) to afford the title compound in 66% yield.

mp: 168 °C-172 °C

This data corresponds to: Ritter, F. O. *J. Am. Chem. Soc.* **1934**, 56, 975-976.

To a solution of 2-bromo-3-hydroxypyridine (1.74 g, 10.0 mmol, 1.00 equiv) in acetone (20 ml) at 0 °C is added K_2CO_3 (1.66 g, 12.0 mmol, 1.20 equiv). To this suspension is added MsCl (1.48 g, 11.0 mmol, 1.10 equiv). The suspension is stirred at 23 °C for 1 h. The suspension is concentrated and ethyl acetate (40 ml) and brine (10 ml) are added. The aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ CH_2Cl_2 (4/1) to afford the title compound in 94% yield.

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 8.34 (*dd*, J = 4.7 Hz, 1.6 Hz, 1 H), 7.76 (*dd*, J = 8.1 Hz, 1.9 Hz, 1 H), 7.36 (*dd*, J = 8.1 Hz, 4.7 Hz, 1 H), 3.33 (*s*, 3 H).

$^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 148.0, 143.9, 135.7, 132.1, 127.8, 39.5.

IR (thin film): 3132, 3032, 2937, 1568, 1409, 1372, 1175, 1062, 860.

MS (EI): Calcd' for ($\text{C}_9\text{H}_9\text{NO}_4$), 250.9252; found, 250.9251.

To a solution of **methanesulfonic acid 2-bromo-pyridin-3-yl ester** (252 mg, 1.00 mmol, 1.00 equiv) in THF (10 ml) at 23 °C is added a freshly prepared solution of LDA in THF (1.80 mmol, 1.80 equiv). After 30 min 5% aqueous NH_4Cl (4 ml) is added followed by ethyl acetate. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na_2SO_4) and

Supporting Information

concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with CH_2Cl_2 / ethyl acetate (3/2) to afford the title compound in 57% yield.

mp: 184 °C

This data corresponds to: Hasseberg, H.-A.; Gerlach, H. *Helv. Chim. Acta* **1988**, *71*, 957-963.

To a solution of 4-hydroxyquinoline (0.500 g, 3.44 mmol, 1.00 equiv) in CH_2Cl_2 (5.0 ml) at 0 °C is added NEt_3 (0.720 ml, 5.17 mmol, 1.50 equiv). To this solution is added MsCl (0.350 ml, 4.31 mmol, 1.25 equiv) over a period of 30 min. The suspension is stirred at 0 °C for additional 15 min. To the suspension is added water (25 ml) and CH_2Cl_2 . The phases are separated and the aqueous phase is extracted with CH_2Cl_2 . The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with toluene/ ethyl acetate (3/1 to 1/1 gradient) to afford the title compounds in 52% and 36%, respectively.

mp: 70 °C

R_f = 0.24 (toluene/ ethyl acetate 3/2).

¹H-NMR (300 MHz, CDCl_3): δ 8.87 (*d*, *J* = 5.0 Hz, 1 H), 8.08 (*m*, 2 H), 7.73 (*m*, 1 H), 7.58 (*m*, 2 H), 7.43 (*d*, *J* = 5.0 Hz, 1 H), 3.26 (*s*, 3 H).

¹³C-NMR (75 MHz, CDCl_3): δ 152.7, 151.0, 150.4, 130.8, 129.8, 127.9, 122.2, 121.5, 112.2, 38.8.

IR (thin film): 3032, 2936, 1599, 1502, 1372, 1181, 1154, 1085, 893.

HRMS (EI): Calcd⁺ for $(\text{C}_{10}\text{H}_9\text{NO}_3\text{S}^+)$, 223.0303; found, 223.0312.

Anal. Calcd. for $\text{C}_{10}\text{H}_9\text{NO}_3\text{S}$: C, 53.80; H, 4.06, N, 6.27; found: C, 53.95; H, 4.13, N, 6.27.

Supporting Information

mp: 145 °C

R_f = 0.17 (toluene/ ethyl acetate 3/2).

¹H-NMR (300 MHz, CDCl₃): δ 8.42 (*dd*, *J* = 7.8 Hz, 1.6 Hz, 1 H), 8.27 (*d*, *J* = 8.7 Hz, 1 H), 8.24 (*m*, 1 H), 7.64 (*m*, 1 H), 7.50 (*m*, 1 H), 6.53 (*d*, *J* = 8.7 Hz, 1 H), 3.37 (*s*, 3 H).

¹³C-NMR (75 MHz, CDCl₃): δ 178.0, 137.2, 137.0, 133.3, 127.6, 126.4, 125.9, 117.5, 112.4, 41.2.

IR (thin film): 3009, 2926, 1640, 1640, 1609, 1468, 1362, 1173, 1104.

HRMS (EI): Calcd^a for (C₁₀H₉NO₃S)⁺, 223.2; found, 223.2.

Anal. Calcd. for C₁₀H₉NO₃S: C, 53.80; H, 4.06, N, 6.27; found: C, 53.88; H, 4.08, N, 6.27.

To a solution of **methanesulfonic acid quinolin-4-yl ester** (112 mg, 0.490 mmol, 1.00 equiv) in THF (1.0 ml) at 0 °C is added a freshly prepared solution of LDA in THF (0.800 mmol, 1.60 equiv). After 5 min 5% aqueous HCl (5 ml) is added followed by CH₂Cl₂. The phases are separated and the aqueous phase is extracted with CH₂Cl₂. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by trituration in acetone to afford the title compound in 68% yield.

mp: 202 °C-206 °C

Proof of purity: **Anal.** Calcd. for C₉H₇NO: C, 74.47; H, 4.86; N, 9.65; found: C, 74.38; H, 5.07; N, 9.86.

This analytical data corresponds to:

Katagiri, N.; Niwa, R.; Furuya, Y.; Kato, T. *Chem. Pharm. Bull.* **1983**, *6*, 1833-1841.

Supporting Information

To a solution of **1-methanesulfonyl-4a,8a-dihydro-1H-quinolin-4-one** (110 mg, 0.500 mmol, 1.00 equiv) in THF (4.0 ml) at 0 °C is added a freshly prepared solution of LDA in THF (0.800 mmol, 1.60 equiv). After 5 min 5% aqueous HCl (5 ml) is added followed by ethyl acetate. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by trituration in acetone to afford the title compound in 72% yield.

mp: 202 °C-206 °C

To a solution of the **1** (1.40 g, 2.63 mmol, 1.00 equiv) in THF (53.0 ml) at -78 °C is added a freshly prepared solution of LDA in THF (10.5 mmol, 4.00 equiv). The suspension is warmed to 0°C. After 5 min 0.5 ml MeOH are added, followed by aqueous NH₄Cl and ethyl acetate. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (1/2 to 1/3 gradient) to afford the title compound in 82% yield.

R_f = 0.31 (hexane/ ethyl acetate 1/2)

α_D = 54°, (CHCl₃, c=0.280, 22.8 °C.

mp: 134 °C.

¹H-NMR (300 MHz, MeOD): δ 7.42 (*m*, 5 H), 6.95 (*d*, *J* = 8.7 Hz), 6.76 (*d*, *J* = 8.7 Hz), 6.15 (*m*, 1 H), 5.99-5.84 (*m*, 2 H), 5.14-4.64 (*m*, 6 H), 3.98 (*m*, 3 H), 3.78 (*m*, 2 H), 3.50 (*m*, 2 H), 3.24-3.78 (*m*, 3 H), 1.72 (*m*, 1 H), 1.36 (*t*, *J* = 7.1 Hz, 3 H), 1.22 (*m*, 1 H), 1.39 (*d*, *J* = 7.1 Hz, 3H).

¹³C-NMR (75 MHz, MeOD): δ 149.1, 145.6, 140.5, 138.6, 137.3, 132.2, 129.6, 129.0, 128.1, 126.7, 123.9, 114.9, 114.4, 102.5, 84.2, 82.7, 68.4, 64.9, 63.4, 44.0, 42.7, 30.8, 24.2, 14.7.

IR (thin film): 3307, 3083, 2978, 1637, 1597, 1488, 1440, 1406, 1295, 1104, 1065, 1050.

HRMS (Maldi): Calcd⁺ for (C₂₇H₃₄O₆Na⁺), 477.2253; found, 477.2243.

Anal. Calcd. for C₂₇H₃₄O₆: C, 71.34; H, 7.54; found: C, 71.17; H, 7.61.

Supporting Information

To a solution of **2** (200 mg, 0.288 mmol, 1.00 equiv) in THF (3.0 ml) at -78 °C is added a freshly prepared solution of LDA in THF (0.461 mmol, 1.60 equiv). After 2 min aqueous NH₄Cl is added followed by ethyl acetate. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (3/2) to afford the title compound in 79% yield.

R_f = 0.31 (hexane/ ethyl acetate 3/2).

α_D = 54°, (CHCl₃, c = 0.395, 28.3 °C).

¹H-NMR (300 MHz, CDCl₃): δ 7.42 (*m*, 5 H), 6.95 (*d*, *J* = 8.7 Hz), 6.82 (*d*, *J* = 8.7 Hz), 6.18 (*m*, 1 H), 5.99-5.84 (*m*, 2 H), 5.14-4.84 (*m*, 6 H), 4.72 (*d*, *J* = 9.3 Hz), 4.63 (*d*, *J* = 9.3 Hz), 3.93 (*q*, *J* = 7.1 Hz, 2 H), 3.74 (*m*, 1 H), 3.59-3.38 (*m*, 2 H), 3.00 (*s*, 3 H), 1.44 (*t*, *J* = 7.1 Hz, 3 H), 1.39 (*d*, *J* = 7.1 Hz, 3H), 1.29 (*s*, 9 H).

¹³C-NMR (75 MHz, CDCl₃): δ 177.8, 148.6, 145.4, 139.5, 137.5, 136.6, 131.6, 129.7, 128.7, 126.6, 124.4, 116.8, 116.6, 114.0, 103.1, 82.4, 81.5, 70.4, 64.9, 43.5, 40.5, 39.1, 39.0, 31.0, 27.6, 23.4, 16.0.

IR (thin film): 3448, 2977, 2935, 1731, 1637, 1481, 1353, 1291, 1174, 913, 897.

HRMS (Maldi): Calcd⁺ for (C₃₃H₄₄O₉SN⁺), 639.2604; found, 639.2590.

To a solution of the aldehyde **6** (24.2 g, 80.0 mmol, 1.00 equiv) in DMF (32 ml) at 0 °C is added Rieke zinc.² To this suspension is added a solution of bromide **5** (37.5 g, 100 mmole, 1.25 equiv) in DMF (10 ml) at 0 °C over a period of 3.5 h. After addition the slurry is warmed to 23 °C and stirred for additional 30 min. The suspension is diluted with ethyl acetate (100 ml) and poured onto aqueous NH₄Cl solution. The phases are separated and the aqueous phase is extracted with ethyl acetate (3* 40 ml). The combined organic phases are washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue is purified by

Supporting Information

chromatography on silica gel eluting with hexane/ ethyl acetate (3/1) collecting fractions at R_f = 0.55 (hexane/ ethyl acetate 3/1). This material is dissolved in THF (120 ml) at 0 °C and TBAF in THF (1.00 M, 76.2 ml, 1.25 equiv) is added. The solution is stirred at 0 °C for 30 min. The solution is diluted with ethyl acetate (250 ml) and aqueous NaHCO_3 solution (250 ml). The phases are separated and the aqueous phase is extracted with ethyl acetate (3* 40 ml). The combined organic phases are washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue is purified by chromatography on silica gel eluting with hexane/ ethyl acetate (3/2) to afford the title compound in 41% yield along with a diastereomer (23%).

R_f = 0.41 (hexane/ ethyl acetate 1/1).

α_D = -5.5°, (CHCl_3 , c = 0.325, 25.2 °C).

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.33 (*d*, J = 9.7 Hz, 1 H), 7.20 (*d*, J = 9.3 Hz, 1 H), 6.37-6.25 (*m*, 1 H), 6.02-5.93 (*m*, 1 H), 5.16-4.85 (*m*, 4 H), 4.02-3.85 (*m*, 5 H), 3.64-3.43 (*m*, 4 H), 3.11 (*s*, 3 H), 2.67 (*s*, 1 H), 2.45 (*s*, 1 H), 2.08 (*dd*, J = 15.3 Hz, 5.7 Hz, 1 H), 1.42-1.15 (*m*, 13 H).

$^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 149.0, 143.4, 141.1, 137.0, 136.6, 132.2, 124.6, 122.1, 118.1, 115.5, 99.9, 78.2, 77.9, 70.4, 67.6, 62.7, 44.6, 37.8, 34.5, 30.4, 30.0, 26.2, 22.5, 15.8.

IR (thin film): 3535, 2979, 2936, 1476, 1369, 1173, 967.

HRMS (Maldi): Calcd¹ for ($\text{C}_{24}\text{H}_{36}\text{O}_6\text{SNa}^+$), 507.2029; found, 507.2018.

¹ A. B. Pangborn, M. A. Giardello, R. H. Grubbs, R. K. Rosen, F. J. Timmers, *Organometallics* **1996**, *15*, 1518.

² R. D. Rieke, P. T. J. Li, T. P. Burns, S. T. Uhm *J. Org. Chem.* **1981**, *46*, 4323-4324.