Supporting Information

A Mild Manganese(III) Acetate Catalyzed Allylic Oxidation: Application to Simple and Complex Alkenes

Tony K. M. Shing, Ying Y. Yeung, and Pak L. Su*

Department of Chemistry, The Chinese University of Hong Kong, Shatin, NT, Hong Kong, China.

E-mail: tonyshing@cuhk.edu.hk

RECEIVED DATE (to be automatically inserted after your manuscript is accepted if
required according to the journal that you are submitting your paper to)

Appendices

S2–S14 ----- **Experimental procedures**

S15–S19 ----- **Table S1: Solvent variation**

Table S2: Oxidation of benzylic carbons

Table S3: More examples of allylic oxidation of simple alkenes

Figure S1: TBHP concentration variation

**Figure S2: Manganese(III) acetate dihydrate concentration
variation**

Figure S3: EPR studies

**Scheme S1: Evidence for the existence of *tert*-butyl peroxy ether
intermediate**

Scheme S2: π -Stabilization of allylic radical for simple alkene

Scheme S3: π -Stabilization of allylic radical for carene

S20–S33 ----- **NMR spectra**

S34–S35 ----- **References**

Experimental procedures

General. Melting points were determined with a Reichert apparatus and are reported in degrees Celsius uncorrected. Optical rotations were measured with a Perkin-Elmer model 341 polarimeter, operating at 589nm. IR spectra were recorded on a Nicolet 205 FT-IR spectrophotometer as thin film on potassium bromide discs. EPR spectra were obtained with a Bruker EMX EPR spectrometer. ^1H NMR and ^{13}C NMR spectra were obtained in CDCl_3 solutions with a Bruker DPX300 spectrophotometer at 300.13 MHz and 75.47 MHz, respectively. NMR Spectra are referenced to residual chloroform (δ 7.26 ppm, ^1H ; δ 77.0 ppm, ^{13}C). Chemical shifts are reported in ppm (δ); multiplicities are indicated by s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet) and br (broad). Coupling constants, J , are reported in Hertz. Gas chromatography was performed on a Hewlett Packard 5890 Series II with 30-m Supelcowax 10 column and a flame ionization detector. Thin-layer chromatography (TLC) was performed on Merck aluminum-precoated plates of silica gel 60 F₂₅₄. Visualization was obtained by spraying with 5% (w/v) dodecamolybdophosphoric acid in ethanol and subsequent heating. Column chromatography was performed with silica gel 60 (230–400) mesh. The *tert*-butyl hydroperoxide (TBHP) 5-6 M solution in Decane was purchased from Aldrich Chemical Co., Milwaukee. Ethyl acetate was dried and distilled from anhydrous magnesium sulfate. Other reagents are purchased from commercial suppliers and are used without purification.

Literature preparations

The following compounds were prepared by literature methods: 3 β ,17 β -diacetoxyandrost-5-ene (**1d**),¹ 3 β -acetyloxypregn-5-ene (**1e**),² 3-[(1,1-dimethylethyl)dimethylsilyl]oxycholest-5-ene (**1f**),³ 3-[(1,1-dimethylethyl)dimethylsilyl]oxy-5-androsten-17-one (**1g**),⁴ 3-[(1,1-dimethylethyl)dimethylsilyl]oxy-pregn-5-en-20-one (**1h**),⁵ 3 β ,17 β -Bis[(1,1-dimethylethyl)dimethylsilyl]oxyandrost-5-ene (**1i**),⁶ 3 β -Acetoxy-17 β -hydroxyandrost-5-ene (**1j**),⁷ 17-oxo-17 α -aza-D-homoandrost-5-en-3 β -yl acetate (**3a**),⁸ 3-acetyloxy-(3 β ,25R)-spirost-5-ene (**5a**),⁹

3-[(1,1-dimethylethyl)dimethylsilyl]-(3 β ,25R)-spirost-5-ene (5b)¹⁰ and 1,4-Dioxaspiro[4.5]dec-6-ene (19).¹¹

General procedure for manganese (III) acetate catalyzed allylic oxidation of alkenes.

Method A. (Entries 1–18): To a solution of alkene (0.27 mmol) in EtOAc (2 ml) was added *tert*-butyl hydroperoxide in decane (1.39 mmol) and 3 Å molecular sieves (100 mg). The mixture was stirred for 30 min under nitrogen atmosphere at RT. Manganese(III) acetate dihydrate (0.027 mmol) was added and the mixture was stirred for 48 h at RT. The solution was filtered through a pad of celite. Concentration of the filtrate followed by column chromatography afforded the α,β -unsaturated ketone.

Method B. (Entries 19–27): To a solution of alkene (0.36 mmol) in EtOAc (3 ml) was added *tert*-butyl hydroperoxide in decane (1.39 mmol) and 3 Å molecular sieves (100 mg). The mixture was stirred for 30 min under nitrogen atmosphere at RT. Manganese(III) acetate dihydrate (0.036 mmol) was added. The solution was degassed and filled with oxygen (balloon). The mixture was stirred for 48 h at RT and filtered through a pad of celite. Concentration of the filtrate followed by column chromatography afforded the α,β -unsaturated ketone.

Representative procedure for complex alkene: A solution of cholesteryl acetate (1a). (100 mg, 0.27 mmol), EtOAc (2 ml) and TBHP in decane (0.25 ml, 1.39 mmol) was stirred with 3 Å molecular sieves (100 mg) for 30 min under nitrogen. Manganese(III) acetate dihydrate (7.4 mg, 0.027 mmol) was added. The reaction was stirred for 48 h, diluted with 2 ml of diethyl ether and filtered through a thin pad of Celite. The filtrate was concentrated under reduced pressure and purified by column chromatography (*n*-hexane:Et₂O = 3:1).

Representative procedure for simple alkene: A solution of (1S)-(–)- α -Pinene (17). (50 mg,

0.36 mmol), EtOAc (3 ml) and TBHP in decane (0.33 ml, 1.39 mmol) was stirred with 3 Å molecular sieves (100 mg) for 30 min under nitrogen. Manganese(III) acetate dihydrate (9.8 mg, 0.0366 mmol) was added. The mixture was degassed and filled with oxygen (balloon). The reaction was stirred for 48 h, diluted with 2 ml of diethyl ether and filtered through a thin pad of Celite. The filtrate was concentrated under reduced pressure and purified by column chromatography (*n*-hexane:Et₂O = 4:1).

Procedure for the one gram scale allylic oxidation of cholesteryl acetate (1a): To a solution of cholesteryl acetate (**1a**) (1.0 g, 2.32 mmol) in EtOAc (20 ml) was added *tert*-butyl hydroperoxide in decane (2.1 mL, 11.6 mmol) and 3 Å molecular sieves (1.0 g). The mixture was stirred for 30 min under nitrogen atmosphere at RT. Manganese(III) acetate dihydrate (62.5 mg, 0.20 mmol) was added in portion (as small amount of heat would be given out) and the mixture was stirred for 48 h at RT. The solution was filtered through a pad of celite. Concentration of the filtrate followed by column chromatography afforded 3β-hydroxycholest-5-en-7-one acetate (**2a**) (883 mg, 85%).

Procedure for the determination of the oxygen gas evolved during allylic oxidation reaction^{12,13}: To a solution of cholesteryl acetate (**1a**) (858 mg, 2 mmol) in EtOAc (20 ml), 3 Å molecular sieves (800 mg) was added *tert*-butyl hydroperoxide in decane (1.8 mL, 10.0 mmol) under nitrogen. The mixture was stirred for 20 min at RT under nitrogen. Manganese(III) acetate dihydrate (74 mg, 0.20 mmol) was added. An empty balloon was equipped to capture oxygen generated during the course of the reaction. After 48 h at RT, the balloon was filled and the gas was passed through aq. potassium pyrogallate. The colour of the solution changed from pale yellow to deep brown which verified the existence of oxygen.

Procedure for the 2,6-di-*tert*-butyl-4-methylphenol (BHT) terminated allylic oxidation: Allylic oxidation of cholesteryl acetate (**1a**) was conducted under the conditions of “Method A”. After 2 h

at RT, 2,6-di-*tert*-butyl-4-methylphenol (BHT) (5 mg) was added to the reaction mixture and the reaction was terminated based on TLC analysis. After 48 h at RT, only trace amount of 3 β -hydroxycholest-5-en-7-one acetate (**2a**) was isolated with starting material recovered.

Procedure for the determination of *tert*-butyloxy radical during the allylic oxidation by gas chromatographic analysis: Allylic oxidation of cyclohexene was conducted under the conditions of “Method B”. After 24 h at RT, the solution was analyzed by gas chromatography and an acetone signal (retention time = 1.707 min) was observed which indicated the existence of *tert*-butyloxy radical.¹⁴

Procedure for the determination of *tert*-butylperoxy radical during the allylic oxidation by EPR analysis: Allylic oxidation of cyclohexene (**34**) was conducted under the conditions of “Method A”. After 5 h at RT, 5,5-dimethyl-1-pyrroline-*N*-Oxide (DMPO) (2 ml, 0.88 M in DMSO) was injected into the reaction mixture and the suspension was stirred for a further 10 min. The solution was then subjected to EPR measurement and the hyperfine constants of the quartet signal ($a^N = 14.0$ G, $a^H = 10.5$ G, $a^H = 1.4$ G) matched with the literature value of the spin adduct of DMPO and *tert*-butylperoxy radical (Figure S3.).³²

3 β -hydroxycholest-5-en-7-one acetate (2a**)¹⁵:** Isolated as a white solid; mp 154–155 °C; $[\alpha]_D^{20}$ –96.6 (c 1.00, CHCl₃); IR (thin film) 2947, 1728, 1668 cm⁻¹; ¹H NMR δ 5.69 (d, 1H, $J = 1.2$ Hz), 4.70 (m, 1H), 2.45 (s, 3H), 1.11 (s, 3H), 0.91 (d, 3H, $J = 6.6$ Hz), 0.85 (d, 3H, $J = 6.6$ Hz), 0.85 (d, 3H, $J = 6.6$ Hz), 0.67 (s, 3H); ¹³C NMR δ 202.4, 170.7, 164.3, 127.1, 72.6, 55.2, 50.4, 50.2, 45.8, 39.9, 39.1, 38.2, 36.6, 36.4, 36.1, 29.0, 28.4, 27.8, 26.7, 24.2, 23.2, 23.0, 21.6, 19.3, 17.7, 12.4. (The data were in full accordance with the literature values.)

3 β -Acetoxy-5-androsten-7,17-dione (2b**)¹⁶:** Isolated as a yellowish solid; mp 175–178 °C;

$[\alpha]_{\text{D}}^{20}$ –85.8 (*c* 1.70, CHCl₃); IR (thin film) 2954, 1742, 1673, 1628 cm⁻¹; ¹H NMR δ 5.72 (s, 1H), 4.68 (m, 1H), 2.02 (s, 3H), 1.20 (s, 3H), 0.80 (s, 3H); ¹³C NMR δ 220.6, 201.1, 170.6, 165.2, 126.8, 72.3, 50.3, 48.2, 46.0, 44.7, 38.8, 38.2, 36.3, 36.0, 31.0, 27.6, 24.5, 21.6, 20.9, 17.7, 14.1. (The data were in full accordance with the literature values.)

3 β -Acetoxypregn-5-ene-7,20-dione (2c)¹⁵: Isolated as a yellowish solid; mp 141–142 °C; $[\alpha]_{\text{D}}^{20}$ –66.6 (*c* 1.50, CHCl₃); IR (thin film) 2946, 1733, 1702, 1671 cm⁻¹; ¹H NMR δ 5.70 (d, 1H, *J* = 1.8 Hz), 4.69 (m, 1H), 2.12 (s, 3H), 2.04 (s, 3H), 1.20 (s, 3H), 0.60 (s, 3H); ¹³C NMR δ 210.2, 201.6, 170.7, 164.6, 126.9, 72.5, 62.7, 50.4, 50.0, 45.6, 44.8, 38.8, 38.2, 38.0, 36.4, 32.0, 27.7, 26.9, 24.0, 21.7, 21.5, 17.7, 13.7. (The data were in full accordance with the literature values.)

3 β ,17 β -diacetoxandrost-5-en-7-one(2d)¹⁵: Isolated as a white solid; mp 210–213 °C; $[\alpha]_{\text{D}}^{20}$ –132.0 (*c* 1.50, CHCl₃); IR (thin film) 2946, 1733, 1664 cm⁻¹; ¹H NMR δ 5.71 (d, 1H, *J* = 1.5 Hz), 4.71 (m, 1H), 4.62 (dd, 1H, *J* = 7.2, 5.4 Hz), 2.04 (s, 3H), 1.17 (s, 3H), 0.80 (s, 3H); ¹³C NMR δ 201.7, 171.6, 164.8, 126.9, 82.4, 72.5, 50.2, 45.4, 45.2, 43.5, 38.8, 38.2, 36.4, 36.2, 28.0, 27.7, 26.3, 21.7, 21.6, 21.1, 17.8, 12.5, 6.5. (The data were in full accordance with the literature values.)

3 β -acetyloxypregn-5-en-7-one 20-ethylene ketal (2e): Isolated as a white solid; mp 149–152 °C; $[\alpha]_{\text{D}}^{20}$ –95.2 (*c* 1.60, CHCl₃); IR (thin film) 2946, 2878, 1734, 1670 cm⁻¹; ¹H NMR δ 5.69 (d, 1H, *J* = 1.2 Hz), 4.70 (m, 1H), 3.97–3.86 (m, 4H), 2.06 (s, 3H), 1.30 (s, 3H), 1.20 (s, 3H), 0.76 (s, 3H); ¹³C NMR δ 202.3, 170.7, 164.4, 127.1, 112.4, 72.6, 65.4, 63.7, 57.2, 50.2, 50.1, 45.4, 43.0, 38.7, 38.2, 36.4, 30.1, 27.7, 26.3, 25.1, 23.9, 21.7, 21.4, 17.7, 13.4; MS (ESI) *m/z* (relative intensity) 439 ([M + Na]⁺, 100); HRMS (ESI) calcd for C₂₅H₃₆O₅ 439.2455 ([M + Na]⁺), found 439.2449. (The compound could be converted to **2c** by literature procedure¹⁷ and the data were in full accordance with those reported in literature.¹⁵)

3-[(1,1-dimethylethyl)dimethylsilyl]oxycholest-5-en-7-one (2f)¹⁸: Isolated as a white solid; mp 209–210 °C; $[\alpha]_D^{20}$ –66.4 (*c* 1.40, CHCl₃); IR (thin film) 2935, 2862, 1667 cm⁻¹; ¹H NMR δ 5.65 (s, 1H), 3.59 (m, 1H), 0.88 (s, 9H), 1.09 (s, 3H), 0.66 (s, 3H), 0.06 (s, 3H), 0.05 (s, 3H); ¹³C NMR δ 202.9, 166.3, 126.2, 71.8, 55.2, 50.4, 45.8, 43.5, 43.0, 39.9, 39.2, 38.8, 36.9, 36.6, 36.1, 32.2, 29.0, 28.4, 26.8, 26.3, 24.2, 23.2, 23.0, 21.6, 19.3, 17.7, 12.4, –4.1. (The data were in full accordance with the literature values.)

3-[(1,1-dimethylethyl)dimethylsilyl]oxy-5-androsten-7,17-dione (2g)¹⁹: Isolated as a white solid; mp 127–129 °C; $[\alpha]_D^{20}$ –51.6 (*c* 1.86, CHCl₃); IR (thin film) 1739, 1670 cm⁻¹; ¹H NMR δ 5.70 (s, 1H), 3.60 (m, 1H), 2.80 (m, 1H), 1.24 (s, 3H), 0.82 (s, 12H), 0.05 (s, 6H); ¹³C NMR δ 220.8, 201.6, 167.3, 126.0, 71.5, 50.5, 48.2, 46.1, 44.7, 43.0, 38.9, 36.7, 36.0, 32.0, 31.1, 26.2, 24.6, 20.9, 18.5, 17.8, 14.1, –4.2. (The data were in full accordance with the literature values.)

3-[(1,1-dimethylethyl)dimethylsilyl]oxypregn-5-ene-7,20-dione (2h): Isolated as a white solid; $[\alpha]_D^{20}$ –12.4 (*c* 0.70, CHCl₃); IR (thin film) 2938, 2856, 1667 cm⁻¹; ¹H NMR δ 5.67 (s, 1H), 3.58 (m, 1H), 2.12 (s, 3H), 1.18 (s, 3H), 0.64 (s, 3H), 0.85 (s, 12H), 0.06 (s, 6H); ¹³C NMR δ 210.1, 202.0, 166.6, 126.0, 71.6, 62.7, 50.4, 50.2, 45.6, 44.8, 42.9, 38.8, 38.1, 36.8, 32.1, 32.0, 26.8, 26.2, 24.0, 21.5, 18.5, 17.7, 13.6, –4.21, –4.24. (The compound **2h** could be converted to **2c** by literature procedure^[20] and the data were in full accordance with those reported in the literature.¹⁵)

3 β ,17 β -Bis[(1,1-dimethylethyl)dimethylsilyl]oxyandrost-5-en-7-one (2i)²¹: Isolated as a white solid; mp 166–168 °C; $[\alpha]_D^{20}$ –62.3 (*c* 0.66, CHCl₃); IR (thin film) 2933, 2856, 1667 cm⁻¹; ¹H NMR δ 5.66 (s, 1H), 3.63–3.52 (m, 2H), 1.23 (s, 3H), 0.95 (s, 9H), 0.93 (s, 9H), 0.71 (s, 3H), 0.06 (s, 6H), 0.00 (s, 6H); ¹³C NMR δ 202.7, 166.7, 126.1, 81.4, 71.7, 50.7, 45.8, 45.0, 44.1, 43.0, 38.9, 36.9,

36.5, 32.2, 31.4, 30.1, 26.3, 21.4, 18.6, 17.8, 11.7, -3.9, -4.1, -4.3. (The data were in full accordance with the literature values.)

3 β -Acetoxy-17 β -hydroxy-5-androsten-7-one (2j)²²: (68%, 31% recovery) Isolated as a white solid; mp 152–153 °C; $[\alpha]_D^{20}$ -100.8 (*c* 1.97, CHCl₃); IR (thin film) 3452, 2941, 2898, 1727 cm⁻¹; ¹H NMR δ 5.69 (d, 1H, *J* = 0.9 Hz), 4.70 (m, 1H), 3.64 (t, 1H, *J* = 8.4 Hz), 2.04 (s, 3H), 1.21 (s, 3H), 0.75 (s, 3H); ¹³C NMR δ 202.0, 170.7, 164.8, 126.8, 81.2, 72.5, 50.3, 45.6, 45.3, 43.7, 38.8, 38.2, 36.4, 35.9, 30.9, 27.7, 26.2, 21.6, 21.2, 17.7, 11.5. (The data were in full accordance with the literature values.)

3-[(1,1-dimethylethyl)dimethylsilyl]oxy-17-oxo-17a-aza-D-homoandrost-5-ene (3b): To a solution of **3a** (14 mg, 0.045 mmol) in MeOH (3 ml) was added K₂CO₃ (7.43 mg, 0.054 mmol) at rt. The reaction mixture was stirred at RT for 7 h. Deionized water (3 ml) was added into the mixture and the product was extracted with CH₂Cl₂ (3 ml $\times$ 4). The organic extracts were collected and concentrated under reduced pressure.

To a solution of the above residue in CH₂Cl₂ (3 ml) was added imidazole (9 mg, 0.13 mmol) and TBSCl (13.5 mg, 0.090 mmol) at RT under nitrogen. The reaction mixture was stirred at RT for further 3 h under nitrogen. Saturated ammonium chloride solution was added and the product was extracted with CH₂Cl₂ (3 ml $\times$ 4). The organic extracts were collected and concentrated under reduced pressure. Column chromatography of the residue using MeOH:CHCl₃ = 3:97 afforded **3b** (16 mg, 88%) as a white solid: mp 289–290 °C; $[\alpha]_D^{20}$ -57.9 (*c* 0.5, CHCl₃); IR (thin film) 3147, 3039, 2932, 1676, 1602 cm⁻¹; ¹H NMR δ 6.12 (br s, 1H), 5.31 (m, 1H), 3.47 (m, 1H), 1.39 (s, 3H), 0.98 (s, 3H), 0.88 (s, 9H), 0.08 (s, 6H); ¹³C NMR δ 141.8, 120.7, 72.8, 54.7, 50.0, 48.5, 42.9, 40.2, 37.5, 37.2, 32.7, 32.3, 31.7, 26.3, 22.5, 21.4, 20.5, 19.7, 18.7, -4.1; MS (ESI) *m/z* (relative intensity) 418 ([M + H]⁺, 100); HRMS (ESI) calcd for C₂₅H₄₃NO₂Si 418.3136 [M + H]⁺, found 418.3124.

7,17-Dioxo-17a-aza-D-homoandrost-5-en-3 β -yl acetate (4a)¹⁶: Isolated as a white solid; mp 279–280 °C; $[\alpha]_{\text{D}}^{20}$ –133.6 (*c* 0.30, CHCl₃); IR (thin film) 3193, 2947, 2872, 1674, 1734 cm⁻¹; ¹H NMR δ 6.54 (s, 1H), 5.72 (d, 1H, *J* = 1.2 Hz), 4.69 (m, 1H), 2.04 (s, 3H), 1.22 (s, 3H), 0.88 (s, 3H); ¹³C NMR δ 200.6, 173.1, 170.1, 163.5, 126.8, 72.5, 54.3, 49.5, 45.1, 41.8, 39.4, 38.8, 37.9, 36.2, 31.1, 27.6, 26.9, 23.1, 21.9, 21.6, 21.0, 17.2. (The data were in full accordance with the literature values.)

3-[(1,1-dimethylethyl)dimethylsilyl]oxy-7,17-Dioxo-17a-aza-D-homoandrost-5-ene (4b):

Isolated as a white solid; $[\alpha]_{\text{D}}^{20}$ –94.5 (*c* 0.20, CHCl₃); IR (thin film) 3197, 2929, 2857, 1688, 1673 cm⁻¹; ¹H NMR δ 6.11 (br s, 1H), 5.69 (s, 1H), 3.59 (s, 1H), 0.88 (s, 9H), 0.05 (s, 6H); ¹³C NMR δ 201.1, 172.9, 165.6, 125.9, 71.7, 54.3, 49.7, 45.1, 42.7, 41.9, 39.4, 39.1, 36.6, 32.0, 31.2, 30.1, 26.2, 23.2, 22.0, 21.1, 18.6, 17.3, –4.1; MS (ESI) *m/z* (relative intensity) 432 ([M + H]⁺, 100); HRMS (ESI) calcd for C₂₅H₄₁NO₃Si 432.2928 [M + H]⁺, found 432.2932. (The compound **4b** could be converted to **4a** by the literature procedure²⁰ and the data were in full accordance with those reported in the literature.¹⁶)

3-acetyloxy-(3 β ,25R)-spirost-5-en-7-one (6a)¹⁵: (72%, 24% recovery) Isolated as a white solid; mp 190–193 °C; $[\alpha]_{\text{D}}^{20}$ –146.8 (*c* 3.60, CHCl₃); IR (thin film) 2948, 2872, 1736, 1672 cm⁻¹; ¹H NMR δ 5.70 (s, 1H), 4.70 (m, 1H), 4.48 (m, 1H), 3.42 (m, 2H), 2.86 (m, 1H), 2.46 (m, 3H), 2.04 (s, 3H), 1.22 (s, 3H), 0.97 (d, 3H, *J* = 6.9 Hz), 0.79 (s, 3H), 0.78 (s, 3H); ¹³C NMR δ 201.7, 170.6, 164.4, 126.5, 109.5, 81.3, 72.5, 67.1, 61.4, 50.0, 49.8, 48.2, 41.9, 41.3, 39.0, 38.8, 38.1, 36.3, 34.0, 31.6, 29.1, 27.6, 21.6, 21.2, 17.6, 17.5, 16.8, 15.0. (The data were in full accordance with the literature values.)

3-[(1,1-dimethylethyl)dimethyl]-(3 β ,25R)-spirost-5-en-7-one (6b): (72%, 25% recovery) Isolated as a white solid; mp 236–238 °C; $[\alpha]_{\text{D}}^{20}$ –125.6 (*c* 0.80, CHCl₃); IR (thin film) 2934, 2860, 1674 cm^{–1}; ¹H NMR δ 5.66 (s, 1H), 4.48 (m, 1H), 3.57 (m, 1H), 3.45–3.38 (m, 2H), 2.86 (m, 1H), 2.41–2.37 (m, 1H), 1.99 (s, 3H), 0.97 (d, 3H, *J* = 7.5 Hz), 0.88 (s, 9H), 0.79 (s, 3H), 0.77 (s, 3H), 0.05 (s, 6H); ¹³C NMR δ 202.3, 166.5, 126.0, 109.6, 81.4, 71.7, 67.2, 61.5, 50.3, 49.9, 45.3, 43.0, 42.0, 41.4, 39.2, 38.9, 36.8, 34.1, 32.1, 31.9, 30.8, 29.2, 26.3, 21.4, 18.6, 17.7, 17.6, 16.9, 15.1, –4.1; MS (ESI) *m/z* (relative intensity) 565 ([M + Na]⁺, 100); HRMS (ESI) calcd for C₃₃H₅₄O₄Si 565.3684 [M + Na]⁺, found 565.3686. (The compound **6b** could be converted to **6a** by the literature procedure²⁰ and the data were in full accordance with those reported in the literature.¹⁵)

3 β ,28-Bis[(1,1-dimethylethyl)dimethylsilyl]oxyurs-12-ene (7): To a solution of uvaol (25 mg, 0.060 mmol) in CH₂Cl₂ (2 ml) was added NEt₃ (0.04 ml, 0.30 mmol) at 0 °C under nitrogen. TBSOTf (0.05 ml, 0.24 mmol) was added and the reaction mixture was stirred in 0 °C for 30 min. Saturated ammonium chloride solution was added and the resulting mixture was extracted by CH₂Cl₂ (2 ml $\times$ 3). The organic extracts were collected and concentrated under reduced pressure. Column chromatography of the residue using Et₂O:*n*-hexane = 2:98 afforded **7** (40 mg, 100%) as a colourless oil: $[\alpha]_{\text{D}}^{20}$ +37.6 (*c* 1.90, CHCl₃); IR (thin film) 2953, 2929, 2856, 1471 cm^{–1}; ¹H NMR δ 5.07 (m, 1H), 3.48 (d, 1H, *J* = 9.6 Hz), 3.18 (dd, 1H, *J* = 10.8, 4.5 Hz), 3.00 (d, 1H, *J* = 9.6 Hz), 1.61 (s, 3H), 0.03 — –0.02 (m, 12H); ¹³C NMR δ 139.5, 125.2, 80.0, 70.1, 55.7, 54.6, 48.2, 42.5, 40.5, 40.0, 39.9, 39.8, 39.3, 38.7, 37.2, 36.2, 33.4, 31.4, 30.2, 29.0, 28.1, 26.4, 26.1, 23.8, 23.7, 21.9, 19.0, 18.6, 17.8, 17.2, 16.6, 16.2, –2.4, –3.2, –4.4, –4.9; MS (EI) *m/z* (relative intensity) 613 ([M – C₄H₉]⁺, 12); HRMS (EI) calcd for C₄₂H₇₈O₂Si₂ 613.4831 [M – C₄H₉]⁺, found 613.4819.

3β,28-Bis[(1,1-dimethylethyl)dimethylsilyl]oxyurs-12-en-11-one (8): (78%, 24% recovery)

Isolated as a colourless oil; $[\alpha]_{\text{D}}^{20} +46.7$ (c 0.66, CHCl_3); IR (thin film) 2933, 2857, 1662 cm^{-1} ; ^1H NMR δ 5.47 (s, 1H), 3.38 (d, 1H, $J = 9.0$ Hz), 3.18 (dd, 1H, $J = 11.2, 4.5$ Hz), 3.00 (d, 1H, $J = 9.0$ Hz), 2.7 (m, 1H), 2.31 (s, 1H), 0.88 (s, 9H), 0.86 (s, 9H), 0.03 (s, 6H), -0.02 (s, 6H); ^{13}C NMR δ 200.3, 164.6, 130.9, 79.8, 70.2, 62.1, 55.4, 54.5, 45.5, 44.1, 40.1, 39.89, 39.7, 39.4, 33.3, 31.0, 28.9, 28.0, 27.0, 26.4, 26.3, 23.1, 21.6, 21.0, 18.8, 18.2, 17.8, 17.0, 16.5, -3.2 , -4.4 , -4.9 ; MS (ESI) m/z (relative intensity) 685 ($[\text{M} + \text{H}]^+$, 100); HRMS (ESI) calcd for $\text{C}_{42}\text{H}_{76}\text{O}_2\text{Si}_2$ 685.5406 $[\text{M} + \text{H}]^+$, found 685.5403. (The compound **8** could be converted to **3β, 28-dihydroxyurs-12-en-11-one (8a)**²³: To a solution of **8** (10 mg, 0.0144 mmol) and THF (0.4 ml) was added TBAF (0.2 ml, 0.2 mmol) and the reaction mixture was stirred in 60 °C for 6 h. The solution was concentrated and column chromatography of the residue using $\text{Et}_2\text{O}:\textit{n}$ -hexane = 1:99 afforded **8a** (4 mg, 64 %) as a white solid.)

3-methyl-2-cyclohexen-1-one (12)²⁴: Isolated as a yellow oil; IR (thin film) 2938, 1664 cm^{-1} ; ^1H NMR δ 5.84 (s, 1H), 2.33–2.23 (m, 4H), 1.96 (m, 2H), 1.92 (s, 3H); ^{13}C NMR δ 200.2, 163.3, 127.0, 37.3, 31.3, 24.8, 22.9. (The data were in full accordance with the literature values.)

3-Phenyl-2-cyclohexen-1-one (13)²⁶: Isolated as a yellow oil; IR (thin film) 1661, 1602 cm^{-1} ; ^1H NMR δ 7.53–7.49 (m, 2H), 7.41–7.31 (m, 3H), 6.40 (m, 1H), 2.75 (dt, 2H, $J = 5.8, 1.5$ Hz), 2.46 (t, 2H, $J = 6.3$ Hz), 2.13 (m, 2H); ^{13}C NMR δ 200.2, 160.1, 139.1, 130.3, 129.1, 126.4, 125.7, 37.6,

28.4, 23.1. (The data were in full accordance with the literature values.)

3-(2-Oxo-cyclohexanyl)-2-cyclohexene-1-one (14)²⁵: Isolated as a colourless oil; IR (thin film) 1710, 1682 cm^{-1} ; ^1H NMR δ 5.83 (s, 1H), 3.19 (dd, 1H, $J = 12.0, 5.0$ Hz), 2.43–2.32 (m, 1H), 2.30 (t, 2H, $J = 5.4$ Hz), 2.12–1.72 (m, 8H); ^{13}C NMR δ 208.6, 199.2, 163.2, 127.6, 59.4, 42.7, 37.9, 31.9, 29.3, 27.9, 25.1, 23.1. (The data were in full accordance with the literature values.)

(S)-(-)- α -Verbinone (18)²⁷: Isolated as a yellow oil; $[\alpha]_{\text{D}}^{20} -126.5$ (c 0.29, CHCl_3); IR (thin film) 2945, 1679 cm^{-1} ; ^1H NMR δ 5.72 (m, 1H); 2.79 (m, 1H); 2.63 (dt, 1H, $J = 5.7, 1.8$ Hz); 2.40 (t, 1H, $J = 5.1$ Hz); 2.07 (d, 1H, $J = 9.0$ Hz), 2.09 (d, 3H, $J = 1.5$ Hz), 1.48 (s, 3H), 1.00 (s, 3H); ^{13}C NMR δ 204.5, 170.6, 121.6, 58.0, 54.5, 50.1, 41.3, 27.0, 24.0, 22.5. (The data were in full accordance with the literature values.)

1,4-Dioxaspiro[4.5]dec-6-en-8-one (20)²⁸: Isolated as a colourless oil; IR (thin film) 2962, 2890, 1684 cm^{-1} ; ^1H NMR δ 6.56 (d, 1H, $J = 10.2$ Hz), 5.94 (dd, 1H, $J = 10.2, 3.0$ Hz), 3.98 (s, 4H), 2.55 (t, 1H, $J = 4.8$ Hz); ^{13}C NMR δ 199.4, 147.1, 131.1, 104.6, 65.7, 35.9, 33.5. (The data were in full accordance with the literature values.)

2,2,6-trimethyl-2-cyclohexen-1,4-dione (22)²⁹: (74%, 15% recovery) Isolated as a yellow oil; IR (thin film) 2970, 1683, 1626 cm^{-1} ; ^1H NMR δ 6.54 (d, 1H, $J = 0.6$ Hz), 2.69 (s, 2H), 1.98 (d, 3H, $J = 1.5$ Hz), 1.22 (s, 6H); ^{13}C NMR δ 203.8, 198.1, 149.3, 137.4, 52.2, 45.5, 26.5, 17.2. (The data were in full accordance with the literature values.)

3-methyl-2-cyclopenten-1-one (24)³⁰: Isolated as a colourless oil; mp 76–77 $^{\circ}\text{C}$; IR (thin film) 2918, 1701, 1621 cm^{-1} ; ^1H NMR δ 5.89 (s, 1H), 2.66 (m, 2H), 2.52 (m, 2H), 2.14 (s, 3H); ^{13}C NMR δ 210.6, 179.3, 131.0, 36.1, 33.4, 19.8. (The data were in full accordance with the literature

values.)

3,7,7-trimethylbicyclo[4.1.0]hept-3-ene-2,5-dione (26)³¹: Isolated as a yellow solid; mp 76–77 °C; $[\alpha]_{\text{D}}^{20} +32.1$ (*c* 0.25, CHCl₃); IR (thin film) 2931, 1660 cm⁻¹; ¹H NMR δ 6.48 (m, 1H), 2.31 (m, 2H), 1.96 (d, 3H, *J* = 1.8 Hz), 1.31 (s, 3H), 1.30 (s, 3H); ¹³C NMR δ 195.4, 194.8, 150.4, 138.0, 40.2, 39.4, 34.0, 29.5, 16.6, 15.8. (The data were in full accordance with the literature values.)

2-cyclohexenone (35): Isolated as a colourless oil; IR (thin film) 2944, 1679 cm⁻¹; ¹H NMR δ 6.97 (dt, 1H, *J* = 10.2, 3.9 Hz), 5.99 (m, 1H), 2.41 (t, 1H, *J* = 6 Hz), 2.35–2.29 (m, 1H), 1.99 (m, 1H); ¹³C NMR δ 200.2, 151.2, 130.3, 38.5, 26.0, 23.1. (The data were in full accordance with the literature values and a commercial available authentic sample.)

3-(*t*-butyl)-2-cyclohexene-1-one (37)²⁴: Isolated as a yellow oil; IR (thin film) 2960, 1670, 1612 cm⁻¹; ¹H NMR δ 5.94 (s, 1H), 2.35 (m, 4H), 1.96 (m, 2H), 1.11 (s, 9H); ¹³C NMR δ 201.3, 174.4, 123.4, 37.9, 28.7, 26.3, 23.7. (The data were in full accordance with the literature values.)

Table S1. Yield as a function of solvent in the allylic oxidation of cholesteryl acetate (**1a**) catalyzed by Manganese(III) acetate dihydrate.

Entry	Solvent ^a	Yield (%)	% Recovery
1	Et ₂ O	23	70
2	THF	0	94
3	EtOAc	87	0
4	CH ₂ Cl ₂	76	0
5	MeCN	77	0

^a Solvent concentration was 0.12 M [substrate] for all entries.

Table S2. Oxidation of benzylic carbons.

Entry	Substrate	Product	Time (h)	Isolated yield (%)
1			48	89
2			48	99
3			48	96

Table S3. More examples of allylic oxidation of simple alkenes.

Entry	Substrate	Product	Time (h)	Isolated yield (%)
1	34	35	48	30 ^a
2	36	37	48	76

^a High volatility of substrate and/or product resulted in material loss during the reaction and workup process, hence low reaction yield.

Figure S1. Isolated yield as a function of equivalent of TBHP in the allylic oxidation of cholesteryl acetate (**1a**) catalyzed by 10 mol% manganese(III) acetate dihydrate in ethyl acetate.

Figure S2. Isolated yield as a function of mole percentage of manganese(III) acetate dihydrate in the allylic oxidation of cholesteryl acetate (**1a**) with 5 eqv. TBHP in ethyl acetate.

Figure S3. EPR spectrum of DMPO-*t*BuOO spin adduct.^a

^a Experimental parameters: Microwave frequency = 9.731 GHz; Power = 0.399 mW; Modulate frequency = 100.00 kHz; Modulate Amplitude = 0.40 G. Hyperfine constants ($a^N = 14.0$ G, $a^H_\beta = 10.5$ G, $a^H_\gamma = 1.4$ G).³²

5,5-dimethyl-1-pyrroline-*N*-Oxide
(DMPO)

DMPO-*t*BuOO

Scheme S1. Evidence for the intermediate of *tert*-butyl peroxy ethers in the allylic oxidation of 1-phenylcyclohexene (**13**) catalyzed by manganese(III) acetate dihydrate.

Further information on allylic oxidation of simple alkenes.

Scheme 2 could rationalize the important role of oxygen, which enhance the possibility to generate the less stable radical from simple alkenes, as the ring is more flexible than that of rigid complex alkenes and hence less syn-parallel π -stabilization (Scheme S2).³³

Scheme S2. π -Stabilization of allylic radical for simple alkene.

The interesting cases (Entries 26, 27) could be explained by the “ π -like” σ -bond syn-parallel extra stabilization of both radicals generated adjacent to the cyclopropane moiety of (+)-3-carene (**27**).^{12,13} This could also rationalize the isomerization and oxidation of (+)-2-carene (**25**) to enedione **26** through a more radical stabilized transition state.

Scheme S3. π -Stabilization of allylic radical for carene.

Current Data Parameters

NAME	p145
EXPNO	1
PROCNO	1

F2 - Acquisition Parameters

Date_	20040908
Time	16.21
INSTRUM	dx300
PROBHD	5 mm BB0 BB-1H
PULPROG	zg
TD	32768
SOLVENT	CDCl3
NS	40
DS	0
SWH	8892.806 Hz
FIDRES	0.274439 Hz
AQ	1.8219508 sec
RG	64
DM	55.600 usec
DE	79.43 usec
TE	683.2 K
D1	1.00000000 sec
MCREST	0.00000000 sec
MCNMR	0.01500000 sec

===== CHANNEL f1 =====

NUC1	¹ H
P1	9.00 usec
PL1	-2.00 dB
SFO1	300.1312000 MHz

F2 - Processing parameters

SI	32768
SF	300.130060 MHz
WDW	EM
SSB	0
LB	0.30 Hz
GB	0
PC	1.00

1D NMR plot parameters

CX	22.00 cm
CY	10.87 cm
F1P	8.500 ppm
F1	2551.10 Hz
F2P	-0.500 ppm
F2	-150.07 Hz
PPHCH	0.40909 ppm/cm
HZCM	122.78046 Hz/cm

Current Data Parameters

NAME p162

EXPNO 1

PROCNO 1

F2 - Acquisition Parameters

Date_ 20041007

Time 10.02

INSTRUM dpx300

PROBHD 5 mm BBO BB-1H

PULPROG zg

TD 32768

SOLVENT CDCl3

NS 35

DS 0

SWH 8992.806 Hz

FIDRES 0.274439 Hz

AQ 1.8219508 sec

RG 64

DM 55.600 usec

DE 79.43 usec

TE 297.2 K

D1 1.00000000 sec

MCHEST 0.00000000 sec

MCMEK 0.01500000 sec

===== CHANNEL f1 =====

NUC1 1H

P1 5.00 usec

PL1 -2.00 dB

SFO1 300.1312000 MHz

F2 - Processing parameters

SI 32768

SF 300.1300063 MHz

WDW EM

SSB 0

LB 0.30 Hz

GB 0

PC 1.00

10 MHz plot parameters

CX 22.00 cm

CY 10.42 cm

F1P 8.500 ppm

F1 2551.10 Hz

F2P -0.500 ppm

F2 -150.07 Hz

PPMCH 0.40309 ppm/cm

HZCM 122.78046 Hz/cm

ppm

210.131

202.058

166.635

126.041

77.921

77.498

77.074

71.674

62.739

50.432

50.276

45.654

44.821

42.994

38.844

38.137

36.856

32.121

32.049

26.899

26.261

24.031

21.557

18.572

17.738

13.685

-4.211

-4.243

Current Data Parameters

NAME p162agatinc13

EXPNO 1

PROCNO 1

F2 - Acquisition Parameters

Date_ 20041222

Time 12.59

INSTRUM dpv300

PROBHD 5 mm BBO BB-1H

PULPROG zgpg

TD 65536

SOLVENT CDCl3

NS 261

DS 0

SWH 22675.736 Hz

FIDRES 0.346004 Hz

AQ 1.445188 sec

RG 8192

DM 22.050 uspc

DE 5.00 uspc

TE 0.0 K

D1 1.00000000 sec

d11 0.03000000 sec

MCREST 0.00000000 sec

MCNRM 0.01500000 sec

===== CHANNEL f1 =====

NUC1 13C

P1 3.00 usec

PL1 -6.00 dB

SFO1 75.474511 MHz

===== CHANNEL f2 =====

CPDPRG2 waltz16

NUC2 1H

PCPD2 100.00 usec

PL2 120.00 dB

PL12 19.00 dB

SFO2 300.1315007 MHz

F2 - Processing parameters

SF 65536

SF 75.4677157 MHz

MW 0

SSB 0

LB 3.00 Hz

GB 0

PC 1.40

===== ID NMR plot parameters =====

CX 23.00 cm

CY 8.39 cm

FAP 220.000 ppm

F1 16602.90 Hz

F2P -10.000 ppm

F2 -754.68 Hz

PCNRM 10.00000 ppm/ci

TCN 0.00000 ppm/ci

Current Data Parameters

NAME p1566again

EXPNO 1

PROCNO 1

F2 - Acquisition Parameters

Date_ 20050517

Time 15.13

INSTRUM gpc300

PROBHD 5 mm BB0 BB-1H

PULPROG zg

TD 32768

SOLVENT CDCl3

NS 20

DS 0

SWH 8992.806 Hz

FIDRES 0.274439 Hz

AQ 1.8219508 sec

RG 181

DW 55.600 usec

DE 79.43 usec

TE 296.2 K

D1 1.00000000 sec

MCREST 0.00000000 sec

MCNKR 0.01500000 sec

===== CHANNEL f1 =====

NUC1 1H

P1 5.00 usec

PL1 -2.00 dB

SFO1 300.1312000 MHz

F2 - Processing parameters

SI 32768

SF 300.1300066 MHz

WDW EM

SSB 0

LB 0.30 Hz

GB 0

PC 1.00

1D NMR plot parameters

CX 22.00 cm

CY 12.00 cm

F1P 8.500 ppm

F1 2551.10 Hz

F2P -0.500 ppm

F2 -150.07 Hz

PPMCH 0.40909 ppm/cm

HZCM 122.78046 Hz/cm

ppm

141.834

120.793

77.926
77.503
77.079
72.820

54.708
50.053
48.545
42.985
40.291
37.514
37.254
32.752
32.395
31.737
26.392
22.584
21.414
20.581
19.772
18.716
-4.125

Current Data Parameters

NAME p156c13

EXPNO 1

PROCNO 1

F2 - Acquisition Parameters

Date_ 20041216

Time 14.39

INSTRUM dpx300

PROBHD 5 mm BB-1H

PULPROG zgpg

TD 65536

SOLVENT CDCl3

NS 956

DS 0

SWH 22675.736 Hz

FIDRES 0.346004 Hz

AQ 1.4451188 sec

RG 8192

DW 22.050 usec

DE 6.00 usec

TE 0.0 K

D1 1.00000000 sec

d11 0.03000000 sec

MCREST 0.00000000 sec

MCIRK 0.01500000 sec

***** CHANNEL f1 *****

NUC1 13C

P1 3.00 usec

PL1 -6.00 dB

SF01 75.4745111 MHz

***** CHANNEL f2 *****

CPDPRG2 waltz16

NUC2 1H

PCPD2 100.00 usec

PL2 120.00 dB

PL12 19.00 dB

SF02 300.1315007 MHz

F2 - Processing parameters

SI 65536

SF 75.4677136 MHz

MDM EW

SSB 0

LB 3.00 Hz

GB 0

PC 1.40

F2 - Acquisition Parameters

CX 23.00 cm

CY 7.07 cm

FIP 220.000 ppm

F1 16602.50 Hz

F2p -10.000 ppm

F2 -754.68 Hz

BPMCH 10.00000 ppm/cm

HZCH 754.67712 Hz/cm

Current Data Parameters

NAME p168

EXPNO 1

PROCNO 1

F2 - Acquisition Parameters

Date_ 20041011

Time 21.11

INSTRUM dx300

PROBHD 5 mm BBO BB-1H

PULPROG zg

TD 32768

SOLVENT CDCl3

NS 91

DS 0

SWH 8992.806 Hz

FIDRES 0.27439 Hz

AQ 1.8219508 sec

RG 228.1

DM 55.600 usec

DE 79.43 usec

TE 297.2 K

D1 1.00000000 sec

WREST 0.00000000 sec

MCWK 0.01500000 sec

***** CHANNEL f1 *****

NUC1 1H

P1 5.00 usec

PL1 -2.00 dB

SFO1 300.1312000 MHz

F2 - Processing parameters

SF 32768

SF 300.1300065 MHz

WDW EM

SSB 0

LB 0.30 Hz

GB 0

PC 1.00

1D NMR Plot parameters

CX 22.00 cm

CY 7.95 cm

F1P 8.500 ppm

F1 2531.10 Hz

F2P -150.07 Hz

F2 0.40909 ppm/cm

PRNCH 122.78046 Hz/cm

H2CM

ppm

201.177

172.922

165.605

125.996

77.926

77.502

77.079

71.761

54.386

49.772

45.105

42.797

41.905

39.456

39.104

36.690

32.043

31.273

30.174

26.293

23.244

22.022

21.151

18.620

17.305

-4.189

4b

Current Data Parameters
NAME DISC13
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20041011
Time 22.53
INSTRUM dpx300
PROBHD 5 mm BBO BB-1H
PULPROG zgpg
TD 65536
SOLVENT CDCl3
NS 2297
DS 0
SWH 22675.736 Hz
FIDRES 0.346004 Hz
AQ 1.4451188 sec
RG 2048
DM 22.050 usec
DE 6.00 usec
TE 297.2 K
D1 1.00000000 sec
d11 0.03000000 sec
MCREST 0.00000000 sec
MCMRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 3.00 usec
PL1 -6.00 dB
SFO1 75.4745111 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 120.00 dB
PL12 19.00 dB
SFO2 300.1315007 MHz

F2 - Processing Parameters
SI 65536
SF 75.4677133 MHz
WDW EM
SSB 0
LB 3.00 Hz
GB 0
PC 1.40

10 NMR Plot Parameters
CX 23.00 cm
CY 8.10 cm
F1P 220.000 ppm
F1 16602.90 Hz
F2P -10.000 ppm
F2 -754.68 Hz
PPMCM 10.00000 ppm/cm
HZCM 754.67713 Hz/cm

Current Data Parameters

NAME	p172
EXPNO	1
PROCNO	1

F2 - Acquisition Parameters

Date_	20041019
Time	20.56
INSTRUM	dpX300
PROBHD	5 mm BBO BB-1H
PULPROG	zg
TD	32768
SOLVENT	CDCl3
NS	17
DS	0
SMH	8992.806 Hz
FIDRES	0.274439 Hz
AQ	1.8219508 sec
RG	161.3
OR	55.600 usec
DE	79.43 usec
TE	0.0 K
D1	1.00000000 sec
MCREST	0.00000000 sec
MCNMR	0.01500000 sec

***** CHANNEL f1 *****

NUC1	¹ H
P1	5.00 usec
PL1	-2.00 dB
SFO1	300.1312000 MHz

F2 - Processing parameters

SF	300.1300066 MHz
WDW	EM
SSB	0
LB	0.30 Hz
GB	0
PC	1.00

10 NMR plot parameters

CX	22.00 cm
CY	8.00 cm
FJP	8.500 ppm
F1	2551.10 Hz
F2P	-0.500 ppm
F2	-150.07 Hz
PPMCM	0.40909 ppm/cm
HZCM	122.78046 Hz/cm

Current Data Parameters

NAME	p155
EXPNO	1
PROCNO	1

F2 - Acquisition Parameters

Date_	20040930
Time	10.10
INSTRUM	dx300
PROBHD	5 mm BBO BB-1H
PULPROG	zg
TD	32768
SOLVENT	CDCl3
NS	64
DS	0
SWH	8992.806 Hz
FIDRES	0.27439 Hz
AQ	1.8219508 sec
RG	287.4
DN	55.500 usec
DE	79.43 usec
TE	297.2 K
D1	1.00000000 sec
MCREST	0.00000000 sec
MCNRC	0.01500000 sec

===== CHANNEL f1 =====

NUC1	¹ H
P1	5.00 usec
PL1	-2.00 dB
SFO1	300.1312000 MHz

F2 - Processing parameters

SI	32768
SF	300.130063 MHz
WGM	EM
SSB	0
LB	0.30 Hz
GB	0
PC	1.00

10 NMR plot parameters

CX	22.00 cm
CY	9.14 cm
FJP	8.500 ppm
F1	2551.10 Hz
F2P	-0.500 ppm
F2	-150.07 Hz
PPMCM	0.40909 ppm/cm
HZCM	122.78046 Hz/cm

ppm

139.515

125.250

Current Data Parameters
NAME p155c13
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters

Date_ 20041004

Time 10.02

INSTRUM gp300

PROBHD 5 mm BBO BB-1H

PULPROG zgpg

TD 65536

SOLVENT CDCl3

NS 1406

DS 0

SWH 22675.736 Hz

FIDRES 0.346004 Hz

AQ 1.445188 sec

RG 8192

DE 22.050 usec

TE 297.2 K

D1 1.00000000 sec

d11 0.03000000 sec

MCHEST 0.00000000 sec

MCMRK 0.01500000 sec

S31

F2 - Processing Parameters

SI 65536

SF 75.4677119 MHz

WDW EM

SSB 0

LB 3.00 Hz

GB 0

PC 1.40

CX 23.00 cm

CY 7.52 cm

F1P 220.000 ppm

F1 16602.90 Hz

F2P -10.000 ppm

F2 -754.68 Hz

PPMKM 10.00000 ppm/cm

H7M 754.57712 Hz/cm

Current Data Parameters
NAME p157
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20041011
Time 20.54
INSTRUM dpX300
PROBHD 5 mm BBO BB-1H
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 101
DS 0
SMH 8992.806 Hz
FIDRES 0.274439 Hz
AQ 1.8219508 sec
RG 574.7
DM 55.600 usec
DE 79.43 usec
TE 297.2 K
D1 1.00000000 sec
MCHES1 0.00000000 sec
MCMRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 5.00 usec
PL1 -2.00 dB
SFO1 300.1312000 MHz

F2 - Processing parameters
SI 32768
SF 300.1300065 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR Plot parameters
CX 22.00 cm
CY 5.47 cm
F1P 8.500 ppm
F1 2551.10 Hz
F2P -0.500 ppm
F2 -150.07 Hz
PPMCM 0.40909 ppm/cm
HZCM 122.78046 Hz/cm

References:

- (1) Zheng, Y.; Li, Y. *J. Org. Chem.* **2003**, *68*, 1603-1606.
- (2) Barton, D. H. R.; Clive, C.; Magnus, P. D. *J. Chem. Soc. Chem. Commun.* **1975**, 432-433.
- (3) Menger, F. M.; McCann, D. J. *J. Org. Chem.* **1985**, *50*, 3928-3930.
- (4) Yu, W.; Jin, Z. *J. Am. Chem. Soc.* **2001**, *123*, 3369-3370.
- (5) Drew, J.; Letellier, M.; Morand, P.; Szabo, A. G. *J. Org. Chem.* **1987**, *52*, 4047-4056.
- (6) Phillipou, G.; Bigham, D. A.; Seamark, R. F. *Steroids* **1975**, *26*, 516-524.
- (7) Akamanchi, K. G.; Patel, H. C.; Meenakshi, R. *Synth. Commun.* **1992**, *22*, 1655-1660.
- (8) Revelli, G. A.; Gros, E. G. *Steroids* **1993**, *58*, 181-184.
- (9) Suhr, R.; Lahmann, M.; Oscarson, S.; Thiem, J. *Eur. J. Org. Chem.* **2003**, *20*, 4003-4011.
- (10) Lahmaun, M.; Gyback, H.; Garegg, P. J.; Oscarson, S.; Suhr, R.; Thiem, J. *Carbohydr. Res.* **2002**, *337* (21-23), 2153-2159.
- (11) Tsunoda, T.; Suzuki, M.; Noyori, R. *Tetrahedron Lett.* **1980**, *21*, 1357-1358.
- (12) (a) Bartlett, P. D.; Günther, P. *J. Am. Chem. Soc.* **1966**, *88*, 3288-3289. (b) Bartlett, P. D.; Guaraldi, G. *J. Am. Chem. Soc.* **1967**, *89*, 4799-4800.
- (13) Catino, A. J.; Forslund, R. E.; Doyle, M. P. *J. Am. Chem. Soc.* **2004**, *126*, 13622-13623.
- (14) Arasasingham, R. D.; Cornman, C. R.; Balch, A. L. *J. Am. Chem. Soc.* **1989**, *111*, 7800-7805.
- (15) Pearson, A. J.; Chen, Y. S.; Han, G. R.; Hsu, S. Y.; Ray, T. *J. Chem. Soc. Perkin Trans. I* **1985**, 267-273.
- (16) Arsenou, E. S.; Koutsourea, A. I.; Foustieris, M. A.; Nikolaropoulos, S. S. *Steroids* **2003**, *68*, 407-414.
- (17) Bauduin, G.; Bondon, D.; Pietrasanta, Y.; Pucci, B. *Tetrahedron* **1978**, *34*, 3269-3274.
- (18) Ciuffreda, P.; Casati, S.; Bollini, D.; Santaniello, E. *Steroids* **2003**, *68*, 193-198.
- (19) Marwah, P.; Marwah, A.; Kneer, N.; Lardy, H. *Steroids* **2001**, *66*, 581-595.
- (20) The 3 β -OTBS was converted into the known 3 β -OAc under classical procedures. (a) Corey, E. J.; Venkateswarlu, A. *J. Am. Chem. Soc.* **1972**, *94*, 6190-6191. (b) Weber, H.; Khorana, H. G. *J. Mol. Biol.* **1972**, *72*, 219-249.

- (21) Zheng, Y.; Li, Y. *J. Org. Chem.* **2003**, *68*, 1603-1606.
- (22) Salvador, J. A. R.; Sa é Melo, M. L.; Compos Neves, A. S. *Tetrahedron Lett.* **1997**, *38*, 119-122.
- (23) Keiichi, N.; Toshio, M.; Hiroshi, N. Jpn. Kokai Tokkyo Koho 2001, 11pp. coden: JKXXAF JP 2001048828 A.
- (24) Wada, E.; Funakoshi, J.; Kanemasa, S. *Bull. Chem. Soc. Jpn.* **1992**, *65*, 2456-2464.
- (25) Magnus, P.; Lacour, J.; Evans, A.; Rigollier, P.; Tobler, H. *J. Am. Chem. Soc.* **1998**, *120*, 12486-12499.
- (26) Yu, J. Q.; Corey, E. J. *Org. Lett.* **2002**, *4*, 2727-2730.
- (27) Watanabe, M.; Zwen, B. Z.; Kato, M. *J. Org. Chem.* **1993**, *58*, 3923-3927.
- (28) Satoh, T.; Kawashima, T.; Takahashi, S.; Sakai, K. *Tetrahedron* **2003**, *59*, 9599-9607.
- (29) Carreno, M. C.; Perez-Gonzalez, M.; Ribagorda, M.; Somoza, A.; Urbano, A. *Chem. Commun.* **2002**, 3052-3053.
- (30) Hendrickson, J. B.; Palumbo, P. S. *J. Org. Chem.* **1985**, *50*, 2110-2112.
- (31) Rotheberg, G.; Yatziv, Y.; Sasson, Y. *Tetrahedron* **1998**, *54*, 593-598.
- (32) (a) Janzen, E. G.; Haire, D. L. *Advances in Free Radical Chemistry, Vol. 1* (Ed.: D. D. Tanner), Jai Press Inc., **1990**, pp. 253-295. (b) Chamulitrat, W.; Takahashi, N.; Mason, R. P. *J. Biol. Chem.* **1989**, *264*, 7889-7899.
- (33) Cainelli, G.; Cardillo, G. *Chromium Oxidations in Organic Chemistry*, (Ed.: G. Cainelli, G. Cardillo), Springer-Verlag, **1984**, pp 41-58.