Mild Organocatalytic α -Methylenation of Aldehydes

Anniina Erkkilä and Petri M. Pihko*

*Laboratory of Organic Chemistry, Helsinki University of Technology, P.O.B. 6100,
FI-02015 TKK, Finland*

Supporting Information

Table of Contents

Table of Contents	S1
General Methods	S2
Procedures for the Preparation of Starting Materials	S2
Synthesis of Catalyst 13 (L-Pro- β -Ala).....	S4
α -Methylenation of Aldehydes, General Procedures and Characterization Data.....	S5
Copies of ^1H and ^{13}C Spectra	S8

General Methods

All aqueous reactions were carried out under ambient atmosphere in capped vials, unless otherwise noted. Under non-aqueous conditions, the reactions were performed under argon, and reagents were dried prior to use and transferred under argon *via* syringe. THF was distilled from Na/benzophenone. DMSO was distilled from CaH₂. MeOH was distilled from Mg(OMe)₂. CH₂Cl₂ was distilled from CaH₂. *N*-Methyl morpholine was distilled over sodium chips. Other solvents and reagents were used as obtained from the supplier, unless otherwise noted. Temperatures refer to bath temperatures. Analytical TLC was performed using Merck silica gel F254 (230-400 mesh) plates and analyzed by UV light or by staining upon heating with vanillin solution (6 g vanillin, 5 mL conc. H₂SO₄, 3 mL glacial acetic acid, 250 mL EtOH) or ninhydrin solution (1 g ninhydrin, 100 mL isopropanol, 5 drops glacial acetic acid). For silica gel chromatography, the flash chromatography technique was used, with Merck silica gel 60 (230-400 mesh) and p.a. grade solvents unless otherwise noted.

The ¹H NMR and ¹³C NMR spectra were recorded in either CDCl₃ or H₂O/D₂O on a 400 MHz (¹H 399.98 MHz; ¹³C 100.59 MHz) spectrometer. The chemical shifts are reported in ppm relative to residual CHCl₃ (δ 7.26) or TMS (δ 0.00) signals for ¹H NMR spectra. The coupling constants are reported in Hz. For the ¹³C NMR spectra, the residual CDCl₃ (δ 77.0) was used as the internal standard.

The enantiomeric excess (ee) of **15** was determined by GC in comparison to the corresponding racemic samples using Chiraldex β-PH cyclodextrin column (30 m x 0.25 mm, 110°C isotherm, carrier pressure 32 psi, FID detection (280°C)).

Procedures for the Preparation of Starting Materials

Aldehyde 28 was obtained by Swern oxidation of the commercially available alcohol **S1**. Data obtained correlates with literature data.¹ ¹H NMR (CDCl₃, 400 MHz): δ (CDCl₃, 400 MHz): δ 9.80 (dd, 1H, *J* = 1.2, 2.0); 4.55 (tt, 1H, *J* = 6.0, 6.8 Hz); 4.18 (dd, 1H, *J* = 6.0; 8.4); 3.58 (dd, 1H, *J* = 6.8, 8.4 Hz); 2.85 (ddd, 1H, *J* = 2.0, 6.8, 17.2 Hz); 2.65 (ddd, 1H, *J* = 1.2, 6.0, 17.2 Hz) 1.41 (s, 3H); 1.36 (s, 3H); ¹³C NMR (CDCl₃, 100 MHz): δ 200.0, 109.2, 70.6, 69.1, 47.8, 26.8, 25.4; [α]_D²⁰ +6.1 (*c* = 2.0, CHCl₃).

Alcohol S3. To a solution of **S2** (3.5 mL, 3.56 g, 39.5 mmol, 100 mol%) in THF (60 mL) at -78°C was added dropwise BuLi (1.58 M, 25.0 mL, 39.5 mmol, 100 mol%), followed by TBDPSCl (10.1 mL, 10.85 g, 39.5 mmol, 100 mol%). The reaction mixture was allowed to warm to rt during 25 min., then brought to reflux for 4 h. The reaction mixture was allowed to cool to rt, then quenched with 7% aq NaHCO₃ solution (60 mL). The mixture was extracted with Et₂O (2 x 30 mL). The combined extracts were dried (Na₂SO₄) and concentrated in vacuo. The crude product was purified with flash chromatography (50% EtOAc in hexanes) to yield 9.40 g (73%) of alcohol **S3** as yellowish oil. Data obtained correlates with literature data.² ¹H NMR (CDCl₃, 400 MHz): δ 7.70-7.65 (m, 4H); 7.47-7.35 (m, 6H); 3.70 (t, 2H, *J* = 5.7 Hz); 3.66 (t, 2H, *J* = 5.7 Hz); 2.03 (br s, 1H); 1.74-1.60 (m, 4H), 1.06 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz): δ 135.5, 133.6, 129.6, 127.6, 64.0, 62.8, 29.8, 29.4, 26.8, 16.1.

¹ Arroyo-Gómez, Y.; López-Sastre, J. A.; Rodríguez-Amo, J. F.; Santos-García, M.; Sanz-Tejedor M. A. *Tetrahedron Asymmetry* **1999**, *10*, 973-990.

² Nacro, K.; Baltas, M.; Gorrichon, L. *Tetrahedron*, **1999**, *55*, 14013-14030.

Aldehyde 24. A solution of oxalyl chloride (1.9 mL, 2.76 g, 21.8 mmol, 110 mol%) in CH_2Cl_2 (50 mL) at -60°C , was added DMSO (3.1 mL, 3.41 g, 43.6 mmol, 220 mol%). After 30 min, a solution of alcohol **S3** (6.50 g, 19.8 mmol, 100 mol%) in CH_2Cl_2 (20 mL) was added and the mixture was stirred for 5 min. Et_3N was then added dropwise. The reaction mixture was allowed to warm to rt and stirred for 70 min. H_2O (100 mL) was added and the separated aqueous phase was extracted with CH_2Cl_2 (50 mL). The combined organic phases were washed with brine, dried (Na_2SO_4) and filtered through thin silica pad. The solution was concentrated in vacuo to afford 5.05 g (78%) of aldehyde **24** as yellowish oil. Data obtained correlates with literature data.³ ^1H NMR (CDCl_3 , 400 MHz): δ 9.81 (t, 1H, $J = 1.6$ Hz); 7.86-7.64 (m, 4H), 7.46-7.36 (m, 6H); 3.70 (t, 2H, $J = 6.0$ Hz); 2.55 (td, 2H, $J = 1.6$ Hz, 7.1 Hz); 1.90 (tt, 2H, $J = 6.0$ Hz, 7.1 Hz); 1.06 (s, 9H) ^{13}C NMR (CDCl_3 , 100 MHz): δ 202.5, 135.5, 133.5, 129.6, 127.6, 62.9, 40.7, 26.8, 25.2, 19.1.

Keto Aldehyde 26. A solution of methyl cyclohexene (1.2 mL, 0.96 g, 10.0 mmol, 100 mol%) in CH_2Cl_2 (70 mL) was cooled to -78°C . A stream of O_3 was bubbled through the solution until a blue colour persisted. A stream of O_2 was then bubbled through the reaction mixture to remove excess ozone. Dimethyl sulfide (1.2 mL, 0.99 g, 16.0 mmol, 160 mol%) was then added. The reaction mixture was stirred for an additional 23 h. H_2O was then added and the mixture was extracted with DCM (2 x 50 mL). The combined extracts were washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The crude product was purified with flash chromatography (40% MTBE in hexanes) to yield 0.38 g (30%) of **26** as pale yellow oil. IR 3419, 2943, 2726, 1712, 1411, 1365, 1161, 1086, 1000, 734 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 9.73 (t, 1H, $J = 1.2$ Hz); 2.45 (m, 4H), 2.14 (s, 3H); 1.61 (m, 4H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 208.4, 202.2, 43.7, 43.2, 29.9, 23.1, 21.5. HRMS (ESI⁺) calc for $[\text{C}_7\text{H}_{12}\text{O}_2 + \text{Na}]$ 151.0735, obsd 151.0742.

³ Freeman, F.; Kim, D. S. H. L. *J. Org. Chem.* **1992**, 57, 1722-1727.

Synthesis of Catalyst 13 (L-Pro-β-Ala)

Cbz-L-prolyl-β-alanine methyl ester S6. To Cbz-L-proline (**S4**) (3.00 g, 12.0 mmol, 100 mol%) in dry THF (120 mL) at $-25\text{ }^{\circ}\text{C}$, *N*-methyl morpholine (NMM) (1.4 mL, 12.6 mmol, 105 mol%) and isobutyl chloroformate (1.7 mL, 12.6 mmol, 105 mol%) were added with stirring. A cloudy mixture was formed. After 10 min, **S5** (1.76 g, 12.6 mmol, 105 mol%) was added, followed by NMM (1.5 mL, 13.2 mmol, 110 mol%). The mixture was allowed to stir for additional 2 h at $-25\text{ }^{\circ}\text{C}$, then quenched with sat. NaHCO_3 (180 mL) and extracted with EtOAc (3 x 100 mL). The combined organic extracts were dried (Na_2SO_4), filtered and concentrated. The crude product was recrystallized from Et₂O/hexanes to afford 4.00 g (quant.) of the dipeptide **S6** as white amorphous crystals. R_f 0.93 (50% MeOH in CH_2Cl_2) $[\alpha]_D^{20} = -71.8$ ($c = 1.0$ in CHCl_3); mp. 55-56 $^{\circ}\text{C}$, IR 3320, 3066, 2953, 2882, 1736, 1704, 1538, 1416, 1356, 1176, 1120, 752, 698; ^1H NMR (CDCl_3 , 400 MHz) δ 7.40-7.27 (m, 5H); 7.02 (br s, 0.5H); 6.49 (br s, 0.5H); 5.19-5.06 (m, 2H); 4.33-4.24 (m, 1H); 3.69-3.36 (m, 7H); 2.56-1.82 (m, 7H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 191.9, 172.6, 151.3, 136.3, 128.5, 128.1, 127.8, 67.2, 60.7, 51.7, 46.9, 34.8, 33.6, 28.6; HRMS (ESI+) calc for $[\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_5 + \text{Na}]$ 357.1426, obsd 357.1422; Anal. calc. for $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_5$: C, 61.07; H, 6.63; N, 8.38. Found: C, 61.00; H, 6.49; N, 8.58.

Cbz-L-prolyl-β-alanine S7. To the methyl ester **S6** (4.25 g, 12.7 mmol, 100 mol%) in 1:1 MeOH/THF (50 mL), was added 1 M NaOH (19.0 mL, 19.0 mmol, 150 mol%) at ambient temperature. The reaction mixture was stirred 3 h, then solvents were evaporated to afford 3.22 g (79%) of **S7** as white crystals. Spectroscopic data obtained for this compound correlates with literature data.⁴ R_f 0.68 (50% MeOH in DCM) $[\alpha]_D^{20} = -31.4^{\circ}$ ($c = 1.0$ in DMF); mp. 138-139 $^{\circ}\text{C}$.

L-prolyl-β-alanine 13. To **S7** (2.73 g, 8.5 mmol, 100 mol%) in MeOH (50 mL) under Ar was added 5% Pd/C (0.27 g, 10 w-%). The reaction mixture was then purged with argon and finally with hydrogen, and the mixture was stirred under H_2 for 1.5 h. After an argon purge, the reaction mixture was filtered through a thin celite pad and washed with MeOH to remove Pd/C. Evaporation of solvents afforded 1.32 g (83%) of **13** as white powder. R_f 0.05 (50% MeOH in CH_2Cl_2). Data obtained correlates with literature data.⁴ mp. 206-207 $^{\circ}\text{C}$; $[\alpha]_D^{20} = -50.8^{\circ}$ ($c = 1.0$ in H_2O); ^1H NMR (H_2O , 400 MHz): δ 4.34-4.30 (m, 1H), 3.49-3.34 (m, 4H), 2.43-2.06 (m, 3H), 2.10-2.98 (m, 3H) ^{13}C NMR (H_2O , 100 MHz): δ 179.9, 169.2, 59.7, 46.3, 36.9, 36.4, 29.6, 23.7.

⁴ Ienaga, K; Higashiura, K.; Kimura, H. *Chem. Pharm. Bull.* **1987**, *35*, 1249-1254.

α -Methylenation of Aldehydes, General Procedures and Characterization Data

General procedure A for the α -methylenation of aldehydes. To a mixture of aqueous formaldehyde solution (37% formaldehyde in water, 1.0 mmol, 100 mol%) and aldehyde (1.0 mmol) in *i*-PrOH (100 μ L) was added propionic acid (0.1 mmol, 10 mol%) and pyrrolidine (0.1 mmol, 10 mol%). The reaction mixture was stirred at 45 °C for 1-25 h. NaHCO₃ was added and the mixture was extracted with CH₂Cl₂ (3 x 5 mL). The combined extracts were washed with brine, dried (Na₂SO₄), and concentrated *in vacuo*. If necessary, the crude product thus obtained was purified by a passage through a short pad of silica gel using Et₂O as the eluent.

General procedure B for the α -methylenation of aldehydes. To catalyst **13** (L-Pro- β -Ala) (0.1 mmol, 10 mol%) in *i*-PrOH (100 μ L) in a 5 mL vial was added aqueous formaldehyde solution (37% formaldehyde in water, 1.0 mmol, 100 mol%) and aldehyde (1.0 mmol, 100 mol%) at rt. The mixture was stirred at 45 °C for 1-25 h. H₂O (5 mL) was then added and the mixture was extracted with CH₂Cl₂ (3 x 5 mL). The combined extracts were washed with brine, dried (Na₂SO₄), and concentrated *in vacuo*. If necessary, the crude product thus obtained was purified by a passage through a short pad of silica gel using Et₂O as the eluent.

Characterization Data for the α -Methylenation Products:

2-methyleneoctanal (17). The reaction was carried out following the General Procedure A to yield 129 mg (92 %) of **17** as a pale yellow oil. Data obtained match the literature data.⁵ ¹H NMR (CDCl₃, 400 MHz): δ 9.48 (s, 1H), 6.19 (s, 1H), 5.93 (s, 1H); 2.18 (t, 2H, *J* = 7.6 Hz), 1.2-1.5 (br m, 8 H), 0.87 (t, 3 H, *J* = 7 Hz); ¹³C NMR (CDCl₃, 100 MHz): δ 194.6, 150.4, 133.7, 31.5, 28.8, 27.7, 27.6, 22.4, 13.9.

2-tert-butylacrolein/3,3-dimethyl-2-methylenebutanal (19). The reaction was carried out following the General Procedure A to yield 97 mg (86%) of **19** as a colorless oil. IR 3380, 2961, 2872, 1702, 1460, 1364, 1306, 1195, 953, 848, 670; ¹H NMR (CDCl₃, 400 MHz): δ 9.54 (s, 1H); 6.30 (s, 1H); 5.90 (s, 1H); 1.19 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz): δ 195.0, 158.1, 133.6, 33.8, 28.5. HRMS (ESI+) calc for [C₇H₁₃O] 113.0966, obsd 133.0931.

(S)-3,7-Dimethyl-2-methylene-6-octanal (15). The reaction was carried out following the General Procedure A to yield 147 mg (89%) of **15** as a pale yellow oil. Data obtained correlates with literature data.⁶ $[\alpha]_D^{20} = +5.2$ (*c* = 13.4 in CHCl₃); ¹H NMR (CDCl₃, 400 MHz): δ 9.53 (s, 1H); 6.23 (s, 1H); 5.98 (s, 1H); 5.07 (tt, 1H, *J* = 1.2, 7.2 Hz); 2.70 (sext., 1H, *J* = 6.8 Hz); 1.92 (m, 2H); 1.67 (s, 3H); 1.56 (s, 3H); 1.51 (m, 1H); 1.37 (m, 1H) 1.06 (d, 3H, *J* = 6.8 Hz) ¹³C NMR (CDCl₃, 100 MHz): δ 194.7; 155.5; 133.2; 124.1; 35.6; 30.9; 25.7; 25.6; 19.5; 17.6. GC analysis (β -PH cyclodextrin, flow rate 32 psi, 110°C isotherm) *t_R* = 5.03 (minor), 5.13 (major) reveals this material to be 99% ee.

⁵ Fearnley, S. P.; Funk, R. L.; Gregg, R. J. *Tetrahedron* **2000**, 56, 10275-10282.

⁶ White, J. D.; Amedio, J. C.; Gut, S.; Ohira, S. Jayasinghe, L. R. *J. Org. Chem.* **1992**, 57, 2270-2284.

2-benzylacrolein/2-methylene-3-phenylpropanal (21). The reaction was carried out following the General Procedure A or General Procedure B to yield 146 mg (99%) of **21** as a pale yellow oil. Data obtained correlates with literature data.⁷ ¹H NMR (CDCl₃, 400 MHz): δ 9.61 (s, 1H); 7.32-7.17 (m, 5H); 6.11 (s, 1H); 6.07 (s, 1H); 3.57 (s, 2H); ¹³C NMR (CDCl₃, 100 MHz): δ 193.9, 149.7, 138.1, 135.2, 129.1, 128.5, 126.4, 34.1.

α-Phenylethylacrolein / 2-(1-phenyl-ethyl)-propenal (23). The reaction was carried out following either the General Procedure A to yield 144 mg (90%) or General Procedure B to provide 122 mg (76%) of **23** as a pale yellow oil. IR: 2986, 2305, 1693, 1421, 1265, 950, 896, 741, 705, 409; ¹H NMR (CDCl₃, 400 MHz): δ 9.54 (s, 1H); δ 7.33-7.16 (m, 5H); 6.24 (s, 1H); 6.08 (s, 1H); 4.06 (q, 1H, *J* = 7.2 Hz); 1.43 (d, 3H, *J* = 7.2 Hz); ¹³C NMR (CDCl₃, 100 MHz): δ 193.8; 154.4; 143.6; 133.6; 128.4; 127.5; 126.4; 37.2; 20.0. HRMS (ESI+) calc for [C₁₁H₁₂O + Na] 183.0786, obsd 183.0765.

2-(2'-(tert-Butyldiphenylsilyloxy)ethyl)propenal (25). The reaction was carried out following the General Procedure A to yield 284 mg (83%) as a pale yellow oil. Data obtained correlates with literature data.⁸ ¹H NMR (CDCl₃, 400 MHz): δ 9.49 (s, 1H); δ 7.67-7.62 (m, 4H); 7.42-7.26 (m, 6H); 6.35 (s, 1H); 6.05 (s, 1H); 3.77 (t, 2H, *J* = 6.4 Hz); 2.51 (t, 2H, *J* = 6.4 Hz); 1.03 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz): δ 194.4, 147.0, 135.8, 135.5, 133.7, 129.6, 127.6, 61.7, 31.0, 26.8, 19.2.

2-methylene-6-oxo-heptanal (27). The reaction was carried out following the General Procedure A to yield 99 mg (74%) of **27** as a pale yellow oil. IR 3412, 2943, 1713, 1691, 1411, 1367, 1143, 953 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 9.49 (s, 1H); 6.26 (s, 1H); 6.00 (s, 1H); 2.42 (t, 2H, *J* = 7.2 Hz); 2.20 (t, 2H, *J* = 7.2 Hz); 2.10 (s, 3H); 1.70 (qn, *J* = 7.2 Hz); ¹³C NMR (CDCl₃, 100 MHz): δ 208.5, 194.5, 149.4, 134.5, 42.8, 29.9, 26.9, 21.7; HRMS (ESI+) calc for [C₈H₁₂O₂ + Na] 163.0735, obsd 163.0731.

(S)-2-(2,2-dimethyl-[1,3]dioxolan-4-yl)-propenal (29). The reaction was carried out following the General Procedure B to yield 126 mg (80%) of **29** as a colorless oil. [α]_D²⁰ +21.0° (*c* = 1.0, CHCl₃); IR 3435, 2987, 1687, 1372, 1217, 1155, 1067, 852 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 9.57 (s, 1H); 6.59 (dd, 1H, *J* = 1.4 Hz); 6.11 (dd, 1H, *J* = 2 x 1.4 Hz); 4.83 (dddd, 1H, *J* = 2 x 1.4, 2 x 6.8 Hz); 4.34 (dd, 1H, *J* = 6.8, 8.4 Hz); 3.49 (dd, 1H, *J* = 6.8, 8.4 Hz); 1.42 (s, 3H); 1.39 (s, 3H); ¹³C NMR (CDCl₃, 100 MHz): δ 193.3, 149.0, 133.2, 109.2, 72.5, 69.5, 26.2, 25.3; HRMS (ESI+) calc for [C₈H₁₂O₃ + Na] 179.0684, obsd 179.0679.

⁷ Nakahira, H.; Ryu, I.; Ikebe, M.; Oku, Y.; Ogawa, A.; Kambe, N.; Sonoda, N.; Murai, S. *J. Org. Chem.* **1992**, 57, 17-28.

⁸ Myers, A. G.; Dragovich, P. S.; Kuox, E. Y. *J. Am. Chem. Soc.* **1992**, 114, 9369 - 9386.

3-benzyloxy-2-methylenepropanal (31). The reaction was carried out following the general General Procedure B to yield 156 mg (89%) of **31** as a pale yellow oil. IR 3436, 3088, 3064, 3031, 2863, 1954, 1693, 1496, 1454, 1368, 1101, 960, 739 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 9.60 (s, 1H); 7.38-7.28 (m, 5H); 6.58 (s, 1H); 6.16 (d, 1H, $J = 1.6$ Hz); 4.59 (s, 2H), 4.26 (t, 2H, $J = 1.6$ Hz); ^{13}C NMR (CDCl_3 , 100 MHz): δ 193.3, 146.7, 137.8, 134.0, 128.4, 127.7, 127.6, 73.0, 65.8; HRMS (ESI+) calc for $[\text{C}_{11}\text{H}_{12}\text{O}_2 + \text{Na}]$ 199.0735, obsd 199.0732.

2-Benzyloxyprop-2-enal/2-benzyloxy-2-methylene-ethanal (33). The reaction was carried out following the General Procedure A to yield 161 mg (98%) of **33** or General Procedure B to yield 103 mg (64%) of **33** as a pale yellow oil. Data obtained correlates with literature data.⁹ ^1H NMR (CDCl_3 , 400 MHz): δ 9.30 (s, 1H); 7.38-7.30 (m, 5H); 5.25 (d, 1H, $J = 3.0$); 5.14 (d, 1H, $J = 3.0$); 4.93 (s, 2H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 188.0, 158.1, 135.5, 128.6, 128.1, 127.3, 103.8, 70.1.

Tert-Butyl-1-formylvinylcarbamate / N-Boc-dehydroalaninal (35): The reaction was carried out following the General Procedure A to yield 146 mg (85%) of **35** or General Procedure B to yield 147 mg (86%) of **35** as a pale yellow oil. Data obtained correlates with literature data:¹⁰ ^1H NMR (CDCl_3 , 400 MHz): δ 9.12 (s, 1H); 6.97 (s, 1H); 6.70 (s, 1H); 5.35 (s, 1H); 1.46 (s, 9H) ^{13}C NMR (CDCl_3 , 100 MHz): δ 188.6, 152.3, 140.2, 114.8, 81.0, 28.1.

⁹ Hon, Y.-S.; Chang, F.-J.; Lu, L.; Lin, W.-C. *Tetrahedron* **1998**, 54, 5233-5246.

¹⁰ *Chem. Pharm. Bull.*, **1996**, 44, 2205-2212.

Copies of ^1H and ^{13}C Spectra

S9

S3

S10

24

S11

S12

S13

S14

S15

S16

S17

CC(=O)C(=C)CCOTBDPS

S19

S20

S21

S22

S23