Mimicking Dihydroxy Acetone Phosphate Utilizing Aldolases Through Organocatalysis: A Facile Route to Carbohydrates and Aminosugars.

Jeff T. Suri, Dhevalapally B. Ramachary, and Carlos F. Barbas III*

Contribution from The Skaggs Institute for Chemical Biology and the Departments of Chemistry and Molecular Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California

Supporting Information

General. Chemicals and solvents were either purchased *puriss p.A.* from commercial suppliers or purified by standard techniques. For thin-layer chromatography (TLC), silica gel plates Merck 60 F254 were used and compounds were visualized by irradiation with UV light and/or by treatment with a solution of *p*-anisaldehyde (23 mL), conc. H₂SO₄ (35 mL), acetic acid (10 mL), and ethanol (900 mL) followed by heating; or with a solution of ninhydrin in EtOH followed by heating. Flash chromatography was performed using silica gel Merck 60 (particle size 0.040-0.063 mm), ¹H NMR and ¹³C NMR spectra were recorded on a Bruker DRX-500 MHz instrument and were referenced internally to the residual solvent peak. HPLC was carried out using a Hitachi organizer consisting of a D-2500 Chromato-Integrator, a L-4000 UV-Detector, and a L-6200A Intelligent Pump. Optical rotations were recorded on a Perkin Elemer 241 Polarimeter (λ=589 nm, 1 dm cell). High-resolution mass spectra were recorded on an IonSpec TOF mass spectrometer. 1,3-Bis(triisopropylsilyloxy)propan-2-one (Table 1, entry 3) and 1,5-Dioxaspiro[5.5]undecan-3-one (Table 1, entries 8,9) were prepared according to the literature procedure.¹ 1,3-Dioxan-5-one (Table 1, entries 4,5) was prepared by PCC oxidation of 1,3-dioxan-5-ol in CH₂Cl₂.

General experimental procedure for the aldol reaction (Table 1): To a glass vial charged with DMF (200 μ L) was added ketone (0.5 mmols), aldehyde (0.1 mmols) and *S*-proline (0.02 mmols, 2.3 mg) the reaction was stirred at ambient temperature or at 4 $^{\circ}$ C for the appropriate time until the reaction was complete by TLC. Then, a half saturated NH_4Cl solution and ethyl acetate were added with vigorous stirring, the layers were separated and the organic phase was washed with brine. The organic phase was dried (MgSO_4), concentrated, and purified by flash column chromatography (silica gel, mixtures of hexanes/ethyl acetate) to afford the desired Aldol product.

General procedure for derivatization of aliphatic substrates for HPLC analysis. The aldol adduct (1 equiv) in CH_2Cl_2 (1 mL/0.5 mmols) was treated with 3,5-dinitrobenzoyl chloride (1.1 equiv) and DMAP (1.1 equiv) and stirred for 1 h. The solution was filtered through a small plug of silica gel and analyzed by HPLC.

(S)-4-((S)-Hydroxy(4-nitrophenyl)methyl)-1,3-dioxan-5-one (Table 1, entry 4):

^1H NMR (CDCl_3 , 500 MHz), major product: δ 8.20 (d, J = 8.7 Hz, 2H), 7.58 (d, J = 8.5 Hz, 2H), 5.07 (d, J = 5.9 Hz, 1H), 4.90 (d, J = 6, Hz 1H), 4.30 (d, J = 2.5 Hz, 1H), 3.76 (bs, 1H), 2.94 (s, 1H), 2.87 (s, 1H). ^{13}C NMR (CDCl_3 , 125 MHz) δ 205.75, 162.61, 145.8, 128.1, 123.26, 91.5, 93.4, 72.99, 72.44. HRMS for $\text{C}_{11}\text{H}_{10}\text{NO}_6$ $[\text{M}-\text{H}]^-$: calcd 252.0514, obsd 252.0511; HPLC (Daicel Chiralcel OJ-H, hexane/isopropanol = 80 : 20, flow rate 1.0 mL/min, λ = 254 nm): t_R = 31.17 min (*anti*, major), t_R = 33.92 min (*anti*, minor), t_R = 36.11 min (*syn*). $[\alpha]_D = -18.42$ (c = 0.73, CHCl_3).

(S)-4-((S)-Hydroxy(4-nitrophenyl)methyl)-2,2-dimethyl-1,3-dioxan-5-one (Table 1, entry

6): Column chromatography provided a separable diastereomeric mixture. ^1H NMR (CDCl_3 , 500 MHz), major product: δ 8.20 (d, J = 8.8 Hz, 2H), 7.59 (d, J = 8.6 Hz, 2H), 4.99 (d, J = 7.9 Hz, 1H), 4.278 (dd, J_1 = 1.4, J_2 = 17.7 Hz, 1H), 4.08 (d, J = 17.7 Hz, 1H), 3.81 (bs, 1H), 1.381 (s, 3H), 1.209 (s, 3H). ^{13}C NMR (CDCl_3 , 125 MHz) δ 210.557, 147.655,

146.432, 127.858, 123.136, 101.386, 75.770, 71.719, 66.554, 23.441, 23.250. HRMS for $C_{13}H_{15}NO_6Na$ (MNa^+): calcd 304.0792, obsd 304.0788; HPLC (Daicel Chirapak OD-H, hexane/isopropanol = 90 : 10, flow rate 1.0 mL/min, λ = 254 nm): t_R = 9.08 min (*anti*, major), t_R = 10.34 min (*anti*, minor), t_R = 13.98 min (*syn*). $[\alpha]_D = -125.1$ ($c = 0.666$, $CHCl_3$).

(S)-4-((S)-hydroxy(4-nitrophenyl)methyl)1,5-Dioxaspiro[5.5]undecan-3-one (Table 1, entry 9): Column chromatography provided a separable diastereomeric mixture.

1H NMR ($CDCl_3$, 500 MHz), major product: δ 8.214 (d, $J = 8.4$, 2H), 7.60 (d, $J = 8.5$, 2H), 5.00 (d, $J = 8.2$, 1H), 4.40-4.41 (m, 2H), 4.09 (d, $J = 17.7$ Hz, 1H), 1.81-1.168 (m, 10H). ^{13}C NMR ($CDCl_3$, 125 MHz) δ 211.176, 147.450, 146.645, 127.913, 123.112, 101.477, 77.477, 75.496, 71.911, 66.320, 32.939, 31.894, 24.874, 22.579, 22.194, δ HRMS for $C_{16}H_{19}NO_6$ (MH^+): calcd 322.1285, obsd 322.1288; HPLC (Daicel Chirapak OD-H, hexane/isopropanol = 95:5, flow rate 0.5 mL/min, λ = 254 nm): t_R = 38.38 min (*anti*, major), t_R = 41.42 min (*anti*, minor), t_R = 48.67 min (*syn*, minor), t_R = 68.64 min (*anti*, minor) $[\alpha]_D = -68.16$ ($c = 0.6$ $CHCl_3$).

(S)-4-((S)-1-hydroxy-3-methylbutyl)-2,2-dimethyl-1,3-dioxan-5-one (Table 2, entry 2):

1H NMR ($CDCl_3$, 500 MHz) : δ 4.24 (dd, $J_1 = 1.4$ Hz, $J_2 = 17.3$ Hz, 1H), 4.05 (dd, $J_1 = 1.3$ Hz, $J_2 = 6.5$ Hz, 1H), 4.00 (d, $J = 17.3$, 1H), 3.972-3.934 (m, 1H), 2.87 (bs, 1H), 1.897-1.842 (m, 1H), 1.456 (s, 3H), 1.428 (s, 3H), 0.934 (d, $J = 6.7$ Hz, 3H), 0.900 (d, $J = 6.9$, 3H). ^{13}C NMR ($CDCl_3$, 125 MHz) δ 210.94, 100.9, 76.507, 68.84, 66.74, 41.19, 24.02, 23.84, 23.749, 23.425, 21.428. HRMS for $C_{11}H_{20}O_4Na$ (MNa^+): calcd 239.1254, obsd 239.1256. $[\alpha]_D = -211.5$ ($c = 0.667$ $CHCl_3$).

HPLC (Daicel Chirapak OJ-H, hexane/*i*-PrOH = 95:5, flow rate 1.0 mL/min, λ = 254 nm): t_R = 31.92 min (major, *anti*), t_R = 35.73 min (minor, *anti*) t_R = 49.07 min (*syn*).

(S)-4-((S)-cyclopentyl(hydroxy)methyl)-2,2-dimethyl-1,3-dioxan-5-one (Table 2, entry 3):

^1H NMR (CDCl_3 , 500 MHz) : δ 4.26 (d, J = 16.4 Hz, 1H), 4.12 (d, J = 6.8 Hz, 1H), 4.01 (d, J = 17.3 Hz, 1H), .83, (m, 1H), 2.96 (d, J = 2.6 Hz), 2.27-2.21 (m, 1H), 1.69-1.49 (m, 8H), 1.47 (s, 3H), 1.44 (s, 3H).

^1H NMR (CDCl_3 , 500 MHz) : δ 9.31 (t, J = 2.1 Hz, 1H), 9.20 (d, J = 2.1, 2H), 5.65 (dd, J_1 = 4.8

Hz, J_2 = 7.4 Hz, 1H), 4.597 (dd, J_1 = 1.0 Hz, J_2 = 4.7 Hz), 4.37 (dd, J_1 = 1.3 Hz, J_2 = 17.2 Hz, 1H), 4.11 (d, J = 17.2 Hz, 1H), 2.72-2.64 (m, 1H), 1.88-1.62 (m, 8H), 1.543 (d, J = 19.7 Hz), 1.55 (s, 3H), 1.52 (s, 3H). ^{13}C NMR (CDCl_3 , 125 MHz) δ 206.4, 162.05, 148.70, 133.99, 129.44, 122.4, 101.0, 77.6, 74.87, 66.988, 40.22, 28.93, 28.48, 25.36,

24.90, 24.15, 23.34. HRMS for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_9\text{Na}$ (MNa^+): calcd 445.1217, obsd 445.1217. HPLC (Daicel Chirapak AD, hexane/*i*-PrOH = 95:5, flow rate 1.0 mL/min, λ = 254 nm): t_R = 12.90 min (major), t_R = 15.12 min (minor). $[\alpha]_D = -88.2$ (c = 0.833 CHCl_3).

2-((S)-2-((S)-2,2-dimethyl-5-oxo-1,3-dioxan-4-yl)-2-hydroxyethyl)isoindoline-1,3-dione

(Table 2, entry 4): ^1H NMR (CDCl_3 , 500 MHz) : δ 7.86 (d, J = 3.0 Hz, 1H), 7.85 (d, J = 3.0 Hz, 1H), 7.72 (d, J = 3.0 Hz, 1H), 7.71 (d, J = 3.0 Hz, 1H), 4.32 (m, 2H), 4.29 (d, J = 18 Hz, 1H), 4.02 (d, J = 17.5 Hz, 1H), 3.97 – 3.93 (m, 2H), 3.24 (d, J = 4.5 Hz, 1H, O-*H*), 1.47 (s, CH_3 , 3H), 1.34 (s, CH_3 , 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 209.5, 168.6, 134.0, 132.0, 123.3, 101.2, 75.6, 68.2, 66.6, 39.9, 23.5, 23.5. HRMS for $\text{C}_{16}\text{H}_{17}\text{NO}_6\text{Na}$

(MNa^+): calcd 342.0948, obsd 342.0943; HPLC (Daicel Chirapak AD, hexane/*i*-PrOH = 80:20, flow rate 1.0 mL/min, λ = 254 nm): t_R = 23.07min (minor, *anti*), t_R = 38.37 min (major, *anti*). $[\alpha]_D = -114.3$ (c = 0.315 CHCl_3).

(S)-2-((S)-2,2-dimethyl-5-oxo-1,3-dioxan-4-yl)-2-hydroxyethyl acetate (Table 2, entry 5):

^1H NMR (CDCl_3 , 500 MHz): δ 4.34 – 4.18 (m, 4H), 4.13 (m, 1H), 4.04 (d, J = 17.5 Hz, 1H), 3.20 (d, J = 3.5 Hz, 1H, O-*H*), 2.08 (s, COCH_3 , 3H), 1.44 (s, CH_3 , 3H), 1.41 (s, CH_3 , 3H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 210.6,

170.9, 101.3, 72.9, 68.8, 66.5, 64.2, 23.6, 23.4, 20.8. HRMS for $C_{10}H_{16}O_6Na$ (MNa^+): calcd 255.0839, obsd 255.0832; $[\alpha]_D = -82.6$ ($c = 0.615$ $CHCl_3$).

1H NMR ($CDCl_3$, 500 MHz): δ 9.22 (br t, $J = 2.0$ Hz, 1H), 9.11 (s, 1H), 9.10 (s, 1H), 5.71 (m, 1H), 4.67 (d, $J = 5.5$ Hz, 1H), 4.45 (m, 2H), 4.31 (d, $J = 17.5$ Hz, 1H), 4.10 (m, 1H), 2.05 (s, $COCH_3$, 3H), 1.49 (s, CH_3 , 3H), 1.46 (s, CH_3 , 3H); ^{13}C NMR ($CDCl_3$, 125

MHz): δ 205.7, 170.4, 161.6, 148.6, 133.2, 129.5, 122.6, 101.6, 72.2, 71.6, 66.7, 61.5, 23.5, 23.4. HRMS for $C_{17}H_{18}N_2O_{11}$ (MH^+): calcd 427.0983, obsd 427.0995; HPLC (Daicel Chirapak AD, hexane/*i*-PrOH = 95:5, flow rate 1.0 mL/min, $\lambda = 254$ nm): $t_R = 45.18$ min (*anti*, major), $t_R = 73.04$ min (*anti*, minor), $t_R = 54.03$ min (*syn*). $[\alpha]_D = -58.5$ ($c = 0.54$ $CHCl_3$).

(S)-4-((S)-((R)-2,2-dimethyl-1,3-dioxolan-4-yl)(hydroxy)methyl)-2,2-dimethyl-1,3-dioxan-5-one (Table 2, entry 6): 1H NMR and ^{13}C NMR were in accordance with lit. values.² 1H NMR

($CDCl_3$, 500 MHz): δ 4.28-4.24 (m, 3H), 4.03 (d, $J = 17.5$ Hz, 1H), 3.98 (dd, $J_1 = 6.6$ Hz, $J_2 = 8.2$ Hz, 1H), 3.86-3.81 (m, 2H), 3.19 (d, 3.5 Hz, 1H), 1.45 (s, 3H), 1.41 (s, 3H), 1.40 (s, 3H), 1.34 (s, 3H); ^{13}C NMR ($CDCl_3$, 125 MHz): δ 210.4, 109.2, 101.3, 75.2, 73.4, 70.1, 66.7, 65.6, 26.3, 25.6, 23.6,

23.5. HRMS for $C_{12}H_{20}O_6$ (MH^+): calcd 261.1333, obsd 261.1328; $[\alpha]_D = -157.9$ ($c = 1.61$ $CHCl_3$), lit. -167.2 ($c = 1.1$ $CHCl_3$). Mp = 102-104 °C, lit. 103-105 °C.

1H NMR ($CDCl_3$, 500 MHz): δ 9.23 (br d, $J = 8.0$ Hz, 1H), 9.16 (s, 1H), 9.13 (s, 1H), 5.51 (br dd, $J = 6.5, 3.5$ Hz, 1H), 4.67 (m, 1H), 4.57 (m, 1H), 4.31 (AB q, $J = 14.0$ Hz, 1H), 4.14 – 3.99 (m, 2H), 3.81 (dd, $J = 7.5, 5.0$ Hz, 1H), 1.53 (s, CH_3 , 3H), 1.47 (s, CH_3 , 3H), 1.44 (s, CH_3 , 3H), 1.34 (s, CH_3 , 3H); ^{13}C NMR ($CDCl_3$, 125 MHz): 207.5,

162.8, 149.5, 134.5, 130.6, 130.4, 110.6, 102.5, 74.6, 73.4, 72.5, 67.7, 66.3, 27.1, 26.0, 25.8, 24.4. HRMS: $C_{19}H_{22}N_2O_{11}Na$ (MNa^+): calcd 477.1116, obsd 477.1110; HPLC (Daicel Chirapak AD, hexane/*i*-PrOH = 90:10, flow rate 1.0 mL/min, $\lambda = 254$ nm): $t_R = 10.6$ min (*anti*, major), $t_R = 19.80$ min (*anti*, minor).

General procedure for deprotection with resin. 2-((S)-2-((S)-2,2-dimethyl-5-oxo-1,3-dioxan-

4-yl)-2-hydroxyethyl)isoindoline-1,3-dione (0.05 mmols, 18 mg) dissolved in H₂O (200 μ L) and THF (300 μ L) was stirred with Dowex 50WX2-100 resin (prewashed with H₂O) for 12 h at RT. The solution was filtered and the filtrate concentrated to give 2-((2S,3S)-2,3,5-trihydroxy-4-oxopentyl)isoindoline-1,3-dione (Yield: 100 %); no purification was necessary. ¹H NMR (dDMSO, 500 MHz): δ 7.88-7.82 (m, 4H), 5.70 (d, J = 5.3 Hz, 1H), 5.31 (d, J = 5.4 Hz, 1H), 4.90 (t, J = 5.9 Hz, 1H) 4.32 (dq, J_1 = 6.0 Hz, J_2 = 19.4 Hz, 2H), 4.03-3.96 (m, 2H), 3.73 (dd, J_1 = 9.2 Hz, J_2 = 13.8 Hz, 1H), 3.56 (dd, J_1 = 3.13 Hz, J_2 = 13.1 Hz, 1H); ¹³C NMR (dDMSO, 125 MHz) δ 211.0, 167.9, 134.1, 131.7, 122.8, 76.3, 69.0, 66.4, 41.0. HRMS for C₁₃H₁₃NNaO₆ (MNa⁺): calcd 302.0635, obsd 302.0634.

2-((S)-2-hydroxy-2-((4S,5S)-5-hydroxy-2,2-dimethyl-1,3-dioxan-4-yl)ethyl)isoindoline-1,3-

dione (1): A solution of 2-((S)-2-((S)-2,2-dimethyl-5-oxo-1,3-dioxan-4-yl)-2-hydroxyethyl)isoindoline-1,3-dione (58 mg, 0.18 mmols) in THF (1.2 mL) was cooled to -78 °C under argon and L-selectride[®] (0.19 mL of 1.0 M in THF) was added dropwise. The solution was stirred and allowed to slowly reach room temperature over 5 h at which time it was quenched with 0.5 mL of an aqueous NH₄Cl (sat.) solution. H₂O₂ (100 μ L of 30 % aq. solution) was added followed by NaOH (0.1 N, 100 μ mL) and the solution stirred for 30 min. The aqueous layer was extracted with EtOAc and dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by column chromatography (gradient elution of 40% EtOAc/hexane to 100 % EtOAc to give a white solid (Yield: 50 %). ¹H NMR (MeOD, 500 MHz): δ 7.86-7.78 (m, 4H), 4.15-4.04 (m, 2H), 3.89 (dd, J_1 = 5.7 Hz, J_2 = 13.9 Hz, 2H), 3.80-3.73 (m, 3H) 3.62 (d, J = 1.5 Hz, 1H), 1.41 (s, 3H), 1.17 (s, 3H); ¹³C NMR (MeOD, 125 MHz) δ 170.1, 135.3, 133.6, 124.0, 99.8, 75.8, 67.2, 67.0, 63.5, 42.8, 29.4, 18.7. HRMS for C₁₆H₁₉NNaO₆ (MNa⁺): calcd 344.1105, obsd 344.1101.

1-Amino-1-Deoxy-D-Lyxitol (2): TFA (23 μ L) was added to a solution of **1** (0.063 mmols, 20.3 mg) in THF/H₂O (3:1, 1 mL) and the mixture was heated at 40 °C for 6 h.

The reaction was quenched with saturated NaHCO₃ (1 mL) and brine (1 mL) and extracted with THF. The organic layer was dried over NaCl and MgSO₄

and concentrated *in vacuo* to give the phthalimide protected intermediate as a crude mixture. The mixture was dissolved in EtOH (1 mL), treated with methylamine (1.5 mL, 30 % in EtOH), and heated to reflux for 3 h.³ The solvent was removed *in vacuo* and the residue purified by column chromatography (dry loading, gradient elution with 10 % MeOH/CH₂Cl₂ to 50 % MeOH/CH₂Cl₂; then with 100 % AcOH) to give a colorless solid (60 % over two steps). The product was isolated as the acetate salt and its spectroscopic data were identical to those in the literature for the HCl salt.⁴ ¹H NMR (D₂O with TSP, 500 MHz): δ 3.94 (m, 2H), 3.67 (d, J = 6.4 Hz, 2H), 3.58 (dd, J_1 = 1.7 Hz, J_2 = 8 Hz, 1H), 3.39 (dd, J_1 = 3.2 Hz, J_2 = 13.2 Hz, 1H), 3.05 (dd, J_1 = 9.2 Hz, J_2 = 13.2 Hz), 2.90 (d, J = 3.2 Hz, 1H, OH); ¹³C NMR (D₂O with TSP, 125 MHz) δ 74.76, 70.06, 65.66, 45.20. HRMS for C₁₇H₂₃NO₄Na (M⁺): calcd 152.0917, obsd 152.0923.

References:

1. Kim, K. S.; Hong, S. D. *Tetrahedron Lett.* **2000**, *41*, 5909-5913.
2. Majewski, M.; Nowak, P. *J. Org. Chem.* **2000**, *65*, 5152-5160.
3. Motawia, M. S.; Meldal, M.; Sofan, M.; Stein, P.; Pedersen, E. B.; Nielsen, C. *Synthesis* **1995**, 265-70.
4. Blanc-Muesser, M.; Defaye, J.; Horton, D. *Carbohydr. Res.* **1979**, *68*, 175-187.

S-11

S-15

S-22

S-24

4.276
4.267
4.264
4.251
4.235
4.053
4.018
3.984
3.971
3.967
3.954
3.860
3.845
3.829
3.825
3.818
3.809
3.196
3.189

1.450
1.410
1.398
1.344

S-26

S-28

S-30

X-ray structure of (S)-4-((S)-Hydroxy(4-nitrophenyl)methyl)-2,2-dimethyl-1,3-dioxan-5-one (Table 1, entry 6).