Modular Synthesis of Chiral Homo- and Hetero-trisoxazolines.[†] Improving the Enantioselectivity in the Asymmetric Michael Addition of Indole to Benzyldiene Malonate

Meng-Chun Ye, Bin Li, Jian Zhou, Xiu-Li Sun and Yong Tang*

State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, 354 Fenglin Lu, Shanghai 200032, China.

tangy@mail.sioc.ac.cn

Supporting Information

Contents

General Methods.....	S2
General Procedure for the Synthesis of 2-Chloromethyl Oxazolines.....	S2
General Procedure for the Synthesis of Bis(oxazolines) 2B and 2C.....	S4
General Procedure for the Synthesis of Homo- and Hetero-chiral Trisoxazolines.....	S5
General Procedures for Catalytic Asymmetric Michael Addition Reaction.....	S9
Copies of ¹ H NMR and ¹³ C NMR of compounds.....	S10
Copies of HPLC spectra of Michael Addition products.....	S27

General Methods. All reactions were carried out under a nitrogen atmosphere in oven-dried glassware with stirring. Dichloromethane and triethylamine were distilled over calcium hydride. Tetrahydrofuran was dried by distillation over sodium / benzophenone ketyl. *tert*-Butyllithium was purchased from Aldrich. Aminoalcohols¹, methyl chloroacetimidate hydrochloride² and bisoxazolines 2A and 2D³ were prepared according to literature procedures.

General Procedure for the Synthesis of 2-Chloromethyl Oxazolines. To a stirred milky solution of methyl chloroacetimidate hydrochloride (5.5 mmol) in CH₂Cl₂ (15 mL) was added a solution of amino alcohol **6** (5 mmol) in CH₂Cl₂ by syringe, followed by triethylamine (0.75 mL, 5.5 mmol) at 0 °C under nitrogen. The resulting mixture was stirred for six to eight hours at 20 °C until the color of reaction mixture became pink. The solvent was removed under reduced pressure and the resulting paste was extracted with ethyl acetate (3 x 10 mL). The combined organic extracts were concentrated in vacuo to afford the crude product, which was purified by flash chromatography (petroleum ether/ethyl acetate = 1/6, containing 3% triethylamine) to give the pure product.

(4S)-2-chloromethyl-4-isopropyl-2-oxazoline (**3a**)⁴: colorless oil (83%); [α]_D²⁰ -103.2 ° (c 5.42, CHCl₃); ¹H NMR (300 MHz, CDCl₃): 4.36 (dd, *J* = 8.4, 9.6 Hz, 1H), 4.12 (s, 2H), 4.07 (t, *J* = 8.4 Hz, 1H), 4.01-3.96 (m, 1H), 1.79 (m, 1H), 0.98 (d, *J* = 6.9 Hz, 3H), 0.90 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃): 162.3, 72.3, 71.1, 36.4, 32.4, 18.6, 18.0.

(4S)-2-chloromethyl-4-*sec*-butyl-2-oxazoline (**3b**): colorless oil (81%); [α]_D²⁰ -95.6 ° (c 7.29, CHCl₃); IR (neat) 1669 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): 4.33-4.24 (m, 1H), 4.06 (s, 2H), 4.02 (dd, *J* = 8.4, 12.9 Hz, 2H), 1.65-1.42 (m, 2H), 1.21-1.07 (m, 1H), 0.88 (t, *J* = 7.2 Hz, 3H), 0.79 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃): δ 162.1, 70.8, 70.6, 38.6, 36.3, 25.8, 14.1, 11.3; LRMS-EI(*m/z*): 175(M⁺); HRMS(*m/z*) calcd for C₈H₁₅NCIO (M+H⁺) 176.0837, found 176.0822.

(4S)-2-chloromethyl-4-*tert*-butyl-2-oxazoline (**3c**): colorless oil (73%); [α]_D²⁰ -105.6 ° (c 5.49, CHCl₃); IR (neat) 1670 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): 4.28 (dd, *J* = 8.7, 10.2 Hz, 1H), 4.15 (t, *J* = 8.4 Hz, 1H), 4.11 (s, 2H), 3.91 (dd, *J* = 8.4,

10.2 Hz, 1H), 0.89 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): δ 162.2, 75.9, 69.6, 36.4, 33.6, 25.6; LRMS-EI(m/z): 176(M^+); HRMS(m/z) calcd for $\text{C}_8\text{H}_{15}\text{NClO}$ ($\text{M}+\text{H}^+$) 176.0837, found 176.0850.

(4S)-2-chloromethyl-4-phenyl-2-oxazoline (**3d**)⁴: colorless oil (85%); $[\alpha]_{\text{D}}^{20}$ -122.2° (c 3.14, CHCl_3); ^1H NMR (300 MHz, CDCl_3): 7.40-7.22 (m, 5H), 5.26 (t, $J = 9.6$ Hz, 1H), 4.74 (dd, $J = 8.7, 10.2$ Hz, 1H), 4.24 (t, $J = 8.4$ Hz, 1H), 4.23 (s, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 163.6, 141.2, 128.7, 127.7, 126.4, 75.5, 69.6, 36.3.

(4R)-2-chloromethyl-4-phenyl-2-oxazoline (**3e**)⁴: colorless oil (74%); $[\alpha]_{\text{D}}^{20}$ $+126.9^\circ$ (c 6.51, CHCl_3); ^1H NMR (300 MHz, CDCl_3): 7.40-7.23 (m, 5H), 5.27 (t, $J = 9.3$ Hz, 1H), 4.74 (dd, $J = 8.4, 9.9$ Hz, 1H), 4.24 (t, $J = 8.1$ Hz, 1H), 4.23 (s, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 163.7, 141.2, 128.8, 127.8, 126.5, 75.6, 69.7, 36.3.

(4S)-2-chloromethyl-4-benzyl-2-oxazoline (**3f**)⁴: colorless oil (79%); $[\alpha]_{\text{D}}^{20}$ -47.2° (c 3.79, CHCl_3); ^1H NMR (300 MHz, CDCl_3): 7.36-7.22 (m, 5H), 4.55-4.45 (m, 1H), 4.33 (t, $J = 8.4$ Hz, 1H), 4.13 (s, 2H), 4.10 (t, $J = 8.1$ Hz, 1H), 3.14 (dd, $J = 5.4, 13.8$ Hz, 1H), 2.73 (dd, $J = 5.4, 13.5$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3): 162.9, 137.3, 129.1, 128.5, 126.6, 72.7, 67.5, 41.2, 36.3.

(4S)-2-chloromethyl-4-isopropyl-5-dimethyl-2-oxazoline (**3g**): colorless oil (28%); $[\alpha]_{\text{D}}^{20}$ -42.6° (c 3.00, CHCl_3); IR (neat) 1664 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): 4.04 (s, 2H), 3.29 (d, $J = 8.7$ Hz, 1H), 1.86-1.74 (m, 1H), 1.45 (s, 3H), 1.30 (s, 3H), 1.05 (d, $J = 6.6$ Hz, 3H), 0.96 (d, $J = 6.6$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3): 160.8, 88.1, 80.2, 37.1, 28.8, 28.7, 20.9, 20.4; LRMS-EI(m/z): 189(M^+); HRMS(m/z) calcd for $\text{C}_9\text{H}_{17}\text{ClNO}$ ($\text{M}+\text{H}^+$) 190.0993, found 190.1001.

(3aS,8aR)-2-chloromethyl-3a,8a-dihydro-8H-indeno[1,2-d]oxazole (**3h**)⁴: white solid (67%); $[\alpha]_{\text{D}}^{20}$ -319.9° (c 2.84, CHCl_3); ^1H NMR (300 MHz, CDCl_3): 7.50-7.46 (m, 1H), 7.30-7.26 (m, 3H), 5.62 (d, $J = 7.8$ Hz, 1H), 5.47-5.42 (m, 1H), 4.10 (ABd, $J = 12.9$ Hz, 1H), 4.05 (ABd, $J = 12.9$ Hz, 1H), 3.46 (dd, $J = 6.6, 18.0$ Hz, 1H), 3.29 (d, $J = 12.9$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3): 162.8, 141.0, 140.0, 128.7, 127.5, 125.4, 125.3, 84.3, 76.7, 39.6, 36.5.

General Procedure for the Synthesis of Bis(oxazolines) 2B and 2C. According to literature procedure,³ but SOCl_2 instead of MsCl as the reaction reagent because the

reaction can not run in the presence of MsCl. To a stirred milky solution of diamide (5 mmol) in dry CHCl_3 (20 mL) was added dropwise SOCl_2 (1.8 mL, 45 mmol) at 0 °C under nitrogen. The mixture was stirred at 0 °C for 20 minutes and then was refluxed for one hour. After cooling to 0 °C, the mixture was diluted with 20 mL CH_2Cl_2 and was washed twice with H_2O (2 x 10 mL). The aqueous layer was extracted with CH_2Cl_2 (20 mL) and the combined organic extracts were dried over Na_2SO_4 , filtered and concentrated to a thick oil. To the crude bischloride was added a solution of NaOH (0.5M, $\text{MeOH}/\text{H}_2\text{O}$ = 1:1, 50 mL) and the resulting mixture was refluxed about three hours then was cooled to room temperature and was extracted with CH_2Cl_2 (3 x 40 mL). The combined organic phase was dried over Na_2SO_4 , filtered and concentrated to a pale yellow oil. The crude product was purified by flash chromatography (petroleum ether/acetone = 1/10) to give the desired product.

1,1'-Bis[2-((4S)-phenyl-1,3-oxazolinyl)]ethane (**2B**): colorless oil (58%); $[\alpha]_{\text{D}}^{20}$ -102.0 ° (c 1.16, CHCl_3); IR (neat) 3030, 2896, 1660, 1567, 1494, 1455, 1355, 1267, 1236, 1188, 1077, 1029, 977, 917, 760, 700 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): 7.34-7.23 (m, 10H), 5.23 (t, J = 9.9 Hz, 2H), 4.66 (t, J = 9.3 Hz, 1H), 4.16 (dt, J = 3.3, 7.5 Hz, 2H), 3.76 (q, J = 7.5 Hz, 1H), 1.65 (d, J = 7.5 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3): 167.1, 166.8, 142.1, 142.1, 128.6, 127.5, 127.5, 126.6, 126.5, 75.2, 75.2, 69.4, 69.4, 34.09, 15.3; LRMS-EI(m/z): 320 (M^+); HRMS(m/z) calcd for $\text{C}_{20}\text{H}_{21}\text{N}_2\text{O}_2$ ($\text{M}+\text{H}^+$) 321.1597, found 321.1585.

1,1'-Bis[2-((4R)-phenyl-1,3-oxazolinyl)]ethane (**2C**): colorless oil (51%); $[\alpha]_{\text{D}}^{20}$ +103.0 ° (c 1.01, CHCl_3); ^1H NMR (300 MHz, CDCl_3): 7.35-7.23 (m, 10H), 5.24 (t, J = 9.6 Hz, 2H), 4.67 (t, J = 9.0 Hz, 1H), 4.16 (dt, J = 3.3, 7.8 Hz, 2H), 3.77 (q, J = 7.5 Hz, 1H), 1.64 (d, J = 6.9 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3): 167.0, 166.8, 142.1, 142.1, 128.6, 127.5, 127.4, 126.6, 126.5, 75.2, 75.2, 69.4, 69.4, 34.0, 15.3.

General Procedure for the Synthesis of Chiral Homo- and Hetero-trisoxazolines.

To a solution of bisoxazoline (**2A-D**) (2 mmol) in dried THF (30 mL) was added dropwise *t*-BuLi (1.3 mL, 1.7 M in hexanes, 2.2 mmol) within 15~20 minutes at -78

under nitrogen. The resulting yellow solution was stirred for an additional one hour at this temperature. Then a solution of 2-chloromethyl oxazoline (2.8 mmol) in THF (10 mL) was added dropwise at -78 °C over 10 minutes. The solution was slowly warmed to room temperature and was stirred for further 10 hours. The mixture was diluted with CH₂Cl₂ (20 mL) and was washed with H₂O (5 mL). The aqueous layer was extracted with CH₂Cl₂ (5 mL) and the combined organic phases was dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography (petroleum ether/acetone = 1/10) to give the desired product.

1,2,2-tris[2-[(4S)-4-isopropyl-1,3-oxazolinyl]]propane (**1a**):⁵ colorless oil (74%); [α]_D²⁰ -47.2 ° (c 3.79, CHCl₃); ¹H NMR (300 MHz, CDCl₃): 4.22-4.07 (m, 3H), 4.00-3.77 (m, 6H), 3.05 (ABd, *J* = 14.7 Hz, 1H), 2.91 (ABd, *J* = 14.4 Hz, 1H), 1.82-1.62 (m, 3H), 1.57 (s, 3H), 0.91 (d, *J* = 6.6 Hz, 6H), 0.87 (d, *J* = 6.9 Hz, 3H), 0.82 (d, *J* = 7.2 Hz, 6H), 0.81 (d, *J* = 7.2 Hz, 3H).

1-[2-[(4S)-4-*sec*-butyl-1,3-oxazolinyl]]-2,2-bis[2-[(4S)-4-isopropyl-1,3-oxazolinyl]]propane (**1Ab**): colorless oil (67%); [α]_D²⁰ -97.0 ° (c 0.89, CHCl₃); IR (neat) 2960, 2903, 2875, 1662, 1466, 1384, 1356, 1236, 1171, 1096, 983, 752 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): 4.21-4.02 (m, 3H), 3.98-3.78 (m, 6H), 3.04 (ABd, *J* = 14.4 Hz, 1H), 2.87 (ABd, *J* = 14.4 Hz, 1H), 1.80-1.66 (m, 2H), 1.55 (s, 3H), 1.52-1.40 (m, 2H), 1.20-1.03 (m, 1H), 0.88 (t, *J* = 6.6 Hz, 3H), 0.86 (d, *J* = 6.9 Hz, 3H), 0.85 (d, *J* = 6.9 Hz, 3H), 0.81 (d, *J* = 6.6 Hz, 3H), 0.80 (d, *J* = 6.9 Hz, 3H), 0.75 (d, *J* = 7.2 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃): 166.8, 166.8, 163.4, 71.9, 71.4, 70.6, 70.3, 69.9, 69.2, 40.7, 38.7, 34.8, 32.3, 32.0, 25.9, 21.2, 18.7, 18.4, 17.7, 17.2, 14.2, 11.3; LRMS-EI(*m/z*): 391 (*M*⁺); HRMS(*m/z*) calcd for C₂₂H₃₈N₃O₃ (*M*+*H*⁺) 392.2908, found 392.2895.

1-[2-[(4S)-4-*tert*-butyl-1,3-oxazolinyl]]-2,2-bis[2-[(4S)-4-isopropyl-1,3-oxazolinyl]]propane (**1Ac**): colorless oil (84%); [α]_D²⁰ -116.1 ° (c 0.91, CHCl₃); IR (neat) 2957, 2904, 2872, 1661, 1479, 1468, 1362, 1236, 1171, 1095, 983, 929 cm⁻¹; ¹H NMR (300

MHz, CDCl₃): 4.23-3.75 (m, 9H), 3.07 (ABd, $J = 15.0$ Hz, 1H), 2.93 (ABd, $J = 15.3$ Hz, 1H), 1.83-1.69 (m, 2H), 1.58 (s, 3H), 0.92 (d, $J = 6.6$ Hz, 3H), 0.89 (d, $J = 6.6$ Hz, 3H), 0.84 (s, 9H), 0.83 (d, $J = 6.6$ Hz, 3H), 0.82 (d, $J = 6.6$ Hz, 3H); ¹³C NMR (75 MHz, CDCl₃): 166.9, 166.9, 163.5, 75.7, 71.9, 71.5, 70.3, 69.9, 68.2, 40.7, 34.8, 33.4, 32.3, 32.1, 25.8, 21.2, 18.8, 18.5, 17.8, 17.3; LRMS-EI(m/z): 391 (M^+); HRMS(m/z) calcd for C₂₂H₃₈N₃O₃ ($M+H^+$) 392.2908, found 392.2878.

1-[2-[(4S)-4-phenyl-1,3-oxazoliny]]-2,2-bis[2-[(4S)-4-isopropyl-1,3-oxazolnyl]]propane (**1Ad**): colorless oil (77%); [α]_D²⁰ -93.3 ° (c 0.93, CHCl₃); IR (neat) 2959, 2902, 1660, 1468, 1356, 1236, 1172, 1096, 983, 757, 701 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): 7.34-7.21 (m, 5H), 5.14 (t, $J = 9.6$ Hz, 1H), 4.53 (dd, $J = 8.1, 9.9$ Hz, 1H), 4.26-4.17 (m, 2H), 4.03-3.89 (m, 5H), 3.18 (ABd, $J = 14.7$ Hz, 1H), 3.04 (ABd, $J = 14.4$ Hz, 1H), 1.84-1.64 (m, 2H), 1.67 (s, 3H), 0.91 (t, $J = 7.2$ Hz, 3H), 0.84 (d, $J = 6.6$ Hz, 6H); ¹³C NMR (75 MHz, CDCl₃): 166.90, 166.8, 165.1, 142.2, 128.5, 127.3, 126.6, 74.4, 71.9, 71.6, 70.4, 70.1, 69.6, 40.9, 35.0, 32.3, 32.1, 21.4, 18.8, 18.5, 17.7, 17.4; LRMS-EI(m/z): 411 (M^+), 231 (100); HRMS(m/z) calcd for C₂₄H₃₄N₃O₃ ($M+H^+$) 412.2600, found 412.2620.

1-[2-[(4R)-4-phenyl-1,3-oxazoliny]]-2,2-bis[2-[(4S)-4-isopropyl-1,3-oxazolnyl]]propane (**1Ae**): colorless oil (55%); [α]_D²⁰ -21.8 ° (c 0.96, CHCl₃); IR (neat) 2959, 2901, 2873, 1661, 1468, 1385, 1357, 1236, 1169, 1096, 982, 757 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): 7.33-7.20 (m, 5H), 5.11 (t, $J = 9.0$ Hz, 1H), 4.53 (dd, $J = 8.7, 10.5$ Hz, 1H), 4.24-4.14 (m, 2H), 4.01-3.85 (m, 5H), 3.14 (ABd, $J = 14.4$ Hz, 1H), 3.06 (ABd, $J = 14.1$ Hz, 1H), 1.83-1.63 (m, 2H), 1.66 (s, 3H), 0.91 (d, $J = 7.2$ Hz, 3H), 0.88 (d, $J = 6.6$ Hz, 3H), 0.83 (d, $J = 6.9$ Hz, 3H), 0.81 (d, $J = 6.6$ Hz, 3H); ¹³C NMR (75 MHz, CDCl₃): 166.8, 166.7, 165.1, 142.2, 128.5, 127.3, 126.6, 74.4, 71.9, 71.4, 70.4, 70.0, 69.6, 40.9, 35.0, 32.4, 32.0, 21.4, 18.8, 18.4, 17.8, 17.3; LRMS-EI(m/z): 411 (M^+); HRMS(m/z) calcd for C₂₄H₃₄N₃O₃ ($M+H^+$) 411.2516, found 411.2522.

1-[2-[(4S)-4-benzyl-1,3-oxazoliny]]-2,2-bis[2-[(4S)-4-isopropyl-1,3-oxazolnyl]]propane

ane (**1Af**): colorless oil (50%); $[\alpha]_{\text{D}}^{20}$ -80.2° (c 1.14, CHCl_3); IR (neat) 2958, 2900, 2873, 1661, 1468, 1356, 1235, 1170, 1095, 980, 931, 701 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): 7.31-7.16 (m, 5H), 4.38-4.15 (m, 3H), 4.08-3.83 (m, 6H), 3.10 (dd, $J = 5.1, 13.5$ Hz, 1H), 3.07 (ABd, $J = 14.1$ Hz, 1H), 2.93 (ABd, $J = 14.4$ Hz, 1H), 2.57 (dd, $J = 9.0, 13.8$ Hz, 1H), 1.85-1.67 (m, 2H), 1.60 (s, 3H), 0.91 (t, $J = 6.0$ Hz, 6H), 0.84 (d, $J = 6.6$ Hz, 6H); ^{13}C NMR (75 MHz, CDCl_3): 166.9, 166.9, 164.4, 138.0, 129.1, 128.5, 126.4, 71.9, 71.6, 71.5, 70.4, 70.0, 67.3, 41.7, 40.82, 34.9, 32.4, 32.1, 21.3, 18.8, 18.5, 17.8, 17.3; LRMS-EI(m/z): 425 (M^+); HRMS(m/z) calcd for $\text{C}_{25}\text{H}_{36}\text{N}_3\text{O}_3$ ($\text{M}+\text{H}^+$) 426.2751, found 426.2755.

1-[2-[(4S)-4-isopropyl-5,5-dimethyl-1,3-oxazoliny]]-2,2-bis[2-[(4S)-4-isopropyl-1,3-oxazoliny]]propane (**1Ag**): colorless oil (36%); $[\alpha]_{\text{D}}^{20}$ -98.6° (c 3.40, CHCl_3); IR (neat) 2959, 2905, 2873, 1661, 1469, 1385, 1370, 1241, 1175, 1096, 983, 941 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): 4.21-4.11 (m, 2H), 4.00-3.82 (m, 4H), 3.17 (d, $J = 8.1$ Hz, 1H), 2.98 (ABd, $J = 14.4$ Hz, 1H), 2.83 (ABd, $J = 14.7$ Hz, 1H), 1.82-1.63 (m, 3H), 1.57 (s, 3H), 1.35 (s, 3H), 1.20 (s, 3H), 1.00 (d, $J = 6.9$ Hz, 3H), 0.94 (d, $J = 6.6$ Hz, 3H), 0.89 (d, $J = 6.3$ Hz, 3H), 0.86 (d, $J = 6.6$ Hz, 3H), 0.81 (d, $J = 6.9$ Hz, 3H), 0.80 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3): 167.0, 166.9, 162.1, 86.0, 80.0, 72.0, 71.4, 70.4, 69.7, 40.9, 35.2, 32.5, 32.0, 29.3, 28.8, 21.3, 21.1, 21.0, 20.6, 18.9, 18.5, 17.8, 17.1; LRMS-EI(m/z): 405 (M^+); HRMS(m/z) calcd for $\text{C}_{23}\text{H}_{39}\text{N}_3\text{O}_3$ ($\text{M}+\text{H}^+$) 406.3064, found 406.3051.

1-[2-[(3aS,8aR)-3a,8a-dihydro-8H-indeno[1,2-d]oxazolyl]]-2,2-bis[2-[(4S)-4-isopropyl-1,3-oxazolyl]]propane (**1Ah**): colorless oil (74%); $[\alpha]_{\text{D}}^{20}$ -149.0° (c 0.82, CHCl_3); IR (neat) 2958, 1659, 1460, 1356, 1235, 1163, 1093, 1005, 981, 753 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): 7.41-7.40 (m, 1H), 7.20-7.14 (m, 3H), 5.43 (d, $J = 7.8$ Hz, 1H), 5.24-5.16 (m, 1H), 4.14-4.05 (m, 1H), 3.98-3.77 (m, 5H), 3.33 (dd, $J = 6.6, 18.0$ Hz, 1H), 3.14 (d, $J = 9.0$ Hz, 1H), 2.98 (ABd, $J = 14.7$ Hz, 1H), 2.85 (ABd, $J = 14.4$ Hz, 1H), 1.74-1.48 (m, 2H), 1.50 (s, 3H), 0.83 (t, $J = 7.5$ Hz, 6H), 0.76 (d, $J = 6.3$ Hz, 6H);

^{13}C NMR (75 MHz, CDCl_3): 166.7, 166.6, 164.2, 141.9, 139.4, 128.1, 127.1, 125.2, 125.0, 82.5, 76.3, 71.7, 71.3, 70.2, 69.7, 40.7, 39.5, 34.8, 32.2, 31.9, 21.3, 18.7, 18.3, 17.7, 17.2; LRMS-EI(m/z): 423 (M^+), 243 (100); HRMS(m/z) calcd for $\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$) 446.2414, found 446.2394.

1-[2-[(4S)-4-isopropyl-1,3-oxazoliny]]-2,2-bis[2-[(4S)-4-phenyl-1,3-oxazoliny]]propane (**1Ba**): colorless oil (52%); $[\alpha]_{\text{D}}^{20} -154.4^\circ$ (c 1.18, CHCl_3); IR (neat) 2959, 2899, 2873, 1658, 1494, 1455, 1356, 1232, 1170, 1097, 981, 921, 760, 700 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): 7.32-7.20 (m, 10H), 5.26-5.16 (m, 2H), 4.70-4.62 (m, 2H), 4.22-4.10 (m, 2H), 3.92-3.85 (m, 2H), 3.21(ABd, $J = 15.0$ Hz, 1H), 3.11 (ABd, $J = 14.7$ Hz, 1H), 1.80-1.59 (m, 1H), 1.78 (s, 3H), 0.95 (d, $J = 6.6$ Hz, 3H), 0.87 (d, $J = 5.7$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3): 168.4, 168.4, 163.1, 142.0, 142.0, 128.4, 128.3, 127.3, 126.6, 126.4, 126.4, 75.4, 75.3, 72.0, 69.7, 69.4, 69.3, 40.9, 34.8, 32.4, 21.2, 18.7, 18.1; LRMS-EI(m/z): 446 (M^+); HRMS(m/z) calcd for $\text{C}_{27}\text{H}_{32}\text{N}_3\text{O}_3$ ($\text{M}+\text{H}^+$) 446.2438, found 446.2413.

1-[2-[(4S)-4-isopropyl-1,3-oxazoliny]]-2,2-bis[2-[(4R)-4-phenyl-1,3-oxazoliny]]propane (**1Ca**): colorless oil (82%); $[\alpha]_{\text{D}}^{20} +96.0^\circ$ (c 1.16, CHCl_3); IR (neat) 2959, 2899, 2873, 1657, 1494, 1455, 1357, 1231, 1169, 1098, 981, 921, 759, 700 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): 7.36-7.23 (m, 10H), 5.25-5.18 (m, 2H), 4.70 (dd, $J = 8.1, 9.3$ Hz, 2H), 4.25-4.14 (m, 2H), 3.94-3.84 (m, 2H), 3.21(ABd, $J = 14.4$ Hz, 1H), 3.12 (ABd, $J = 14.4$ Hz, 1H), 1.78 (s, 3H), 1.78-1.67 (m, 1H), 0.97 (d, $J = 6.9$ Hz, 3H), 0.88 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3): 168.6, 163.5, 142.2, 128.6, 128.5, 127.5, 126.8, 126.6, 75.6, 75.5, 72.2, 70.0, 69.6, 69.5, 41.3, 35.1, 32.6, 21.5, 18.9, 18.3; LRMS-EI(m/z): 446 (M^+); HRMS(m/z) calcd for $\text{C}_{27}\text{H}_{32}\text{N}_3\text{O}_3$ ($\text{M}+\text{H}^+$) 446.2438, found 446.2418.

1,2,2-tris[2-[(4S)-4-*tert*-butyl-1,3-oxazoliny]]propane (**1Dc**): white solid (42%); $[\alpha]_{\text{D}}^{20} -114.6^\circ$ (c 1.05, CHCl_3); IR (neat) 2955, 2904, 2870, 1663, 1479, 1394, 1364,

1238, 1209, 1171, 1100, 980, 926 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): 4.19-3.77 (m, 9H), 3.14 (ABd, $J = 14.4$ Hz, 1H), 2.91 (ABd, $J = 14.7$ Hz, 1H), 1.60 (s, 3H), 0.86 (s, 3H), 0.85 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3): 166.9, 166.6, 163.5, 75.6, 75.4, 75.3, 68.9, 68.9, 68.1, 40.6, 34.7, 33.8, 33.6, 33.3, 25.7, 25.7, 25.6, 21.1; LRMS-EI(m/z): 419 (M^+); HRMS(m/z) calcd for $\text{C}_{24}\text{H}_{42}\text{N}_3\text{O}_3$ ($\text{M}+\text{H}^+$) 420.3221, found 420.3198.

General Procedures for Using Trisoxazolines/ $\text{Cu}(\text{OTf})_2$ as the Chiral Catalysts in Catalytic Asymmetric Michael Addition Reaction. To a Schlenk tube was added $\text{Cu}(\text{OTf})_2$ (0.10 eq), followed by trisoxazoline **1** (0.12 eq) and then $i\text{BuOH}$ (2.5 ml) under air atmosphere. The resulting blue-green solution was stirred at room temperature (10-25°C) for one hour before alkylidene malonate (1.0 eq) in $i\text{BuOH}$ (2.5 ml) was added. The mixture was stirred at room temperature for 15 minutes, and then the indole (1.2 eq) was added. After three hours, the reaction mixture was concentrated under reduced pressure at room temperature, and the residue was purified by flash column chromatography on silica gel [eluted with CH_2Cl_2 /petroleum ether (1/1, v/v) then pure CH_2Cl_2] to afford the desired product.

(S)-Ethyl 2-ethoxycarbonyl-3-(3-indolyl)-3-phenyl propanoate **8⁵:** ^1H NMR (300 MHz, CDCl_3): δ 8.06 (brs, 1H), 7.55 (d, $J = 7.8$ Hz, 1H), 7.36 (d, $J = 6.9$ Hz, 2H), 7.10-7.30 (m, 6H), 7.03-7.06 (m, 1H), 5.08 (d, $J = 11.7$ Hz, 1H), 4.29 (d, $J = 11.4$ Hz, 1H), 3.99 (m, 4H), 1.00 (m, 6H).

References:

1. Abiko, A.; Masamune, S. *Tetrahedron Lett.* **1992**, 33, 5517.
2. Dox, A. M. *Org. Synth. Coll. Vol I*, **1941**, 5.
3. Denmark, S. E.; Stiff, C. M. *J. Org. Chem.* **2000**, 65, 5875.
4. Kamata, K.; Sato, H.; Takagi, E.; Agata, I.; Meyers, A. I. *Heterocycles* **1999**, 51, 373.
5. Zhou, J.; Ye, M.-C.; Huang, Z.-Z. Tang, Y. *J. Org. Chem.* **2004**, 69, 1309.

1.

2.

3.

4.

5.

6.

7.

Test REPORT

Peak #	Component Name	Time [min]	Area [uV*sec]	Height [uV]	Area [%]	Norm. Area [%]	BL	Adjusted Amount
1		15.055	1099781.74	22765.29	7.19	7.19	MM	1.0998
2		18.375	14206041.58	214953.53	92.81	92.81	MM	14.2060
3		23.234	42.08	63.07	0.00	0.00	BB	0.0000
			15305865.40	237781.89	100.00	100.00		15.3059

8.

Test REPORT

Peak #	Component Name	Time [min]	Area [uV*sec]	Height [uV]	Area [%]	Norm. Area [%]	BL	Adjusted Amount
1		14.660	1363810.76	29184.42	4.97	4.97	MM	1.3638
2		17.632	26100812.51	382208.98	95.03	95.03	MM	26.1008
			27464623.26	411393.41	100.00	100.00		27.4646

9.

Test REPORT

Peak #	Component Name	Time [min]	Area [uV*sec]	Height [uV]	Area [%]	Norm. Area [%]	BL	Adjusted Amount
1		14.953	1548062.25	32738.92	7.46	7.46	MM	1.5481
2		18.202	19201317.99	280417.60	92.54	92.54	MM	19.2013
			20749380.23	313156.51	100.00	100.00		20.7494

10.

Test REPORT

Peak #	Component Name	Time [min]	Area [uV*sec]	Height [uV]	Area [%]	Norm. Area [%]	BL	Adjusted Amount
1		15.020	1272836.13	26896.79	4.59	4.59	MM	1.2728
2		18.433	26457128.98	370558.52	95.41	95.41	MM	26.4571
			27729965.11	397455.30	100.00	100.00		27.7300

11.

Test REPORT

Peak #	Component Name	Time [min]	Area [uV*sec]	Height [uV]	Area [%]	Norm. Area [%]	BL	Adjusted Amount
1		14.410	1685694.70	36633.90	13.47	13.47	MM	1.6857
2		17.643	10831306.64	175377.83	86.53	86.53	MM	10.8313
			12517001.34	212011.73	100.00	100.00		12.5170

12.

Test REPORT

Peak #	Component Name	Time [min]	Area [uV*sec]	Height [uV]	Area [%]	Norm. Area [%]	BL	Adjusted Amount
1		14.568	8293504.94	169864.62	80.43	80.43	MM	8.2935
2		18.532	2017311.96	32953.79	19.57	19.57	MM	2.0173
			10310816.90	202818.41	100.00	100.00		10.3108

13.