A multicomponent synthesis of spiroisoxazolines

Mauro F. A. Adamo, Donato Donati, Eleanor F. Duffy and Piero Sarti-Fantoni

Table of Contents

S2: General experimental.

S3-9: spectroscopic data for compounds **5d-p** and **3d-m**.

S10-26: ^1H -NMR spectra showing the epimerisation of **5a** to **4-*epi*-5a**.

S27-39: ^1H -NMR of compounds **5b-p**.

S40-52: ^1H -NMR of compounds **3a-m**.

S53-54: ^1H -NMR (400 MHz) and peak assignments of compounds **5a** and **4-*epi*-5a**.

S55-58: ^1H - ^1H correlation and ^1H - ^{13}C correlation NMR of compound **5a** and **4-*epi*-5a**.

S59: n.O.e. experiments for the assignment of **5a** and **4-*epi*-5a** (reference n. 9).

General experimental: ^1H and ^{13}C Spectra were recorded on a 200 or a 400 MHz spectrometers at ambient temperatures. ^1H NMR spectral assignments are supported by ^1H - ^1H COSY and ^{13}C - ^1H COSY where necessary. For ^1H NMR recorded in CDCl_3 chemical shifts (δ_{H}) are quoted in parts per million (ppm) and are referenced to the residual solvent peak. The following abbreviations are used: s, singlet, d, doublet, t, triplet, dd, doublet of doublets, dt, doublet of triplets, tt, triplet of triplets, m, multiplet and br, broad. Coupling constants (J) were recorded in Hertz (Hz) to the nearest 0.5Hz. Carbon spectra are supported by DEPT analysis where necessary. Infrared (IR) spectra were recorded as thin films between NaCl plates. Absorption maximum (ν_{max}) was reported in wave numbers (cm^{-1}) and only selected peaks are reported. The following abbreviations are used: w, weak, m, medium, s, strong and br, broad.

Flash chromatography was carried out using *silica gel 60* (0.040-0.063mm, 230-400 mesh) as the stationary phase. Thin layer chromatography was carried out on aluminium backed plates pre-coated with *silica gel 60*, which were visualized by quenching of u.v. fluorescence ($\lambda_{\text{max}} = 254 \text{ nm}$) or by staining with either 10% w/v ammonium molybdate in 2M sulphuric acid or basic potassium permanganate solution (followed by heat) as appropriate. Retention factors (R_f) are reported to ± 0.5 .

Spectroscopic data for compounds 5d-p and 3d-m:

8-Acetyl-3-methyl-4-nitro-9-(4-nitro-phenyl)-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5d: colourless solid (932 mg, 83% yield) m.p. 181-2 °C (ethanol); $\nu_{\max}$ (Film)/cm⁻¹: 1720 s, 1560s; ¹H-NMR δ : (200 MHz, CD₃COCD₃) 7.96 (2H, d, J = 8), 7.48 (2H, d, J = 8), 5.80 (1H, s), 4.35 (1H, d, J = 12), 3.70 (1H, td, J = 12, J = 4), 3.05 (1H, d, J = 14), 2.53-2.43 (2H, m), 1.90 (3H, s), 1.81 (3H, s), 1.74-1.65 (1H, m); ¹³C-NMR δ : (80MHz, CD₃COCD₃) 203.7, 201.9, 200.8, 153.3, 153.1, 146.5, 146.4, 139.5, 138.9, 128.8, 128.1, 123.7, 123.3, 97.7, 97.5, 88.2, 66.6, 66.1, 48.7, 44.6, 41.1, 40.7, 40.5, 35.6, 29.5, 10.3. HRMS found: MH⁺ 376.1143, C₁₇H₁₈N₃O₇ requires 376.1145; m/z : 376 (85%, MH⁺).

3-Methyl-4-nitro-7-(4-nitro-phenyl)-9-oxo-1-oxa-2-aza-spiro[4.5]dec-2-ene-8-carboxylic acid ethyl ester 5e: colorless solid (862 mg, 71% yield), m.p. 152-3 °C (ethanol), $\nu_{\max}$ (Film)/cm⁻¹: 1731s, 1570s; ¹H-NMR δ : (200 MHz, CDCl₃) 8.20 (2H, d, J = 8), 7.47 (2H, d, J = 8), 5.39 (1H, s), 4.20-3.90 (3H, m), 3.74 (1H, d, J = 14), 2.86 (1H, d, J = 16), 2.76 (1H, d, J = 16), 2.31-2.08 (2H, m), 2.17 (3H, s), 1.13 (3H, t, J = 7); ¹³C-NMR δ : (80MHz, CD₃COCD₃) 197.2, 166.9, 152.2, 151.4, 146.1, 145.2, 138.8, 138.1, 128.9, 123.5, 122.1, 97.6, 97.3, 88.1, 62.1, 61.8, 59.2, 48.0, 44.2, 42.5, 42.1, 41.8, 37.8, 12.5, 10.1; HRMS found: MH⁺ 406.1249, C₁₈H₂₀N₃O₈ requires 406.1250; m/z : 406 (100%, MH⁺).

3-Methyl-4-nitro-9-(4-nitro-phenyl)-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5f: colourless solid (779 mg, 78% yield), m.p. 148-9 °C (ethanol), $\nu_{\max}$ (Film)/cm⁻¹: 1715s, 1575s; ¹H-NMR δ : (200 MHz, CDCl₃) 7.99 (2H, d, J = 8), 7.49 (2H, d, J = 8), 5.78 (1H, s), 3.37 (1H, td, J = 14, J = 4), 2.94-2.66 (2H, m), 2.41-2.04 (3H, m), 1.89 (3H, s) 1.84-1.62 (1H, m); ¹³C-NMR δ : (80Mhz, CD₃COCD₃) 202.3, 202.1, 150.3, 150.2, 146.7, 141.2, 127.9, 127.8, 123.4, 123.2, 98.1, 97.7, 88.5, 48.2, 46.0, 45.6, 43.2, 39.3, 38.3, 38.1, 36.2, 10.7. HRMS found: MH⁺ 334.1038, C₁₅H₁₆N₃O₆ requires 334.1039; m/z : 334 (100%, MH⁺).

8-Acetyl-9-(4-chloro-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5g: colourless solid (873 mg, 80% yield), m.p. 184-5 °C (ethanol); ν_{max} (Film)/cm⁻¹: 1725s, 1560s; ¹H-NMR δ : (200 MHz, CDCl₃) 7.29 (2H, d, J = 8), 7.15 (2H, d, J = 8), 5.36 (1H, s), 3.85 (1H, td, J = 12, J = 4), 3.75 (1H, d, J = 12), 2.84-2.75 (2H, m), 2.26-2.01 (2H, m), 2.16 (3H, s), 2.01 (3H, s); ¹³C-NMR δ : (80MHz, CD₃COCD₃) 202.8, 201.9, 201.8, 150.4, 150.2, 140.2, 140.1, 132.0, 131.9, 128.8, 128.7, 128.3, 97.8, 97.4, 88.0, 66.3, 65.9, 48.6, 44.6, 40.6, 40.3, 40.1, 35.9, 29.6, 10.7. HRMS found: MH⁺ 365.0902, C₁₇H₁₈ClN₂O₅ requires 365.0904; m/z : 365 (100%, MH⁺).

7-(4-Chloro-phenyl)-3-methyl-4-nitro-9-oxo-1-oxa-2-aza-spiro[4.5]dec-2-ene-8-carboxylic acid ethyl ester 5h: colourless solid (815 mg, 69% yield), m.p. 151-2°C (ethanol), ν_{max} (Film)/cm⁻¹: 1728s, 1576s; ¹H-NMR δ : (200 MHz, CDCl₃) 7.07 (2H, d, J = 8), 6.95 (2H, d, J = 8), 5.14 (1H, s), 3.89-3.76 (2H, m), 3.76-3.53 (1H, m), 3.43 (1H, d, J = 13), 2.58 (1H, d, J = 15), 2.48 (1H, dd, J = 15, J = 2), 2.05-2.91 (2H, m), 1.91 (3H, s), 0.86 (3H, t, J = 7); ¹³C-NMR δ : (80MHz, CDCl₃) 197.0, 196.6, 166.0, 148.8, 148.2, 137.4, 132.6, 128.0, 127.5, 97.6, 97.3, 86.9, 61.5, 61.3, 60.2, 48.0, 44.3, 39.9, 39.8, 39.7, 34.9, 12.8, 10.9. HRMS found: MH⁺ 395.1009, C₁₈H₂₀ClN₂O₆ requires 395.1001; m/z : 395 (80%, MH⁺).

9-(4-Chloro-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5i: colourless solid (830 mg, 86% yield), m.p. 164-5 °C (ethanol), ν_{max} (Film)/cm⁻¹: 1720s, 1572s; ¹H-NMR δ : (200 MHz, CD₃COCD₃) 7.22-7.11 (4H, m), 5.75 (1H, s), 3.18 (1H, tt, J = 12, J = 4), 2.87 (1H, d, J = 15), 2.67 (1H, d, J = 14), 2.34-2.08 (3H, m), 1.89 (3H, s) 1.83-1.85-1.65 (1H, m); ¹³C-NMR δ : (80MHz, CD₃COCD₃) 203.5, 202.7, 150.3, 150.0, 141.7, 141.6, 131.8, 131.7, 128.3, 128.2, 128.1, 98.2, 97.8, 88.6, 48.2, 46.6, 46.1, 44.1, 39.9, 37.9, 37.7, 35.0, 10.7. HRMS found: MH⁺ 323.0798, C₁₅H₁₆ClN₂O₄ requires 323.0799; m/z : 323 (80%, MH⁺).

8-Acetyl-9-(2-chloro-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5k: colourless solid (709 mg, 65% yield), m.p. 142-3 °C (ethanol); $\nu_{\max}$ (Film)/cm⁻¹: 1725s, 1570s; ¹H-NMR δ : (200 MHz, CD₃COCD₃) 7.23-7.06 (4H, m), 5.60 (1H, s), 4.42-4.32 (1H, m), 3.00 (1H, d, J = 15), 2.29-2.12 (2H, m), 1.90 (3H, s), 1.65-1.46 (1H, m), 1.43 (3H, s); ¹³C-NMR δ : (80MHz, CD₃COCD₃) 198.9, 178.1, 150.1, 142.2, 132.1, 129.3, 128.4, 127.9, 127.8, 109.0, 97.6, 85.1, 38.2, 35.0, 24.6, 17.5, 10.7; HRMS found: MH⁺ 365.0905, C₁₇H₁₈ClN₂O₅ requires 365.0904; m/z : 365 (100%, MH⁺).

7-(2-Chloro-phenyl)-3-methyl-4-nitro-9-oxo-1-oxa-2-aza-spiro[4.5]dec-2-ene-8-carboxylic acid ethyl ester 5l: colourless solid (697 mg, 59% yield), m.p. 131-2 °C (ethanol); $\nu_{\max}$ (Film)/cm⁻¹: 1731s, 1570s; ¹H-NMR δ : (200 MHz, CDCl₃) 7.40-7.18 (4H, m), 5.35 (1H, s), 4.60-4.23 (1H, m), 4.20-3.82 (3H, m), 2.83 (1H, d, J = 14), 2.73 (1H, dd, J = 14, J = 2), 2.39-2.12 (2H, m), 2.14 (3H, s), 1.06 (3H, t, J = 7); ¹³C-NMR δ : (80MHz, CDCl₃) 199.1, 167.1, 150.2, 150.1, 139.4, 132.4, 129.3, 128.2, 128.1, 127.5, 127.3, 127.1, 127.0, 97.9, 96.7, 88.4, 88.3, 61.8, 61.5, 59.2, 48.2, 43.2, 41.9, 41.5, 40.1, 35.6, 13.1, 10.7; HRMS found: MH⁺ 395.1002, C₁₈H₂₀ClN₂O₆ requires 395.1001; m/z : 395 (100%, MH⁺).

9-(2-Chloro-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5m: colourless solid (656 mg, 68% yield), m.p. 125-6 °C (ethanol), $\nu_{\max}$ (Film)/cm⁻¹: 1702s, 1570s; ¹H-NMR δ : (200 MHz, CDCl₃) 7.32 (1H, d, J = 7), 7.23- 7.05 (3H, m), 5.76 (1H, s), 3.70 (1H, tt, J = 12, J = 4), 2.90 (1H, d, J = 14), 2.72 (1H, t, J = 14), 2.40-2.06 (3H, m), 1.90 (3H, s), 1.89-1.65 (1H, m); ¹³C-NMR δ : (80MHz, CD₃COCD₃) 202.8, 202.7, 150.2, 150.1, 139.5, 132.6, 129.3, 128.1, 128.0, 127.3, 127.2, 127.1, 127.0, 97.9, 96.7, 88.5, 88.4, 48.4, 45.0, 44.6, 43.5, 38.6, 34.3, 34.2, 33.7, 10.8. HRMS found: MH⁺ 323.0798, C₁₅H₁₆ClN₂O₄ requires 323.0799; m/z : 323 (80%, MH⁺).

8-Acetyl-9-(4-methoxy-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5n: colourless solid (918 mg, 85% yield), m.p. 168-9 °C (ethanol); $\nu_{\max}$ (Film)/cm⁻¹: 1721s, 1570s; ¹H-

NMR δ : (200 MHz, CDCl_3) 7.10 (2H, d, $J = 8$), 6.80 (2H, d, $J = 8$), 5.41 (1H, s), 3.80 (1H, d, $J = 15$), 3.75-3.68 (4H, m), 2.74 (2H, m), 2.28-2.05 (2H, m), 2.12 (3H, s), 2.00 (3H, s); ^{13}C -NMR δ : (80MHz, CDCl_3) 202.9, 202.4, 202.2, 158.5, 158.4, 150.4, 150.2, 132.9, 132.8, 128.0, 127.9, 113.7, 113.6, 97.8, 97.4, 88.1, 66.8, 66.4, 54.1, 48.5, 44.6, 40.8, 40.4, 40.3, 36.0, 29.9, 10.7. HRMS found: MH^+ 361.1400, $\text{C}_{18}\text{H}_{21}\text{N}_2\text{O}_6$ requires 361.1399; m/z : 361 (100%, MH^+).

7-(4-Methoxy-phenyl)-3-methyl-4-nitro-9-oxo-1-oxa-2-aza-spiro[4.5]dec-2-ene-8-carboxylic acid ethyl ester 5o: colourless solid (842 mg, 72% yield), m.p. 151-2 °C (ethanol); ν_{max} (Film)/ cm^{-1} : 1722s, 1572s; ^1H -NMR δ : (200 MHz, CDCl_3) 7.15 (2H, d, $J = 8$), 6.85 (2H, d, $J = 8$), 5.35 (1H, s), 4.20-3.98 (2H, m), 3.90-3.73 (1H, m), 3.78 (3H, s), 3.65 (1H, d, $J = 13$), 2.79 (1H, d, $J = 15$), 2.69 (1H, d, $J = 15$), 2.28-2.07 (2H, m), 2.15 (3H, s), 1.08 (3H, t, $J = 7$); ^{13}C -NMR δ : (80MHz, CDCl_3) 197.9, 166.5, 157.9, 148.4, 131.0, 127.2, 127.1, 113.3, 97.5, 97.3, 87.2, 62.0, 61.8, 60.0, 54.2, 48.0, 44.4, 40.4, 39.9, 39.7, 35.1, 12.9, 10.7. HRMS found: MH^+ 391.1506, $\text{C}_{19}\text{H}_{23}\text{N}_2\text{O}_7$ requires 391.1505; m/z : 391 (100%, MH^+).

9-(4-Methoxy-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5p: colourless solid (735 mg, 77% yield), m.p. 146-7 °C (ethanol); ν_{max} (Film)/ cm^{-1} : 1709s, 1570s; ^1H -NMR δ : (200 MHz, CDCl_3) 7.15 (2H, d, $J = 8$), 6.88 (2H, d, $J = 8$), 5.38 (1H, s), 3.80 (3H, s), 3.46 (1H, tt, $J = 12$, $J = 4$), 2.76-2.47 (4H, m), 2.25-2.01 (2H, m), 2.13 (3H, s); ^{13}C -NMR δ : (80MHz, CDCl_3) 203.4, 202.7, 158.3, 150.7, 150.2, 134.7, 134.4, 127.4, 127.3, 113.7, 113.6, 98.0, 97.7, 88.7, 54.1, 48.2, 47.1, 46.6, 43.4, 40.4, 37.8, 37.6, 35.6, 10.8. HRMS found: MH^+ 319.1295, $\text{C}_{16}\text{H}_{19}\text{N}_2\text{O}_5$ requires 319.1294; m/z : 319 (90%, MH^+).

3-[2-(3-Methyl-4-nitro-isoxazol-5-yl)-1-(4-nitro-phenyl)-ethyl]-pentane-2,4-dione 3d: colourless solid (920 mg, 82% yield), $R_f = 0.2$ (ethyl acetate : petroleum ether 1:3), m.p. 131-2 °C (ethanol); ν_{max} (Film)/ cm^{-1} : 1710s, 1600s, 1560m; ^1H -NMR δ : (400 MHz, CDCl_3) 8.15 (2H, $J = 8$), 7.37 (2H, $J = 8$),

4.35-4.28 (2H, m), 3.52 (1H, dd, $J = 14, J = 9$), 3.35 (1H, dd, $J = 14, J = 5$), 2.49 (3H, s), 2.36 (3H, s), 1.95 (3H, s); ^{13}C -NMR δ : (100MHz, CDCl_3) 200.7, 200.3, 170.4, 155.2, 147.0, 145.3, 130.9, 128.4, 123.8, 73.7, 41.7, 31.4, 29.3, 29.1, 11.5; HRMS found: MH^+ 376.1146, $\text{C}_{17}\text{H}_{18}\text{N}_3\text{O}_7$ requires 376.1145; m/z : 376 (100%, MH^+).

2-Acetyl-4-(3-methyl-4-nitro-isoxazol-5-yl)-3-(4-nitro-phenyl)-butyric acid ethyl ester 3e:

colourless solid (875 mg, 74% yield), $R_f = 0.4$ (ethyl acetate : petroleum ether 1:3), m.p. 95-8 °C (ethanol); ν_{max} (Film)/ cm^{-1} : 1725s, 1600s, 1560m; ^1H -NMR δ : (200 MHz, CDCl_3) 8.01 (2H, d, $J = 8$), 7.37 (2H, d, $J = 8$), 4.31-3.81 (4H, m), 3.74-3.46 (2H, m), 2.44 (3H, s), 2.34 (1.5H, s), 2.06 (1.5H, s), 1.27 (1.5H, t, $J = 7$), 0.98 (1.5H, t, $J = 7$); ^{13}C -NMR δ : (80MHz, CDCl_3) 198.9, 198.6, 170.2, 166.0, 165.5, 154.4, 154.3, 146.3, 145.3, 129.4, 127.9, 122.8, 63.5, 63.2, 61.3, 60.8, 30.5, 30.1, 28.9, 28.3, 12.8, 12.5, 10.2; HRMS found: MH^+ 406.1248, $\text{C}_{18}\text{H}_{20}\text{N}_3\text{O}_8$ requires 406.1250; m/z : 406 (100%, MH^+).

5-(3-Methyl-4-nitro-isoxazol-5-yl)-4-(4-nitro-phenyl)-pentan-2-one 3f:

colourless oil (798 mg, 80% yield), $R_f = 0.4$ (ethyl acetate : petroleum ether 1:4); ν_{max} (Film)/ cm^{-1} : 1710s, 1601s, 1580s; ^1H -NMR δ : (400 MHz, CDCl_3) 8.17 (2H, d, $J = 8$), 7.42 (2H, d, $J = 8$), 4.03 (1H, quintet, $J = 7$), 3.66 (1H, dd, $J = 14, J = 7$), 3.53 (1H, dd, $J = 14, J = 7$), 2.99 (1H, dd, $J = 14, J = 4$), 2.95 (1H, dd, $J = 14, J = 4$), 2.52 (3H, s), 2.14 (3H, s); ^{13}C -NMR δ : (100MHz, CDCl_3) 204.4, 171.4, 155.2, 148.7, 146.2, 130.2, 127.8, 123.7, 48.2, 37.7, 32.8, 29.8, 11.1; HRMS found: MH^+ 334.1037, $\text{C}_{15}\text{H}_{16}\text{N}_3\text{O}_6$ requires 334.1039; m/z : 334 (100%, MH^+).

3-[1-(4-Chloro-phenyl)-2-(3-methyl-4-nitro-isoxazol-5-yl)-ethyl]-pentane-2,4-dione 3g:

colourless solid (775 mg, 71% yield), $R_f = 0.3$ (ethyl acetate : petroleum ether 1:4), m.p. 89-90 °C (ethanol); ν_{max} (Film)/ cm^{-1} : 1705 s, 1601s, 1520m, ^1H -NMR δ : (400 MHz, CDCl_3) 7.22 (2H, $J = 8$), 6.98 (2H, $J = 8$), 4.26 (1H, d, $J = 11$), 4.18-4.12 (1H, m), 3.52 (1H, dd, $J = 14, J = 9$), 3.35 (1H, dd, $J = 14, J = 5$), 2.48

(3H, s), 2.32 (3H, s), 1.90 (3H, s); ^{13}C -NMR δ : (100MHz, CDCl_3) 201.7, 201.3, 171.0, 155.0, 136.2, 133.0, 131.0, 129.3, 129.0, 74.1, 41.5, 31.7, 29.6, 29.1, 11.1; HRMS found: MH^+ 365.0906, $\text{C}_{17}\text{H}_{18}\text{ClN}_2\text{O}_5$ requires 365.0904; m/z : 365 (100%, MH^+).

2-Acetyl-3-(4-chloro-phenyl)-4-(3-methyl-4-nitro-isoxazol-5-yl)-butyric acid ethyl ester 3h:

colourless solid (920 mg, 78% yield), R_f = 0.3 (ethyl acetate : petroleum ether 1:4), m.p. 94-5 °C (ethanol); IR: 1742s, 1601s, 1562s; ^1H -NMR δ : (200MHz, CDCl_3) 7.21 (2H, J = 8), 7.10 (2H, J = 8), 4.09-3.94 (2H, m), 3.92 (2H, q, J = 7), 3.62 (1H, dd, J = 14, J = 9), 3.45 (1H, dd, J = 14, J = 5), 2.46 (3H, s), 2.32 (3H, s), 0.99 (3H, t, J = 7); ^{13}C -NMR δ : (80MHz, CDCl_3) 199.6, 170.8, 165.5, 154.4, 136.0, 132.5, 130.5, 128.2, 127.8, 64.2, 60.7, 40.3, 30.8, 28.4, 16.6, 11.4; HRMS found: MH^+ 395.1009, $\text{C}_{18}\text{H}_{20}\text{ClN}_2\text{O}_6$ requires 395.1001; m/z : 395 (70%, MH^+).

4-(4-Chloro-phenyl)-5-(3-methyl-4-nitro-isoxazol-5-yl)-pentan-2-one 3i: colourless oil (782 mg, 81% yield), R_f = 0.3 (ethyl acetate : petroleum ether 1:5): ν_{max} (Film)/ cm^{-1} : 1715s, 1605s, 1565s; ^1H -NMR δ : (400 MHz, CDCl_3) 7.25 (2H, d, J = 8), 7.15 (2H, d, J = 8), 3.82 (1H, quintet, J = 7), 3.57 (1H, dd, J = 14, J = 7), 3.44 (1H, dd, J = 14, J = 7), 2.89 (2H, app. septet, J = 17, J = 7), 2.50 (3H, s), 2.10 (3H, s); ^{13}C -NMR δ : (100MHz, CDCl_3) 205.1, 172.1, 155.1, 139.7, 132.6, 130.1, 128.55, 128.1, 48.6, 37.4, 33.2, 30.0, 11.2; HRMS found: MH^+ 323.0801, $\text{C}_{15}\text{H}_{16}\text{ClN}_2\text{O}_4$ requires 323.0799; m/z : 323 (100%, MH^+).

3-[1-(2-Chloro-phenyl)-2-(3-methyl-4-nitro-isoxazol-5-yl)-ethyl]-pentane-2,4-dione 3k: colourless solid (677 mg, 62% yield), R_f = 0.3 (ethyl acetate : petroleum ether 1:3), m.p. 85-6 °C (ethanol), ν_{max} (Film)/ cm^{-1} : 1705s, 1605s, 1535m; ^1H -NMR δ : (400 MHz, CDCl_3) 7.34-7.19 (4H, m), 4.85-4.60 (1H, m), 4.45-4.39 (1H, m), 3.59-3.48 (2H, m), 2.48 (3H, s), 2.33 (3H, s), 1.94 (3H, s); ^{13}C -NMR δ : (100MHz, CDCl_3) 202.1, 201.7, 172.1, 155.4, 138.4, 133.2, 130.6, 129.8, 128.5, 127.1, 126.5, 75.1,

42.6, 32.0, 29.5, 29.3, 11.4; HRMS found: MH^+ 365.0905, $\text{C}_{17}\text{H}_{18}\text{ClN}_2\text{O}_5$ requires 365.0904; m/z : 365 (100%, MH^+).

2-Acetyl-3-(2-chloro-phenyl)-4-(3-methyl-4-nitro-isoxazol-5-yl)-butyric acid ethyl ester 3l:

colourless solid (614 mg, 52% yield), $R_f = 0.3$ (ethyl acetate : petroleum ether 1:3), m.p. 64-5 °C (ethanol), ν_{max} (Film)/ cm^{-1} : 1720, 1601, 1525; $^1\text{H-NMR}$ δ : (200 MHz, CDCl_3) 7.32-7.15 (5H, m), 4.73-4.52 (1H, m), 4.32-4.09 (2H, m), 3.92 (1H, q, $J = 7$), 3.80-3.54 (2H, m), 2.46 (3H, s), 2.35 (1.6H, s), 2.11 (1.4H, s), 1.30 (1.4H, t, $J = 7$), 0.96 (1.6H, t, $J = 7$); $^{13}\text{C-NMR}$ δ : (80MHz, CDCl_3) 199.5, 199.2, 170.7, 166.6, 165.7, 154.2, 135.2, 134.8, 133.1, 129.5, 130.0, 127.9, 127.8, 126.2, 125.9, 62.7, 62.2, 60.9, 60.6, 29.5, 29.1, 28.3, 28.0, 12.9, 12.4, 10.3; HRMS found: MH^+ 395.1009, $\text{C}_{18}\text{H}_{20}\text{ClN}_2\text{O}_6$ requires 395.1001; m/z : 395 (100%, MH^+).

4-(2-Chloro-phenyl)-5-(3-methyl-4-nitro-isoxazol-5-yl)-pentan-2-one 3m: colourless oil (695 mg, 72% yield), $R_f = 0.3$ (ethyl acetate : petroleum ether 1:4), ν_{max} (Film)/ cm^{-1} : 1715s, 1600s, 1565s; $^1\text{H-NMR}$ δ : (400 MHz, CDCl_3) 7.34 (1H, d, $J = 8$), 7.25-7.15 (3H, m), 4.38 (1H, quintet, $J = 7$), 3.60 (2H, d, $J = 7$), 2.96 (2H, d, $J = 6$), 2.51 (3H, s), 2.15 (3H, s); $^{13}\text{C-NMR}$ δ : (100MHz, CDCl_3) 205.2, 172.0, 155.0, 138.2, 133.2, 130.4, 129.7, 128.0, 127.1, 126.8, 47.4, 34.9, 32.0, 30.0, 11.6; HRMS found: MH^+ 323.0797, $\text{C}_{15}\text{H}_{16}\text{ClN}_2\text{O}_4$ requires 323.0798; m/z : 323 (100%, MH^+).

8-Acetyl-3-methyl-4-nitro-9-phenyl-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5a

1

t=0

2

STANDARD 1H OBSERVE

Pulse Sequence: zgpg30
Solvent: Acetone
Ambient temperature
GEMINI-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec
Pulse 40.7 degrees
Acq. time 3.744 sec
Width 3000.0 Hz
16 repetitions
OBSERVE N H1, 199.9707954 MHz
DATA PROCESSING
Line broadening 1.0 Hz
Fw size 32768
Total time 0 min, 18 sec

6-30"

3

STANDARD 1H OBSERVE

Pulse Sequence: zgpg30

Solvent: Acetone

Ambient temperature

GENI-200 "gemini200"

PULSE PROGRAM

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707954 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

(- = 120°)

1

STANDARD 1H OBSERVE

Pulse Sequence: s2pul

Solvent: Acetone

Ambient temperature

GEVINI-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707952 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

C-24.0"

12

STANDARD 1H OBSERVE

Pulse Sequence: zgpg1
Solvent: Acetone
Ambient temperature
GEMINI-200 "gemin1200"

PULSE SEQUENCE
Relax. delay 1.000 sec
Pulse 40.7 degrees
Acq. time 3.744 sec
Width 3000.0 Hz
16 repetitions
OBSERVE H1, 199.9707937 MHz
DATA PROCESSING
Line broadening 1.0 Hz
Fw size 32768
Total time 0 min, 18 sec

180°

6

STANDARD 1H OBSERVE

Pulse Sequence: zgpg30

Solvent: Acetone

Ambient temperature

GE4000-200 "gemin4200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

6 repetitions

OBSERVE H1, 199.9707954 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

6.0

2

STANDARD 1H OBSERVE

Pulse Sequence: zgpg30

Solvent: Acetone

Ambient temperature

gpg30-200 "gpg30-200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707954 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

C-3011

8

STANDARD 1H OBSERVE

Pulse Sequence: zgpg1

Solvent: Acetone

Ambient temperature

GEMINI-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707954 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

$\epsilon = 120''$

9

STANDARD 1H OBSERVE

Pulse Sequence: zgpg1

Solvent: Acetone

Ambient temperature

GEMINI-200 "gemin1200"

RELAX. DELAY 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707952 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

$t = 240''$

10

STANDARD 1H OBSERVE

Pulse Sequence: zgpg1

Solvent: Acetone

Ambient temperature

GE1200-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707937 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

4.80"

11

STANDARD 1H OBSERVE

Pulse Sequence: zgpg1

Solvent: Acetone

Ambient temperature

GEKINI-200 "gemini200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

6 repetitions

OBSERVE H1, 199.9707954 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

t=0

12

STANDARD 1H OBSERVE

Pulse Sequence: zgpg1

Solvent: Acetone

Ambient temperature

GEMINI-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707954 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

$\tau = 30$

13

STANDARD 1H OBSERVE

Pulse Sequence: zgpg1

Solvent: Acetone

Ambient temperature

GEVINI-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707954 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

$t = 120''$

(14)

STANDARD 1H OBSERVE

Pulse Sequence: zgpg1

Solvent: Acetone

Ambient temperature

GEMINI-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707952 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

$\delta = 240$ "

15

STANDARD 1H OBSERVE

Pulse Sequence: zgpg30

Solvent: Acetone

Ambient temperature

GERMINI-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707937 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

480 "

16

STANDARD 1H OBSERVE

Pulse Sequence: szpul

Solvent: Acetone

Ambient temperature

GEMINI-200 "gemin1200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

6 repetitions

OBSERVE H1, 199.9707954 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

6-0

17

STANDARD 1H OBSERVE

Pulse Sequence: s2pul

Solvent: Acetone

Ambient temperature

GEVINI-200 "gemin200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9707952 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

C-240"

3-Methyl-4-nitro-7-oxo-9-phenyl-1-oxa-2-aza-spiro[4.5]dec-2-ene-8-carboxylic acid ethyl ester 5b

3-Methyl-4-nitro-9-phenyl-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5c

8-Acetyl-3-methyl-4-nitro-9-(4-nitro-phenyl)-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5d

3-Methyl-4-nitro-7-(4-nitro-phenyl)-9-oxo-1-oxa-2-aza-spiro[4.5]dec-2-ene-8-carboxylic acid ethyl ester 5e

3-Methyl-4-nitro-9-(4-nitro-phenyl)-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5f

8-Acetyl-9-(4-chloro-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5g

7-(4-Chloro-phenyl)-3-methyl-4-nitro-9-oxo-1-oxa-2-aza-spiro[4.5]dec-2-ene-8-carboxylic acid ethyl ester 5h

9-(4-Chloro-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5i

8-Acetyl-9-(2-chloro-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5k

9-(2-Chloro-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5m

8-Acetyl-9-(4-methoxy-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5n

7-(4-Methoxy-phenyl)-3-methyl-4-nitro-9-oxo-1-oxa-2-aza-spiro[4.5]dec-2-ene-8-carboxylic acid ethyl ester 50

9-(4-Methoxy-phenyl)-3-methyl-4-nitro-1-oxa-2-aza-spiro[4.5]dec-2-en-7-one 5p

3-[2-(3-Methyl-4-nitro-isoxazol-5-yl)-1-phenyl-ethyl]-pentane-2,4-dione 3a

ED 4.9

Pulse Sequence: s2pul

Solvent: CDCl₃

Ambient temperature

GEMINI-200 "gemin200"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 40.7 degrees

Acq. time 3.744 sec

Width 3000.0 Hz

16 repetitions

OBSERVE H1, 199.9450247 MHz

DATA PROCESSING

Line broadening 1.0 Hz

FT size 32768

Total time 0 min, 18 sec

2-Acetyl-4-(3-methyl-4-nitro-isoxazol-5-yl)-3-phenyl-butyric acid ethyl ester 3b

5-(3-Methyl-4-nitro-isoxazol-5-yl)-4-phenyl-pentan-2-one 3c

column of ether layer

*** Current Data Parameters ***
 NAME : 56el3
 EXPNO : 1
 PROCNO : 1
 *** Acquisition Parameters ***
 DATED : Jan 16 2004
 NS : 16
 NUCLEUS : off
 SF01 : 400.1352017 MHz
 *** 1D NMR Plot Parameters ***
 NUCLEUS : off

3-[2-(3-Methyl-4-nitro-isoxazol-5-yl)-1-(4-nitro-phenyl)-ethyl]-pentane-2,4-dione 3d

*** Current Data Parameters ***

NAME	:	71e2y
EXPNO	:	1
PROCNO	:	1

*** Acquisition Parameters ***

DATE	:	Apr 14 2004
NS	:	16
NUCLEUS	:	off
SFO1	:	400.1336012 MHz

*** 1D NMR Plot Parameters ***

NUCLEUS	:	off
---------	---	-----

2-Acetyl-4-(3-methyl-4-nitro-isoxazol-5-yl)-3-(4-nitro-phenyl)-butyric acid ethyl ester 3e

5-(3-Methyl-4-nitro-isoxazol-5-yl)-4-(4-nitro-phenyl)-pentan-2-one 3f

*** Current Data Parameters ***

NAME	:	76yc13
EXPNO	:	1
PROCNO	:	1

*** Acquisition Parameters ***

DATE	:	Apr 21 2004
NS	:	16
NUCLEUS	:	off
SFO1	:	400.1336012 MHz

*** 1D NMR Plot Parameters ***

NUCLEUS	:	off
---------	---	-----

3-[1-(4-Chloro-phenyl)-2-(3-methyl-4-nitro-isoxazol-5-yl)-ethyl]-pentane-2,4-dione 3g

*** Current Data Parameters ***

NAME	:	70ey
EXPNO	:	1
PROCNO	:	1

*** Acquisition Parameters ***

DATE	:	Apr 01 2004
NS	:	16
NUCLEUS	:	off
SFO1	:	400.1336012 MHz

*** 1D NMR Plot Parameters ***

NUCLEUS	:	off
---------	---	-----

2-Acetyl-3-(4-chloro-phenyl)-4-(3-methyl-4-nitro-isoxazol-5-yl)-butyric acid ethyl ester 3h

4-(4-Chloro-phenyl)-5-(3-methyl-4-nitro-isoxazol-5-yl)-pentan-2-one 3i

*** Current Data Parameters ***

NAME	:	75ebv
EXPNO	:	1
PROCNO	:	1

*** Acquisition Parameters ***

DATE	:	Apr 14 2004
NS	:	16
NUCLEUS	:	off
SFO1	:	400.1336012 MHz

*** 1D NMR Plot Parameters ***

NUCLEUS	:	off
---------	---	-----

3-[1-(2-Chloro-phenyl)-2-(3-methyl-4-nitro-isoxazol-5-yl)-ethyl]-pentane-2,4-dione 3k

*** Current Data Parameters ***

NAME	EXPNO	PROCNO
69e0x	1	1

*** Acquisition Parameters ***

DATE_1	TIME_1	DATE_2	TIME_2
Apr 06 2004	16		

NUCLEUS : off

SFO1 : 400.1336012 MHz

*** 1D NMR Plot Parameters ***

NUCLEUS : off

2-Acetyl-3-(2-chloro-phenyl)-4-(3-methyl-4-nitro-isoxazol-5-yl)-butyric acid ethyl ester 3l

4-(2-Chloro-phenyl)-5-(3-methyl-4-nitro-isoxazol-5-yl)-pentan-2-one 3m

NMR Peak Assignments for 5a and 4-*epi*-5a: δ_H (400 MHz, CD_3COCD_3) for **5a**: 7.41-7.23 (5H, m, Ph), 6.02 (1H, s, CHNO₂), 4.43 (1H, d, $J = 12$, O=CCHC=O), 3.79 (1H, td, $J = 12$, $J = 4$, CHPh), 3.24 (1H, d, $J = 14$, CH_{ax}H_{eq}C=O), 2.73 (1H, dd, CH_{ax}H_{eq}C=O, $J = 12$, $J = 2.5$) 2.63 (1H, t, $J = 12$, CH_{ax}H_{eq}CHPh), 2.02 (3H, s CH₃C=N), 1.88 (3H, s, CH₃C=O), 1.89 (1H, ddd, $J = 12$, $J = 4$, $J = 2.5$, CH_{ax}H_{eq}CHPh); δ_H (400 MHz, CD_3COCD_3) for **4-*epi*-5a**: 7.41-7.23 (5H, m, Ph), 5.88 (1H, s, CHNO₂), 4.43 (1H, d, $J = 12$, O=CCHC=O), 3.79 (1H, td, $J = 12$, $J = 4$, CHPh), 3.28 (1H, d, $J = 14$, CH_{ax}H_{eq}C=O), 2.52 (1H, t, $J = 12$, CH_{ax}H_{eq}CHPh), 2.37 (1H, dd, $J = 12$, $J = 2.5$, CH_{ax}H_{eq}C=O), 2.18 (1H, ddd, $J = 12$, $J = 4$, $J = 2.5$, CH_{ax}H_{eq}CHPh), 2.02 (3H, s CH₃C=N), 1.88 (3H, s, CH₃C=O); δ_C (80Mhz, CD_3COCD_3) 202.7 (C=O), 201.9 and 201.8 (C=O), 151.2 and 151.0 (CH₃C=N), 141.2 (Ph), 128.3 (Ph), 127.8 (Ph), 126.9 and 126.8 (Ph), 97.8 and 97.5 (CHNO₂), 88.1 (O₂NCHCO), 66.5 and 66.4 (O=CCHC=O), 48.6 and 44.6 (CH₂C=O), 41.0 and 35.9 (CH₂CHPh), 37.5 and 36.9 (CHPh), 29.8 (CH₃C=O), 10.7 (CH₃C=N).

Figure 1: coupling between protons H₁-H₆

Figure 2: ¹H-NMR chemical shifts of 5a

Figure 3: ¹H-NMR chemical shifts of 4-*epi*- 5a

^1H - ^1H correlation for compound 5a

^1H - ^1H correlation for compound 5a

^{13}C - ^1H correlation for compound 5a

Figure 4: ^{13}C -NMR chemical shifts of 4-*epi*- 5a

^{13}C - ^1H correlation for compound 5a

n.O.e. experiments for the assignment of 5a and 4-epi-5a (reference n. 9)

File 1 : C:\USERS\ASPX32\RCMASA\001001.1R
 File 2 : C:\USERS\ASPX32\RCMASA\100001.1R
 File 3 : C:\USERS\ASPX32\RCMASA\102001.1R