NaIO₄/LiBr-mediated diastereoselective dihydroxylation of olefins: a catalytic approach to Prevost – Woodward reaction

Supporting Information

*Lourdusamy Emmanuvel, Tanveer Mahammad Ali Shaikh and Arumugam sudalai**

Chemical Engineering-Process Development Division, National Chemical Laboratory,

Pashan Road, pune, India 411008

General information:

Solvents were purified and dried by standard procedures before use. petroleum ether of boiling range 60-80 °C was used. Melting points are uncorrected. Infrared spectra were recorded on a Shimadzu FTIR-8400 spectrometer. ^1H NMR and ^{13}C NMR were recorded on Bruker AC-200 NMR spectrometer. Mass spectra were obtained with a Finnegan MAT-1020 B-70 eV mass spectrometer. Elemental analysis was carried out on a Carlo Erba CHN-O analyzer.

General experimental procedure:

A mixture of olefin (3 mmol), NaIO_4 (30 mol %), LiBr (20 mol %) was taken in 25 ml R round bottomed flask and glacial acetic acid (5 ml) was added. The reaction mixture was heated at 95 °C (using oil bath) for 18 hours. The light yellow colored reaction mixture became purple in color after the completion of the reaction. The reaction mixture was then extracted with EtOAc (30 mL X 3) and the combined organic phase was washed with saturated sodium thiosulfate solution, water and aq. NaHCO_3 . The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give crude product which was subjected to basic hydrolysis without purification.

General procedure for the basic hydrolysis:

The crude product as obtained above was stirred with K_2CO_3 (1.5 equiv) in methanol (20 ml) at 25 °C for 24 hours. After completion of the reaction, methanol was removed under reduced pressure and the reaction mixture was extracted with EtOAc (30 mL X 3). The combined organic phase was washed with water and brine. The organic layer was then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give crude diol, which was purified by column chromatography packed with silica gel using pet ether and EtOAc (7:3) as eluents to afford pure diol.

General procedure for the acetylation of polyols: (5d and 5o-5r)

To the crude polyol was added acetic anhydride (1 mL) and pyridine (3 mL) and stirred at 90 °C for 10 hours. After completion of the reaction it was extracted with EtOAc and thoroughly washed with water and brine successively. Then the organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give crude

poly acetate, which was purified by column chromatography packed with silica gel using pet ether and EtOAc (9:1) as eluents to afford pure acetate derivatives.

1-Phenyl-1,2-ethanediol (5a)

Yield: 87%. m.p.: 64 - 65 °C. ^1H NMR (200 MHz, CDCl_3): δ 3.57 (m, 2H), 4.51(br, s, 2H), 4.69 (dd, J = 3.8 & 8.2 Hz, 1H), 7.24 (s, 5H). ^{13}C NMR (200 MHz, CDCl_3): δ 59.02, 73.33, 125.15, 126.08, 126.96, and 140.93. Analysis: $\text{C}_8\text{H}_{10}\text{O}_2$ requires C: 69.54, H: 7.31, found C: 69.53, H: 7.29 %.

1-(4-Methylphenyl)-1,2 ethanediol (5b):

Yield: 89%. m.p.: 76 - 77 °C. ^1H NMR (200 MHz, CDCl_3): δ 2.22 (s, 3H), 3.46 (m, 2H), 4.58 (dd, J = 4.1 & 8.0 Hz, 1H), 5.05(br, s, 2H), 6.97(d, J = 7.9, 2H), 7.05(d, J = 8.1, 2H). ^{13}C NMR (200 MHz, CDCl_3): δ 21.19, 67.98, 74.59, 126.11, 129.05, 137.16, 137.6. Analysis: $\text{C}_9\text{H}_{12}\text{O}_2$ requires C: 71.02, H: 7.96, found C: 70.96, H: 8.09 %.

1-(4-Bromophenyl)-1,2ethanediol (5c):

Yield: 90%. m.p.: 81 °C. ^1H NMR (200 MHz, CDCl_3 and DMSO-d_6): δ 3.60 (m, 2H), 4.72(dd, J = 3.8 & 8.2 Hz, 1H), 7.25(d, J = 8.34 Hz, 2H), 7.46(d, J = 8.5 Hz, 2H). ^{13}C NMR (200 MHz, CDCl_3 and DMSO-d_6): δ 82.67, 88.59, 135.29, 143.83, 146.0, 157.93. Analysis: $\text{C}_8\text{H}_9\text{O}_2\text{Br}$ requires C: 44.27, H: 4.19, Br: 36.81 found C: 44.26, H: 4.16, Br: 36.79 %.

1-(4-Acetoxyphenyl)-1,2- diacetoxyethane (5d)

Yield: 78%. m.p.: gum. ^1H NMR (200 MHz, CDCl_3): δ 2.05 (s, 3H), 2.10 (s, 3H), 2.28 (s, 3H) 4.27 (m, 2H), 6.0 (dd, $J = 4.0$ & 8.0 Hz). ^{13}C NMR (200 MHz, CDCl_3): 19.0, 19.32, 64.36, 71.13, 120.38, 126.41, 132.64, 149.36, 167.33, 168.12, 168.70. Analysis: $\text{C}_{12}\text{H}_{14}\text{O}_5$ requires C: 60.50, H: 5.94, found C: 60.58, H: 5.79 %.

1-Phenyl-1,2-propane-cis-diol (5e):

Yield: 84%. m.p.: gum. ^1H NMR (200 MHz, CDCl_3): δ 0.91 (d, $J = 6.3$ Hz, 3H), 3.74 (m, 1H), 4.10 (br, s, 2H), 4.2(d, $J = 7.9$ Hz, 1H), 7.25 (s, 5H). ^{13}C NMR (200 MHz, CDCl_3): δ 18.82, 72.17, 79.57, 127.10, 127.90, 128.36, 141.31 Analysis: $\text{C}_9\text{H}_{11}\text{O}_2$ requires C: 71.50, H: 7.33, found C: 71.32, H: 7.41 %.

meso-1,2-Diphenyl-1,2-ethanediol (5f & 6c)

Yield: 87 & 87%. m.p.: 98 °C. ^1H NMR (200 MHz, CDCl_3): δ 3.04 (s, 2H), 4.61 (s, 2H), 7.050 (m, 4H), 7.19 (m, 6H) ^{13}C NMR (200 MHz, CDCl_3): δ 78.64, 126.90, 127.31, 140.75 Analysis: $\text{C}_{14}\text{H}_{14}\text{O}_2$ requires C: 78.48, H: 6.58 found C: 78.40, H: 6.60 %.

1,2-Diphenyl-1,2-ethanediol (5g & 6b)

Yield: 79%. m.p.: 116 - 117 °C. ^1H NMR (200 MHz, CDCl_3): δ 2.79 (br, s, 2H), 4.61(s, 2H), 7.06 (m, 4H), 7.21 (m, 6H). ^{13}C NMR (200 MHz, CDCl_3 and DMSO-d_6): δ 78.35, 126.61, 127.02, 140.47. Analysis: $\text{C}_{14}\text{H}_{14}\text{O}_2$ requires C: 78.48, H: 6.58, found C: 78.56, H: 6.71 %.

Indane-1,2-cis-diol (5h)

Yield: 87%. m.p.: 108 - 109 °C. ^1H NMR (200 MHz, CDCl_3): δ 2.72 (br, s, 1H), 2.79 (br, s, 1H), 2.90(dd, J = 3.6 & 16.42 Hz, 1H), 3.08(dd, J = 5.7 & 16.3 Hz, 1H), 4.41(m, 1H), 4.91(d, J = 4.1 Hz, 1H), 7.24 (m, 3H), 7.38 (m, 1H). ^{13}C NMR (200 MHz, DMSO- d_6 and CDCl_3): δ 38.49, 73.03, 75.27, 124.93, 126.40, 127.76, 140.70, 143.99. Analysis: $\text{C}_9\text{H}_{10}\text{O}_2$ requires C: 71.98, H: 6.73 found C: 71.79, H: 6.65 %.

Tetrahydro naphthalene 1,2-cis-diol (5i)

Yield: 79%. m.p.: 100 - 102 °C. ^1H NMR (200 MHz, $\text{MeOH}-\text{d}_4$): δ 1.85 (m, 1H), 2.13 (m, 1H), 2.90 (m, 2H), 3.82 (m, 1H), 4.50 (d, J = 7.1 Hz, 1H), 7.13 (m, 3H), 7.55 (m, 1H). ^{13}C NMR (200 MHz, DMSO- d_6 and CDCl_3): δ 38.49, 73.03, 75.27, 124.93, 126.40, 127.76, 140.70, 143.99. Analysis: $\text{C}_{10}\text{H}_{12}\text{O}_2$ requires C: 73.14, H: 7.36 found C: 73.24, H: 7.33 %.

1-(Phenyl)-1,2,3-triacetoxy propane (5j):

Yield: 77%. m.p.: gum. ^1H NMR (200 MHz CDCl_3): δ 2.05 (s, 6H), 2.06(s, 3H), 3.76(dd, J = 4.2 & 12.0 Hz, 1H), 4.25 (dd, J = 4.03 & 12.1 Hz, 1H), 5.4 (m, 1H), 5.94(d, J = 7.9 Hz, 1H), 7.34(s, 5H) ^{13}C NMR (200 MHz, CDCl_3): δ 20.50, 20.60, 20.77, 62.04, 72.23, 73.71, 127.14, 128.66, 128.80, 169.86, 170.17, 170.86 136.42, Analysis: $\text{C}_{15}\text{H}_{18}\text{O}_6$ requires C: 61.21, H: 6.17 found C: 61.35, H: 6.01 %.

3-(Phenoxy)-1,2-propane diol (5k)

Yield: 80%. m.p.: 50 - 52 °C. ^1H NMR (200 MHz, CDCl_3): δ 3.70 (m, 2H), 3.90 (d, J = 5.31 Hz, 2H), 4.05 (m, 1H), 4.31 (br, s 2H), 6.85 (m, 3H), 7.19 (m, 2H). ^{13}C NMR (200 MHz, CDCl_3): δ 63.57, 68.82, 70.59, 114.54, 121.04, 129.37, 158.46. Analysis: $\text{C}_9\text{H}_{12}\text{O}_3$ requires C: 64.27, H: 7.21 found C: 64.20, H: 7.09 %.

3-Phenyl-3-acetoxy-2-hydroxy methyl propionate (5l)

Yield: 65%. m.p.: gum. ^1H NMR (200MHz, CDCl_3): δ 2.10(s, 3H), 3.77 (s, 3H), 4.35(d, J = 2.6 Hz, 1H), 6.03 (d, J = 2.9 Hz, 1H), 7.33 (m, 5H). ^{13}C NMR (200MHz, CDCl_3): δ 20.74, 52.67, 73.48, 75.49, 126.19, 127.02, 128.38, 136.42, 169.45, 172.30. Analysis: $\text{C}_{12}\text{H}_{14}\text{O}_5$ requires C: 60.49, H: 5.92 found C: 60.31, H: 5.77 %.

2-(4-Chlorophenyl)-propane-1,2-diol (5m)

Yield: 82%. m.p.: gum. ^1H NMR (200 MHz, CDCl_3): δ 1.45 (s, 3H), 2.69(br, s, 1H), 3.07(br, s, 1H), 3.42 (dd, J = 1.4 & 11.0 Hz, 1H) 3.66(dd, J = 1.83 & 11.84 Hz, 1H), 3.66(d, J = 11.2Hz), 7.3(m, 4H). ^{13}C NMR (200 MHz, CDCl_3): δ 36.25, 80.78, 84.72, 106.42, 137.09, 138.43, 142.86, 154.54 Analysis: $\text{C}_9\text{H}_{11}\text{ClO}_2$ requires C: 57.91, H: 5.90, Cl: 18.99, found C: 57.98, H: 5.81 %.

1-Cyclohexyl-1,2-ethane-diol (5n)

Yield: 85%. m.p.: gum. ^1H NMR (200 MHz, CDCl_3): δ 0.91-1.45, (m, 6H); 1.59-1.87, (m, 5H); 3.27 (br, s, 2H), 3.33-3.51, (m, 2H); 3.64 (dd, $J = 2.02$ & 10.3 Hz, 1H). ^{13}C NMR (200 MHz, CDCl_3): δ 26.12, 26.40, 28.65, 29.04, 40.68, 64.58, and 76.35. Analysis: $\text{C}_8\text{H}_{16}\text{O}_2$ requires C: 66.61, H: 11.18, found C: 66.78, H: 11.29 %.

1,2,4-Triacetoxy butane (5o)

Yield: 91%. m.p.: gum. ^1H NMR (200 MHz, CDCl_3): δ 1.94 (m, 2H), 2.05(s, 3H), 2.07(s, 6H), 4.09(m, 3H), 4.28(dd, $J = 3.5$ & 11.5 Hz, 1H), 5.17 (m, 1H) ^{13}C NMR (200 MHz, CDCl_3): δ 20.49, 20.73, 29.76, 59.99, 64.62, 68.40, 169.94, 170.17, 170.41. Analysis: $\text{C}_{10}\text{H}_{16}\text{O}_6$ requires C: 51.72, H: 6.94 found C: 51.66, H: 6.93 %.

1,2,3,4-Tetra acetoxy butane (5p)

Yield: 83% m.p.: gum. ^1H NMR (200 MHz, CDCl_3): δ 2.07-2.11 (m, 9H), 2.13(s, 3H), 4.21 (m, 2H), 4.32 (m, 4H). ^{13}C NMR (200 MHz, CDCl_3): δ 20.28, 20.36, 26.24, 65.08, 65.33, 68.65, 169.15, 169.36, 169.47, 169.66. Analysis: $\text{C}_{12}\text{H}_{18}\text{O}_8$ requires C: 49.65: H: 6.25, found C: 49.78, H: 6.14 %.

1,2,3-Triacetoxy propane (5q & 5r)

Yield: (5q = 86, 5r = 79 %) m.p.: gum. ^1H NMR (200 MHz, CDCl_3): δ 2.07 (s, 6H), 2.09 (s, 3H), 4.13 (dd, $J = 6.0$ & 10.0 Hz, 2H) 4.28 (dd, $J = 4.0$ & 12.0 Hz, 2H), 5.19 (m, 1H) ^{13}C NMR (200 MHz, CDCl_3): δ 20.09, 20.30, 61.76, 68.68, 169.27, 169.63. Analysis: $\text{C}_9\text{H}_{14}\text{O}_6$ requires C: 49.54, H: 6.46 found C: 49.69, H: 6.32 %.

Cyclohexane-1, 2-*cis*-diol (5s):

Yield: 86%. m.p.: 92 - 93 °C. ^1H NMR (200 MHz, CDCl_3): δ 1.30 (m, 2H), 1.61 (m, 6H), 3.73 (m, 2H), 4.05 (br, s, 2H). ^{13}C -NMR(200 MHz, CDCl_3): δ 21.58, 29.75, 70.64. Analysis: $\text{C}_6\text{H}_{12}\text{O}_2$ requires C: 62.04, H: 10.41 found C: 61.92, H: 10.36 %.

Cyclooctane-1, 2-*cis*-diol (5t):

Yield: 83%. m.p.: 75 °C. ^1H NMR (200 MHz, CDCl_3): δ 1.52-1.91 (m, 14H), 2.76 (br, s, 2H), 3.87 (m, 2H). ^{13}C NMR (200 MHz, CDCl_3): δ 23.70, 26.37, 29.96, 72.93. Analysis: $\text{C}_8\text{H}_{16}\text{O}_2$ requires C: 66.62, H: 11.18 found C: 66.78, H: 11.03 %.

***n*-Oct-1, 2-diol(5u)**

Yield: 84%. m.p.: 30 °C. ^1H NMR (200 MHz, CDCl_3): δ 0.89 (m, 3H), 1.25-1.41 (m, 10H), 3.02 (br, s, 2H), 3.39 (m, 1H), 3.62 (m, 2H). ^{13}C NMR (200 MHz, CDCl_3): δ 14.02, 22.59, 25.64, 29.42, 31.81, 33.06, 66.54, 72.23. Analysis: $\text{C}_8\text{H}_{18}\text{O}_2$ requires C: 65.71, H: 12.40, found C: 65.78, H: 12.31 %.

Indane-1, 2- *cis*-diol(6a)

Yield: 79%. m.p.: 182 - 183 °C. ^1H NMR (200 MHz, CDCl_3): δ 2.76 (dd, J = 8.08 & 15.4 Hz, 1H), 3.17 (dd, J = 7.45, & 15.4 Hz, 1H), 4.28 (m, 1H), 4.92(d, J = 6.0Hz, 1H), 7.20(m, 3H), 7.36(m, 1H). ^{13}C NMR (200 MHz, CDCl_3 and DMSO-d_6): δ 36.49, 79.03, 79.57, 122.75, 123.04, 125.01, 126.11, 137.83, 142.19. Analysis: $\text{C}_9\text{H}_{10}\text{O}_2$ requires C: 71.98, H: 6.73 found C: 71.82, H: 6.78 %.

Cyclic voltogram of NaIO₄ mediated reaction

(Dewkar, G. K.; Narina, S. V.; Sudalai, A. *Org. Lett.* **2003**, 5, 4501-4504.)

oxidation peak at $E_{pa} = 0.565$ V

reduction peaks at $E_{pc} = 0.720$ V, 0.490 V and 0.390 V

Voltage vs. Ag/AgCl reference