Nankakurine A, a Novel C₁₆N₂ Type Alkaloid from *Lycopodium hamiltonii*

Yusuke Hirasawa, Hiroshi Morita, and Jun'ichi Kobayashi*

Graduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan

Supporting Information

Figure S1. ¹H NMR spectrum of nankakurine A (**1**) in CD₃OD.

Figure S2. ¹³C NMR spectrum of nankakurine A (**1**) in CD₃OD.

Figure S3. ¹H-¹H COSY spectrum of nankakurine A (**1**) in CD₃OD.

Figure S4. HOHAHA spectrum of nankakurine A (**1**) in CD₃OD.

Figure S5. NOESY spectrum of nankakurine A (**1**) in CD₃OD.

Figure S6. HMQC spectrum of nankakurine A (**1**) in CD₃OD.

Figure S7. HMBC spectrum of nankakurine A (**1**) in CD₃OD.

Figure S1. ^1H NMR spectrum of nankakurine A (1) in CD_3OD .

Figure S2. ^{13}C NMR spectrum of nankakurine A (1) in CD_3OD .

Figure S3. ^1H - ^1H COSY spectrum of nankakurine A (**1**) in CD_3OD .

Figure S4. HOHAHA spectrum of nankakurine A (**1**) in CD₃OD.

Figure S5. NOESY spectrum of nankakurine A (**1**) in CD₃OD.

Figure S6. HMQC spectrum of nankakurine A (**1**) in CD₃OD.

Figure S7. HMBC spectrum of nankakurine A (**1**) in CD₃OD.