Supporting Information
for
Nanosized Gradient π -Conjugated Thienylethynylene Dendrimers for
Light-Harvesting: Synthesis and Properties

Jin-Liang Wang, Jia Luo, Liang-Hui Liu, Qi-Feng Zhou, Yuguo Ma,* and Jian Pei*

The Key Laboratory of Bioorganic Chemistry and Molecular Engineering and the Key Laboratory of
Polymer Chemistry and Physics of Ministry of Education, College of Chemistry, Peking University,
Beijing 100871, China

jianpei@pku.edu.cn, ygma@pku.edu.cn

Table of Contents

1. Experimental Section
2. UV-vis and PL Spectra for all compounds
3. Copies of ^1H and ^{13}C NMR spectra

1. Experimental Section

General Methods. Chemicals were purchased and used as received. All reactions were performed under a nitrogen atmosphere unless stated otherwise. Reagent grade diethyl ether and tetrahydrofuran (THF) were distilled from sodium benzophenone ketyl. ^1H and ^{13}C NMR spectra were recorded on a 200 (200 MHz) or 300 (300 MHz) and 400 (400 MHz) NMR instrument using CDCl_3 as solvent unless otherwise noted. Chemical shifts were reported in parts per million (ppm) with referencing to TMS. UV-vis spectra were recorded on PerkinElmer Lambda 35 UV-vis Spectrometer. PL spectra were carried out on PerkinElmer LS55 Luminescence Spectrometer. MALDI-TOF mass spectra were recorded on a time-of-flight (TOF) mass spectrometer using a 337 nm nitrogen laser with dithranol as matrix. Elemental analysis was performed using an elemental analyzer.

Compound 2: To a solution of 1.09 g (1.0 mmol) of **1** in 10 mL of methylene chloride was added 10 mL of acetic acid. The mixture was shielded from light and cooled in an ice bath and 740 mg (3.3 mmol) of NIS (*N*-Iodosuccinimide) was added stepwise. After 2h, the solvents were evaporated under vacuum and the residue was washed with methanol to give 1.35 g product as white powders. (Yield: 92%). ^1H NMR (CDCl_3 , 300MHz, ppm): δ 8.33-8.36 (d, $J = 8.4$ Hz, 3H, Ar-H), 7.58-7.62 (d, s, $J = 8.4$ Hz, 6H, Ar-H), 7.29-7.30 (d, $J = 3.6$ Hz, 3H, Th-H), 7.13-7.14 (d, $J = 3.6$ Hz, 3H, Th-H), 2.92-2.94 (m, 6H, CH_2), 2.10-2.11 (m, 6H, CH_2), 0.84-0.91 (m, 36H, CH_2), 0.56-0.58 (m, 30H, CH_2 , CH_3). ^{13}C NMR (CDCl_3 , 75 MHz, ppm): δ 154.4, 150.8, 145.3, 140.0, 137.9, 137.8, 131.7, 125.0, 124.3, 124.0, 119.1, 72.1, 55.7, 37.0,

31.4, 29.4, 23.9, 22.2, 13.9. MALDI-TOF MS: m/e 1470.3 (M^+). Elemental Analysis: Calcd for $C_{75}H_{93}I_3S_2$: C, 61.22; H, 6.37; S, 6.54. Found: C, 61.18; H, 6.32; S, 6.55.

General Procedure for the Sonogashira Coupling Reaction:

A flask with a side arm was charged with acetylene derivatives (1.3 eq. per reaction site), aromatic iodide compound (1 eq.), $Pd_2(dba)_3$ (0.02 eq.), triphenylphosphine (0.1 eq.), copper(I) iodide (0.02 eq.) and dry triethylamine (30 mL per mmol of iodide compound). The flask was then evacuated and back-filled with nitrogen three times, capped and stirred at 40 °C. After completion of reaction, the reaction mixture was diluted with dichloromethane. After extraction, the combined organic layers were washed with water and brine, followed by removal of solvents. The residue was purified by flash chromatography to afford the desired product.

Compound 4: The general procedure for the Sonogashira coupling was followed and purified by flash chromatography (hexanes/ CH_2Cl_2 , 5:1) to afford **4** (87%). 1H NMR ($CDCl_3$, 400 MHz, ppm): δ 8.36-8.39 (d, $J = 8.4$ Hz, 3H, Ar-H), 7.67-7.68 (d, s, $J = 8.4$ Hz, 6H, Ar-H), 7.38-7.39 (d, $J = 3.6$ Hz, 3H, Th-H), 7.31-7.34 (m, 9H, Th-H), 7.16-7.18 (d, d, $J = 3.6$ Hz, $J = 3.6$ Hz, 6H, Th-H), 7.01-7.04 (dd, $J = 4.8$ Hz, $J = 3.6$ Hz, 3H, Th-H), 2.95-2.97 (m, 6H, CH_2), 2.10-2.15 (m, 6H, CH_2), 0.84-0.91

(m, 36H, CH₂), 0.56-0.58 (m, 30H, CH₂, CH₃). ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 154.4, 147.1, 145.5, 140.3, 137.9, 133.6, 132.5, 131.9, 131.8, 131.7, 127.9, 127.2, 124.9, 124.6, 124.5, 124.1, 123.0, 122.5, 121.2, 119.2, 87.9, 87.5, 86.7, 85.8, 55.8, 37.0, 31.4, 29.4, 23.9, 22.2, 13.8. MALDI-TOF MS: *m/e* 1728.6 (M⁺). Elemental Analysis: Calcd for C₁₁₁H₁₀₈S₉: C, 77.03; H, 6.29; S, 16.68. Found: C, 76.62; H, 6.20; S, 17.18.

Compound 5: To a solution of diisopropylamine (267.3 mg, 2.7 mmol) in 5 mL of THF at 0 °C was added dropwise *n*-BuLi (1.8 mmol). The solution was cooled to -78 °C. The compound **4** (0.3 mmol, 0.455 g) in THF was then added dropwise using a cannula, and stirred at -78 °C for 1 h. Iodine (914mg, 3.6 mmol) was then added to the mixture as a solid. And the reaction was stirred at -78 °C for 1 h, and then quenched with NH₄Cl solution. The aqueous layer was extracted with EtOAc. The organic layers were washed with brine. The solvents were removed by rotary evaporation, and the residue was purified by column chromatography (hexanes/CH₂Cl₂, 5:1) to provide the desired product as yellow solids (400 mg, 76

%). ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 8.39-8.42 (d, J = 8.4 Hz, 3H, Ar-H), 7.69-7.71 (d, s, J = 8.4 Hz, 6H, Ar-H), 7.41-7.42 (d, J = 3.6 Hz, 3H, Th-H), 7.36-7.38 (d, J = 3.6 Hz, 3H, Th-H), 7.18-7.22 (m, 9H, Th-H), 6.98-6.97 (d, J = 3.6 Hz, 3H, Th-H), 2.95-2.97 (m, 6H, CH_2), 2.10-2.15 (m, 6H, CH_2), 0.84-0.91 (m, 36H, CH_2), 0.56-0.58 (m, 30H, CH_2 , CH_3). ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 154.5, 147.2, 145.6, 140.4, 137.9, 137.3, 137.3, 133.8, 133.7, 132.4, 131.9, 131.8, 128.7, 125.1, 124.2, 124.0, 124.1, 123.1, 121.2, 119.3, 88.3, 87.8, 86.7, 86.3, 75.7, 55.9, 37.1, 31.5, 29.5, 23.9, 22.3, 13.9. MALDI-TOF MS: m/e 2213.6 ($[\text{M}+\text{Ag}]^+$). Elemental Analysis: Calcd for $\text{C}_{111}\text{H}_{105}\text{I}_3\text{S}_9$: C, 63.23; H, 5.02; S, 13.69. Found: C, 62.88; H, 5.14; S, 13.80.

General procedure for the preparation of alkynes from aldehydes:

When a suspension of the bromomethyltriphenylphosphonium bromide (1.5 eq.) in dry THF was cooled to -78°C under an inert gas atmosphere, potassium *tert*-butoxide (1.5 eq.) was added into the mixture as a solid. The yellow mixture was stirred at -78°C for 0.5 h, then at 0°C for 0.5 h. After the mixture was cooled to -78°C again, the solution of aldehydes (1.0 eq.) in dry THF was transferred dropwise to the mixture. The resulting mixture was stirred for 1 h at -78°C . Second portion of potassium *tert*-butoxide (1.5 eq.) was then added in the mixture as a solid at -78°C . The mixture was stirred for 1 h at -78°C , then warm to room temperature. The mixture was quenched by pouring into aqueous saturated ammonium chloride solution. The water layer was extracted with 100 mL EtOAc. The combined organic layers were washed with brine, dried over MgSO_4 . After evaporation of the solvents, the residue was purified by flash chromatography to give the product.

Compound 7: The general procedure for the preparation of alkynes from aldehydes was followed and purified by flash chromatography (hexanes/CH₂Cl₂, 20:1) to make **7** in 72 % yield. ¹H NMR (CDCl₃, 300 MHz, ppm): δ 8.36-8.38 (d, *J* = 8.4 Hz, 3H, Ar-H), 7.66-7.70 (d, s, *J* = 8.4 Hz, 6H, Ar-H), 7.47-7.48 (d, *J* = 3.6 Hz, 2H, Th-H), 7.33-7.35 (d, d, *J* = 4.8 Hz, *J* = 3.6 Hz, 4H, Th-H), 7.14-7.17 (dd, *J* = 4.8 Hz, *J* = 3.6 Hz, 2H, Th-H), 3.44(1H, s, CCH), 2.91-2.97 (m, 6H, CH₂), 2.10-2.15 (m, 6H, CH₂), 0.84-0.91 (m, 36H, CH₂), 0.56-0.58 (m, 30H, CH₂, CH₃). ¹³C NMR (CDCl₃, 75 MHz, ppm): δ 154.5, 154.3, 146.7, 145.3, 145.2, 145.0, 144.8, 140, 139.6, 138.1, 138.0, 137.7, 134.2, 132.5, 131.5, 128.1, 124.9, 124.5, 124.1, 122.9, 122.5, 120.7, 119.2, 81.9, 77.2, 55.7, 37.0, 31.4, 29.4, 23.9, 22.2, 13.8. MALDI-TOF MS: *m/e* 1117.0 (M⁺). Elemental Analysis: Calcd for C₇₇H₉₆S₃: C, 82.74; H, 8.66; S, 8.61. Found: C, 82.74; H, 8.66; S, 8.60.

Compound 8: To a solution of 1.15 g (1.0 mmol) of **6** in 15 mL of methylene chloride was added 15 mL of acetic acid. The mixture was shielded from light and cooled in an ice bath and 492 mg (2.2 mmol) of NIS was added stepwise. After 2 h the mixture was evaporated and the residue was washed with MeOH to give white powders. (1.35 g, 98%). ¹H NMR (CDCl₃, 300 MHz, ppm): δ 9.95 (s, 1H, CHO), 8.41-8.44 (d, *J* = 8.4 Hz, 1H, Ar-H), 8.34-8.38 (d, *J* = 8.4 Hz, 2H, Ar-H), 7.82-7.83 (d, *J* = 3.6 Hz, 1H, Th-H), 7.76-7.79 (d, s, *J* = 8.4 Hz, 2H, Ar-H), 7.60-7.63 (d, *J* = 8.4 Hz, 4H, Ar-H), 7.57-7.58 (d, *J* = 3.6 Hz, 1H, Th-H), 7.30-7.31 (d, *J* = 3.6 Hz, 2H, Th-H), 7.14-7.15 (d, *J* = 3.6 Hz, 2H, Th-H), 2.95-2.97 (m, 6H, CH₂), 2.10-2.15 (m, 6H, CH₂), 0.84-0.91 (m, 36H, CH₂), 0.56-0.58 (m, 30H, CH₂, CH₃). ¹³C NMR (CDCl₃, 75 MHz, ppm): δ 182.7, 154.8, 154.6, 154.4, 150.7, 145.8, 145.6, 145.4, 142.1, 141.7, 139.9, 138.1, 137.9,

137.6, 131.8, 131.1, 125.1, 125.0, 124.7, 124.4, 124.0, 123.9, 119.8, 119.1, 72.1, 55.8, 37.0, 31.4, 29.4, 23.9, 22.2, 13.9. MALDI-TOF MS: m/e 1479.3 ($[M^+ + Ag]^+$). Elemental Analysis: Calcd for $C_{76}H_{94}I_2OS_3$: C, 66.46; H, 6.90; S, 7.00. Found: C, 66.30; H, 6.88; S, 7.22.

Compound 9: The general procedure for the Sonogashira coupling was followed and purified by flash chromatography (hexanes/EtOAc, 6:1) to make **9** in 87 % yield. 1H NMR ($CDCl_3$, 400 MHz, ppm): δ 9.95 (s, 1H, CHO), 8.36-8.42 (m, 9H, Ar-H), 7.81-7.83 (d, $J = 3.6$ Hz, 1H, Th-H), 7.77-7.79 (m, 2H, Ar-H), 7.65-7.75 (m, 16H, Ar-H), 7.57-7.58 (d, $J = 3.6$ Hz, 1H, Th-H), 7.47-7.48 (d, d, $J = 3.6$ Hz, 4H, Th-H), 7.40-7.42 (d, d, $J = 3.6$ Hz, $J = 3.6$ Hz, 4H, Th-H), 7.35-7.37 (d, d, $J = 3.6$ Hz, $J = 3.6$ Hz, 4H, Th-H), 7.32-7.34 (dd, $J = 4.8$ Hz, $J = 0.88$ Hz, 4H, Th-H), 7.14-7.16 (dd, $J = 4.8$ Hz, $J = 3.6$ Hz, 4H, Th-H), 2.95-2.97 (m, 18H, CH_2), 2.10-2.15 (m, 18H, CH_2), 0.84-0.91 (m, 108H, CH_2), 0.56-0.58 (m, 90H, CH_2 , CH_3). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm): δ 182.7, 154.8, 154.7, 154.6, 154.5, 154.4, 154.3, 146.9, 146.7, 146.0, 145.8, 145.6, 145.4, 145.3, 145.1, 144.9, 144.8, 142.2, 141.7, 140.5, 140.2, 140.1, 139.7, 139.6, 138.2, 138.1, 138.0, 137.8, 137.7, 137.5, 133.3, 132.6, 132.0, 131.8, 131.2, 128.1, 125.1, 125.0, 124.6, 124.3, 124.2, 123.9, 123.1, 123.0, 122.0, 121.8,

119.3, 119.2, 87.7, 87.6, 55.9, 55.8, 37.1, 31.5, 29.5, 23.9, 22.3, 13.9. MALDI-TOF MS: m/e 3461.4 ($[M^+ + Ag]^+$). Elemental Analysis: Calcd for $C_{230}H_{284}OS_9$: C, 82.38; H, 8.54; S, 8.61. Found: C, 82.20; H, 8.62; S, 8.68.

Compound 10: The general procedure for the preparation of alkynes from aldehydes was followed and purified by flash chromatography (hexanes/ CH_2Cl_2 , 10:1) to give 10 in **71** % yield. 1H NMR ($CDCl_3$, 400 MHz, ppm): δ 8.36-8.39 (d, $J = 8.4$ Hz, 9H, Ar-H), 7.70-7.71 (d, s, $J = 8.4$ Hz, 18H, Ar-H), 7.48-7.49 (d, $J = 3.6$ Hz, 4H, Th-H), 7.41-7.42 (d, $J = 3.6$ Hz, 4H, Th-H), 7.33-7.37 (m, 10H, Th-H), 7.13-7.17 (dd, $J = 4.8$ Hz, $J = 3.6$ Hz, 4H, Th-H), 3.45 (1H, s, CH-C), 2.95-2.97 (m, 18H, CH_2), 2.10-2.15 (m, 18H, CH_2), 0.84-0.91 (m, 108H, CH_2), 0.56-0.58 (m, 90H, CH_2 , CH_3). ^{13}C NMR ($CDCl_3$, 100MHz, ppm): δ 154.6, 154.5, 154.4, 154.3, 146.9, 146.7, 146.6, 145.5, 145.4, 145.3, 145.2, 145.1, 144.9, 140.5, 140.3, 139.7, 139.6, 138.1, 138.0, 137.9, 137.8, 134.3, 133.3, 132.6, 131.9, 131.8, 128.1, 125.1, 125.0, 124.6, 124.2, 124.1, 123.1, 123.0, 122.9, 122.7, 121.9, 121.8, 120.8, 119.3, 119.2, 87.7, 87.6, 81.9, 77.2, 55.9, 55.8, 37.1, 31.5, 29.5, 23.9, 22.3, 13.9. MALDI-TOF MS: m/e 3457.7 ($[M^+ + Ag]^+$). Elemental Analysis: Calcd for $C_{231}H_{284}S_9$: C, 82.84; H, 8.55; S, 8.62.

Found: C, 82.51; H, 8.51; S, 8.98.

G0: The general procedure for the Sonogashira coupling reaction was followed and purified by flash chromatography (hexanes/CH₂Cl₂, 8:1) to give **G0** in 90% yield. ¹H NMR (CDCl₃, 400 MHz, ppm): δ 8.36-8.42 (m, 12H, Ar-H), 7.69-7.74 (m, 24H, Ar-H), 7.47-7.48 (d, d, *J* = 3.6 Hz, *J* = 0.75 Hz, 6H, Th-H), 7.40-7.42 (d, d, *J* = 3.6 Hz, 6H, Th-H), 7.35-7.37 (d, d, *J* = 3.6 Hz, 6H, Th-H), 7.32-7.34 (dd, *J* = 4.8 Hz, *J* = 0.88 Hz, 6H, Th-H), 7.14-7.16 (dd, *J* = 4.8 Hz, *J* = 3.6 Hz, 6H, Th-H), 2.95-2.97 (m, 24H, CH₂), 2.10-2.15 (m, 24H, CH₂), 0.84-0.91 (m, 144H, CH₂), 0.56-0.58 (m, 120H, CH₂, CH₃). ¹³C NMR (CDCl₃, 100 MHz, ppm): δ 154.54, 154.51, 154.38, 154.34, 146.8, 146.7, 145.5, 145.3, 145.2, 145.1, 144.9, 140.4, 140.3, 139.7, 139.6, 138.1, 138.0, 137.9, 137.8, 133.3, 132.5, 131.9, 131.7, 128.1, 125.0, 124.9, 124.6, 124.2, 124.1,

123.1, 123.0, 122.9, 121.8, 121.7, 119.3, 119.2, 87.7, 87.6, 55.9, 55.8, 37.1, 31.4, 29.5, 23.9, 22.2, 13.9. MALDI-TOF MS: m/e 4549.1 ($[M^+ + Ag]^+$). Elemental Analysis: Calcd for $C_{306}H_{378}S_{12}$: C, 82.76; H, 8.58; S, 8.66. Found: C, 82.33; H, 8.58; S, 9.09.

G1: The general procedure for the Sonogashira coupling reaction was followed and purified by flash chromatography (hexanes/ CH_2Cl_2 , 6:1) to give **G1** in 82% yield. 1H NMR ($CDCl_3$, 400MHz, ppm): δ 8.36-8.42 (m, 30H, Ar-H), 7.69-7.74 (m, 60H, Ar-H), 7.47-7.48 (d, $J = 3.6$ Hz, 12H, Th-H), 7.40-7.42 (m, 18H, Th-H), 7.33-7.36 (m, 30H, Th-H), 7.22-7.23 (m, 12H, Th-H), 7.14-7.16 (dd, $J = 4.8$ Hz, $J = 3.6$ Hz, 6H, Th-H), 2.95-2.97 (m, 60H, CH_2), 2.10-2.15 (m, 60H, CH_2), 0.84-0.91 (m, 360H, CH_2), 0.56-0.58 (m, 300 H, CH_2 , CH_3). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm): δ 154.6, 154.4, 154.3, 147.3, 147.2, 146.9, 146.8, 145.6, 145.4, 145.3, 145.1, 144.9, 140.5, 140.3, 139.7, 139.6, 138.1, 138.0, 137.9, 133.8, 133.3, 132.6, 132.5, 132.1, 131.9,

131.8, 128.1, 125.1, 125.0, 124.9, 124.6, 124.4, 124.2, 124.1, 123.1, 123.0, 122.9, 121.9, 121.7, 121.3, 119.3, 119.3, 88.4, 87.7, 87.6, 87.3, 86.8, 55.9, 55.8, 37.1, 31.5, 29.8, 24.0, 22.3, 13.9. MALDI-TOF MS: m/e 11770 (M^+). Elemental Analysis: Calcd for $C_{804}H_{954}S_{36}$: C, 82.03; H, 8.17; S, 9.81. Found: C, 81.63; H, 8.22; S, 10.12.

2. UV-vis and PL Spectra for dendrimer

Figure S1. Comparison of absorption and excitation spectra of **G0** in dilute solutions. The spectra were normalized at the 380 nm-450 nm absorption region.

Figure S2. Comparison of absorption and excitation spectra of **G1** in dilute solutions. The spectra were normalized at the 380 nm - 450 nm absorption region.

Figure S3. Comparison of absorption spectra of dendrons and dendrimers in thin films.

Figure S4. Comparison of emission spectra of dendrons and dendrimers in thin films.

Table S1. Molecular weights and optical properties of all dendrons and dendrimers.

Cmps.	ϵ $M^{-1}m^{-1}$	Abs. λ_{max}/nm Sol.	Abs. λ_{max}/nm Films	Emis. λ_{max}/nm Sol.	Emis. λ_{max}/nm Films	Molecular mass $g \cdot mol^{-1}$	MALDI-TOF $g \cdot mol^{-1}$
1	1.8×10^7	343	343, 362	365, 383	369, 386	1093.7	1093.0
7	1.8×10^7	343,	345,	377, 397	462, 491	1117.8	1117.0
	1.6×10^7	355	358				
	1.1×10^8	343,	345, 357,	444, 475	508	4441.1	4549.1
G0	1.0×10^8	355,	402				($M^+ + Ag$)
	9.6×10^7	398					
4	2.4×10^7	398	404	435, 464	485	1730.6	1728.6
	3.2×10^7	343,	346, 357,	443, 473	455, 483	3349.3	3457.7
10	2.8×10^7	355,	396				($M^+ + Ag$)
	2.4×10^7	395					
	2.3×10^8	343,	346, 358,	468, 498	521	11772.5	11770
G1	2.4×10^8	355,	404				
	2.7×10^8	410					

3. Copies of ^1H and ^{13}C NMR spectra

^1H NMR spectrum for compound 2.

^{13}C NMR spectrum for compound 2.

¹H NMR spectrum for compound 4.

^{13}C NMR spectrum for compound 4.

¹H NMR spectrum for compound 5.

¹H NMR spectrum for compound **7**.

^{13}C NMR spectrum for compound 7.

^1H NMR spectrum for compound 8.

¹³C NMR spectrum for compound **8**.

Current Data Parameters
NAME n27670
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050729
Time 9:51
INSTRUM ARK400
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 64
DS 0
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5959540 sec
RG 180
DW 79.000 usec
DE 97.50 usec
TE 300.0 K
D1 2.00000000 sec
P1 4.00 usec
DE 97.50 usec
SF01 400.1317965 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 16384
SF 400.1300124 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 4.00

1D NMR plot parameters
CX 20.00 cm
F1p 12.000 ppm
F1 4801.56 Hz
F2p -0.500 ppm
F2 -200.07 Hz
PPMCK 0.62500 ppm/cm
HZCM 250.08125 Hz/cm

¹H NMR spectrum for compound 9.

¹³C NMR spectrum for compound **9**.

Current Data Parameters	
NAME	C27507
EXPNO	1
PROCNO	1
F2 - Acquisition Parameters	
Date_	20050802
Time	7.30
INSTRUM	ARR400
PROBHD	5 mm Multinu
PULPROG	zgpg
TD	32768
SOLVENT	CDCl3
NS	3031
DS	0
SWH	29411.766 Hz
FIDRES	0.897576 Hz
AQ	0.5571050 sec
RG	32768
DW	17.000 usec
DE	21.25 usec
TE	300.0 K
D2	0.0000000 sec
OL5	24.00 dB
CHOPRG	mlt216
Pd1	100.00 usec
Q1	2.0000000 sec
P1	4.00 usec
DE	21.25 usec
SP01	100.6240000 MHz
NUC1EUS	13C
Q11	0.0300000 sec
F2 - Processing parameters	
SI	32768
SF	100.627720 MHz
WIM	EM
SSB	0
LB	1.00 Hz
GB	0
PC	0.30
10 NMR plot parameters	
CX	20.00 cm
FP	230.000 ppm
F1	22134.31 Hz
F2P	-5.000 ppm
F2	-503.07 Hz
PPHCH	11.25000 ppm/cm
HZCM	131.89358 Hz/cm

¹³C NMR spectrum for compound **10**.

Current Data Parameters
NAME h27648
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050727
Time 14.05
INSTRUM ARX400
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 64
DS 0
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 128
DW 76.000 usec
DE 97.50 usec
TE 300.0 K
D1 2.00000000 sec
P1 4.00 usec
DE 97.50 usec
SFO1 400.1317965 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 16384
SF 400.1300132 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 4.00

1D NMR plot parameters
CX 20.00 cm
FIP 10.000 ppm
F1 4001.30 Hz
F2 -0.500 ppm
F2 -200.07 Hz
PPMCH 0.52500 ppm/cm
HZCM 210.06825 Hz/cm

¹H NMR spectrum for compound G0.

Current Data Parameters
NAME C27659
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050728
Time 16.31
INSTRUM AFX400
PROBHD 5 mm NULHnu
PULPROG zgpg
TD 32768
SOLVENT CDCl3
NS 1000
DS 2
SWH 25000.000 Hz
FIDRES 0.762939 Hz
AQ 0.6534100 sec
RG 32768
DM 20.000 usec
DE 25.00 usec
TE 300.0 K
D12 0.0002000 sec
DL5 24.00 dB
COPROG waltz16
P31 100.00 usec
D1 1.50000000 sec
P1 4.00 usec
DE 25.00 usec
SFO1 100.623283 MHz
NUCLEUS 13C
D11 0.03000000 sec

F2 - Processing parameters
SI 32768
SF 100.612754 MHz
MD 0
SSB 0
LB 0.50 Hz
GB 0
PC 1.00

10 NMR plot parameters
CX 20.00 cm
F1p 200.000 ppm
F1 20122.56 Hz
F2p -5.000 ppm
F2 -503.06 Hz
PPHCK 10.25000 ppm/cm
HZCN 1031.28101 Hz/cm

^{13}C NMR spectrum for compound G0.

S30

¹³C NMR spectrum for compound **G1**.