Supporting information for:

New Annulations via Platinum-Catalyzed Enyne Cyclization and Cyclopropane Cleavage

Cristina Nevado, Catalina Ferrer, and Antonio M. Echavarren*

*Institute of Chemical Research of Catalonia (ICIQ), 43007 Tarragona and
Departamento de Química Orgánica, Universidad Autónoma de Madrid, 28049 Madrid,
Spain*

aechavarren@iciq.es

General: All reactions were carried out under Ar in dry freshly distilled solvents under anhydrous conditions. THF was dried by using 4Å molecular sieves. Toluene and DMF were distilled from sodium/benzophenone and CaH₂ respectively and were kept with 4Å. CH₂Cl₂, CH₃CN, Toluene and THF for oxidation and benzannulation reactions were used without further purification from commercial sources. Aldehydes were distilled from NaHCO₃ under Ar or reduced pressure depending on the boiling points.

Thin layer chromatography was carried out using TLC-aluminium sheets with 0.2 mm of silica gel (Merk GF₂₃₄). Chromatography purifications were carried out using flash grade silica gel (SDS Chromatogel 60 ACC, 40-60 µm) with distilled solvents. NMR spectra were recorded at 23°C.

General procedure for the preparation of alcohols¹ precursors of compounds 3a-g, 4a-c, 5a-b: A solution containing 1.2 equiv of the corresponding enol ether in dried THF (volume of THF necessary to make the concentration of enol ether 2 M) was cooled at -78°C and 1 equiv of *t*-BuLi was added dropwise. The mixture was warmed up to -5°C and then stirred at that temperature for 5 h. The solution was then cooled at -78°C and 1 equiv of the aldehyde was slowly added. The mixture was warmed up to room temperature overnight and then quenched with 4 mL of a saturated NH₄Cl solution. The mixture was extracted with 3 x 20 mL of Et₂O and the combined organic layers were dried over MgSO₄. Solvent was removed under reduced pressure and the

(1) Baldwin, E. J.; Höfle, A. G.; Lever, O.W. *J. Am. Chem. Soc.* **1974**, 96, 7125-7127.

residue was purified by silica gel chromatography (15:1 hexane:EtOAc containing 5% Et₃N). Reactions were carried out in a 15-30 mmol scale.

Because of its lability, the alcohol precursor of compound **6f** was not purified and was used in the alkylation without further purification after extractive work up and evaporation of the solvent. Alcohols precursors of compounds **6a-d**, **7a**, **8a** have been previously described.²

(2E)-1-(5,6-Dihydro-4H-pyran-2-yl)but-2-en-1-ol: Yield: 59% (3.20 g, 20.1 mmol). Yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 5.76 (dq, *J* = 15.4, 6.5, 0.8 Hz, 1H), 5.59 (ddq, *J* = 15.4, 6.7, 1.5 Hz, 1H), 4.77 (t, *J* = 3.8 Hz, 1H), 4.38 (br m, 1H), 4.15-3.97 (m, 2H), 2.05-2.00 (m, 2H), 1.85-1.77 (m, 2H), 1.72 (d, *J* = 6.5 Hz, 3H) (the OH signal was not observed); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 154.22 (C), 130.49 (CH), 128.43 (CH), 96.88 (CH), 73.26 (CH), 66.45 (CH₂), 22.33 (CH₂), 19.88 (CH₂), 17.76 (CH₃). EI-HMRS Calcd. for C₉H₁₄O₂: 154.0993. Found: 154.0992.

1-(5,6-dihydro-4H-pyran-2-yl)-2-methylpropan-1-ol:³ Yield: 67% (3.70 g, 23.7 mmol). Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 4.68 (dd, *J* = 3.84, 3.84 Hz, 1H), 4.05-3.91 (m, 2H), 3.48 (t, *J* = 7.8 Hz, 1H), 2.04-1.99 (m, 2H), 1.86-1.66 (m, 4H), 0.95 (d, *J* = 6.7 Hz, 3H), 0.84 (d, *J* = 6.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 154.49, 98.32, 79.51, 66.85, 32.42, 23.15, 20.58, 19.89, 19.01.

(4,5-dihydrofuran-2-yl)(*p*-tolyl)methanol: Yield: 50% (4.00 g, 21.0 mmol). Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.32 (d, *J* = 8.1 Hz, 2H), 7.17 (d, *J* = 8.1 Hz, 2H), 5.23 (br s, 1H), 4.80 (ddd, *J* = 2.7, 2.7, 1.1 Hz, 1H), 4.38 (t, *J* = 9.4 Hz, 2H), 2.70-2.58 (m, 2H), 2.35 (s, 3H) (the OH signal was not observed); ¹³C NMR (75 MHz, CDCl₃) δ 159.74, 138.36, 129.70, 127.20, 97.39, 71.26, 70.82, 30.48, 21.82. (one C signal was not observed due to overlapping). EI-HMRS Calcd. for C₁₂H₁₄O₂: 190.0994 Found: 190.0997.

1-(4,5-Dihydrofuran-2-yl)-2,2-dimethylpropan-1-ol: Yield: 42% (2.80 g, 17.3 mmol). Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 4.79 (t, *J* = 2.4 Hz, 1H), 4.38-4.24 (m, 2H), 3.84 (dd, *J* = 6.1, 0.8 Hz, 1H), 2.62 (ddd, *J* = 9.3, 2.4, 0.8 Hz, 2H), 1.97 (m, 1H), 0.93 (s,

(2) Nevado, C.; Echavarren, A.M. *Tetrahedron*, **2004**, *60*, in press.

(3) Ley, S. V.; Lygo, B.K.; Sternfeld, F.; Wonnacott, A. *Tetrahedron* **1986**, *42*, 4333-4342.

9H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 158.85 (C), 97.44 (CH), 76.94 (CH), 70.50 (CH_2), 35.50 (C), 30.37 (CH_2), 26.60 (CH_3). EI-HMRS Calcd. for $\text{C}_9\text{H}_{16}\text{O}_2$: 156.1150. Found: 156.1144.

2-Ethoxy-4-methylpent-1-en-3-ol: Yield: 46% (2.50 g, 17.5 mmol). Yellow oil. ^1H NMR (300 MHz, CDCl_3) δ 4.06 (d, $J = 2.1$ Hz, 1H), 3.98 (d, $J = 2.1$ Hz, 1H), 3.75 (q, $J = 7.0$ Hz, 2H), 3.65 (t, $J = 7.0$ Hz, 1H), 1.87 (octet, $J = 7.0$ Hz, 1H), 1.30 (t, $J = 7.0$ Hz, 3H), 0.96 (d, $J = 7.0$ Hz, 3H), 0.87 (d, $J = 7.0$ Hz, 3H) (the OH signal was not observed); ^{13}C NMR (75 MHz, CDCl_3) δ 166.78, 86.59, 83.27, 67.22, 36.50, 23.53, 22.29, 18.76. Anal. Calcd. For $\text{C}_8\text{H}_{16}\text{O}_2$: C, 66.63; H, 11.18. Found: C, 66.59; H, 11.39.

General procedure for the preparation of substrates 3a-g, 4a-c, 5a-b: A solution containing 1.2 equiv of NaH (60% in mineral oil) in dried DMF (volume of DMF necessary to make the concentration of NaH 1.0 M) was cooled at 0°C . A 1 equiv solution of the corresponding benzylic alcohol in DMF (volume of DMF necessary to make the concentration of the alcohol 1.0 M) was added dropwise. The mixture was warmed up to room temperature and stirred for 20 min and 1 equiv of the alkylating agent was added. The reaction was stirred at room temperature for 4 h and then quenched with a water-ice mixture and diluted with Et_2O . The organic layer was washed several times with water and then dried over MgSO_4 . Solvent was removed under reduced pressure and the residue was purified by silica gel chromatography (100:1 hexane:EtOAc containing 5% Et_3N). All reactions were carried out in a 5-15 mmol scale.

Enyne **3a** has been previously described.²

2-Napht-6-(1-(but-2-ynyloxy)ethyl)-3,4-dihydro-2H-pyran (3b): Yield: 89% (2.90 g, 9.9 mmol). Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 8.00 (d, $J = 1.0$ Hz, 1H), 7.95-7.88 (m, 3H), 7.65 (dd, $J = 8.5, 1.8$ Hz, 1H), 7.57-7.49 (m, 2H), 5.17 (s, 1H), 5.07 (t, $J = 3.8$ Hz, 1H), 4.33 (dq, $J = 12.9, 2.2$ Hz, 1H), 4.27 (dq, $J = 12.9, 2.2$ Hz, 1H), 4.09-3.99 (m, 2H), 2.18-2.12 (m, 2H), 1.94 (t, $J = 2.2$ Hz, 3H), 1.89-1.84 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 153.03 (C), 137.38 (C), 133.91 (C), 133.81 (C), 128.81 (CH), 128.52 (CH), 128.33 (CH), 126.97 (CH), 126.63 (CH), 126.51 (CH), 125.90 (CH), 100.56 (CH), 83.36 (C), 80.92 (CH), 75.83 (C), 67.14 (CH_2), 57.03 (CH_2), 22.99 (CH_2),

20.86 (CH₂), 4.46 (CH₃). EI-HMRS Calcd. for C₂₀H₂₀O₂: 292.1463. Found: 292.1468. Anal. Calcd. For C₂₀H₂₀O₂: C, 82.16; H, 6.89. Found: C, 82.08; H, 6.95.

3,4-Dihydro-6-((pent-2-ynyloxy)(phenyl)methyl)-2H-pyran (3c): Yield: 90% (5.30 g, 20.6 mmol). Colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.31-7.34 (m, 2H), 7.28-7.19 (m, 3H), 4.58-4.83 (m, 2H), 4.12 (dt, *J* = 15.3, 2.2 Hz, 1H), 4.07 (dt, *J* = 15.3, 2.2 Hz, 1H), 3.95-3.85 (m, 2H), 2.15 (qt, *J* = 7.6, 2.2 Hz, 2H), 2.02-1.97 (m, 2H), 1.75-1.68 (m, 2H), 1.06 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 152.22, 139.21, 128.06, 127.58, 127.25, 99.62, 88.42, 79.97, 75.17, 66.37, 56.25, 22.27, 20.11, 13.78, 12.50.

6-((3-Phenylprop-2-ynyloxy)(phenyl)methyl)-3,4-dihydro-2H-pyran (3d): Yield: 57% (1.80 g, 6.0 mmol). White vitreous solid. ¹H NMR (300 MHz, CDCl₃) δ 7.53-7.30 (m, 10H), 5.05 (s, 1H), 5.01 (t, *J* = 4.0 Hz, 1H), 4.52 (d, *J* = 15.8 Hz, 1H), 4.44 (d, *J* = 15.8 Hz, 1H), 4.11-3.99 (m, 2H), 2.17-2.09 (m, 2H), 1.90-1.82 (m, 2H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 152.78 (C), 139.71 (C), 132.43 (CH), 128.98 (CH), 128.88 (CH), 128.81 (CH), 128.37 (CH), 127.97 (CH), 123.47 (C), 100.65 (CH), 87.01 (C), 85.84 (C), 81.04 (CH), 67.10 (CH₂), 57.16 (CH₂), 22.93 (CH₂), 20.80 (CH₂). EI-HMRS Calcd. for C₂₁H₂₀O₂: 304.1463. Found: 304.1461.

5-((*E*)-1-(But-2-ynyloxy)but-2-enyl)-2,3-dihydro-2H-pyran (3e): Yield: 73% (943 mg, 4.6 mmol). Yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 5.8 (dc, *J* = 15.4, 6.5 Hz, 1H), 5.53 (ddq, *J* = 15.4, 7.5, 1.4 Hz, 1H), 4.81 (t, *J* = 3.6 Hz, 1H), 4.20 (d, *J* = 7.5 Hz, 1H), 4.07 (q, *J* = 2.3 Hz, 2H), 4.04-3.94 (m, 2H), 2.06-2.01 (m, 2H), 1.82 (t, *J* = 2.2 Hz, 3H), 1.83-1.76 (m overlapped, 2H), 1.71 (dd, *J* = 6.23, 1.0 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 152.21 (C), 129.73 (CH), 128.37 (CH), 98.97 (CH), 82.07 (C), 79.37 (CH), 75.29 (C), 66.34 (CH₂), 55.38 (CH₂), 22.30 (CH₂), 20.30 (CH₂), 17.79 (CH₃), 3.62 (CH₃). EI-HMRS Calcd. for C₁₃H₁₈O₂: 206.1306. Found: 206.1297.

6-((*E*)-1-(But-2-ynyloxy)-3-phenylallyl)-3,4-dihydro-2H-pyran (3f): Yield: 19% (353 mg, 1.3 mmol). Yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 7.41-7.37 (m, 1H), 7.32-7.19 (m, 3H), 6.64 (d, *J* = 16.2 Hz, 1H), 6.25 (dd, *J* = 16.2, 7.3 Hz, 1H), 4.89 (t, *J* = 5.0 Hz, 1H), 4.45 (d, *J* = 7.3 Hz, 1H), 4.15 (q, *J* = 2.8 Hz, 2H), 4.07-3.96 (m, 2H), 2.08-2.02 (m, 2H), 1.84 (t, *J* = 2.8 Hz, 3H), 1.81-1.77 (m, 2H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 152.50 (C), 137.33 (C), 133.42 (CH), 129.10 (CH), 128.35 (CH), 127.43 (CH), 127.29 (CH), 100.28 (CH), 83.10 (C), 80.03 (CH), 75.79 (C), 67.09 (CH₂), 56.35 (CH₂), 22.91

(CH₂), 20.73 (CH₂), 4.36 (CH₃). EI-HMRS Calcd. for C₁₈H₁₉O₂ (M⁺-1): 267.1385. Found: 267.1378.

6-(1-(But-2-ynyloxy)-2-methylpropyl)-3,4-dihydro-2H-pyran (3g): Yield: 70% (561 mg, 2.7 mmol). Pale yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 4.73 (t, *J* = 3.6 Hz, 1H), 4.21-4.13 (m, 2H), 4.05-3.91 (m, 2H), 3.27 (d, *J* = 8.5 Hz, 1H), 2.07-1.75 (m, 8H), 0.98 (d, *J* = 6.7 Hz, 3H), 0.85 (d, *J* = 6.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 151.39, 101.25, 85.84, 82.18, 82.05, 76.42, 66.74, 56.64, 30.81, 23.13, 20.81, 19.91, 19.84. EI-HMRS Calcd. for C₁₈H₁₉O₂ (M⁺-1): 267.1385. Found: 267.1378.

5-((But-2-ynyloxy)(phenyl)methyl)-2,3-dihydrofuran (4a): Yield: 69% (7.60 g, 35.4 mmol). Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.43 (dd, *J* = 7.9, 1.4 Hz, 2H), 7.39-7.29 (m, 3H), 5.14 (br s, 1H), 4.87 (td, *J* = 2.4, 0.8 Hz, 1H), 4.37 (t, *J* = 9.5 Hz, 2H), 4.18 (dq, *J* = 15.3, 2.4 Hz, 1H), 4.06 (dq, *J* = 15.3, 2.4 Hz, 1H), 2.62 (tdd, *J* = 9.5, 2.4, 1.2 Hz, 2H), 1.85 (t, *J* = 2.4 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 156.98, 138.33, 128.23, 128.03, 127.45, 98.39, 82.77, 75.49, 74.76, 70.44, 56.41, 29.69, 3.62. EI-HMRS Calcd. for C₁₅H₁₆O₂: 228.1150. Found: 228.1143. Anal. Calcd. for C₁₅H₁₆O₂: C, 78.92; H, 7.06. Found: C, 78.86; H, 7.07.

2,3-Dihydro-5-((pent-2-ynyloxy)(*p*-tolyl)methyl)furan (4b): Yield: 59% (1.20 g, 4.7 mmol). Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.31 (d, *J* = 8.1 Hz, 2H), 7.16 (d, *J* = 8.1 Hz, 2H), 5.12 (br s, 1H), 4.88 (td, *J* = 2.4, 1.0 Hz, 1H), 4.40-4.33 (m, 2H), 4.18 (dt, *J* = 4.5, 2.2, 1H), 4.06 (dt, *J* = 4.5, 2.2 Hz, 1H), 2.66-2.59 (m, 2H), 2.34 (s, 3H), 2.24 (qt, *J* = 7.7, 2.2 Hz, 2H), 1.14 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 157.90, 138.44, 135.98, 129.63, 128.14, 98.74, 89.29, 75.99, 75.65, 71.13, 56.99, 30.37, 21.82, 14.42, 13.15. EI-HMRS Calcd. for C₁₇H₁₉O₂ (M⁺-1): 255.1385. Found: 255.1373.

5-(1-(But-2-ynyloxy)-2,2-dimethylpropyl)-2,3-dihydrofuran (4c): Yield: 53% (1.60 g, 6.8 mmol). Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 4.89 (t, *J* = 2.4 Hz, 1H), 4.43-4.27 (m, 2H), 4.22 (dq, *J* = 15.4, 2.4 Hz, 1H), 4.00 (dq, *J* = 15.4, 2.4 Hz, 1H), 3.70 (s, 1H), 2.68-2.62 (m, 2H), 1.87 (t, *J* = 2.4 Hz, 3H), 0.98 (s, 9H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 156.54 (C), 99.87 (CH), 82.31 (CH), 77.22 (C), 76.22 (C), 70.52 (CH₂), 57.32 (CH₂), 34.95 (C), 30.24 (CH₂), 27.03 (CH₃), 4.20 (CH₃). FAB-HMRS Calcd. for C₁₃H₁₉O₂ (M⁺+1): 209.1541. Found: 209.1543.

1-(1-But-2-ynyloxy)-2-ethoxyallyl)benzene (5a): Yield: 49% (600 mg, 2.6 mmol). Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.50-7.29 (m, 5H), 4.99 (s, 1H), 4.40 (d, J = 2.0 Hz, 1H), 4.20 (q, J = 2.4 Hz, 2H), 4.16 (d, J = 2.0 Hz, 1H), 3.83-3.74 (m, 2H), 1.90 (t, J = 2.4 Hz, 3H), 1.28 (t, J = 7.3 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 161.42 (C), 140.11 (C), 128.73 (CH), 128.34 (CH), 127.98 (CH), 83.91 (CH_2), 83.16 (C), 81.04 (CH), 75.71 (C), 63.75 (CH_2), 57.17 (CH_2), 14.89 (CH_3), 4.35 (CH_3). CI-HMRS Calcd. for $\text{C}_{15}\text{H}_{19}\text{O}_2$ ($\text{M}^+ + 1$): 231.1385. Found: 231.1384.

3-(But-2-ynyloxy)-2-ethoxy-4-methylpent-1-ene (5b): Yield: 70% (561 mg, 2.7 mmol). Pale yellow oil. ^1H NMR (300 MHz, CDCl_3) δ 4.15 (dq, J = 15.0, 2.4 Hz, 1H), 4.05 (q, J = 2.0 Hz, 2H), 3.94 (dq, J = 15.0, 2.4 Hz, 1H), 3.71 (q, J = 6.9 Hz, 2H), 3.39 (d, J = 7.7 Hz, 1H), 1.90 (m, 1H), 1.82 (t, J = 2.0 Hz, 3H), 1.27 (t, J = 6.9 Hz, 3H), 0.94 (d, J = 6.5 Hz, 3H), 0.83 (d, J = 6.5 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 160.22 (C), 85.87 (CH), 84.92 (CH_2), 78.08 (C), 76.30 (C), 63.39 (CH_2), 57.06 (CH_2), 31.29 (CH), 19.69 (CH_3), 19.41 (CH_3), 15.00 (CH_3), 4.27 (CH_3). Anal. Calcd. for $\text{C}_{12}\text{H}_{20}\text{O}_2$: C, 73.43; H, 10.27. Found: C, 73.63; H, 10.57.

General procedure for the cyclization of substrates 3a-g, 4a-c, 5a-b to give 3-oxa-bicyclo[4.1.0]hept-4-ene derivatives 6a-g, 7a-c, 8a-b: A mixture of enyne (0.5 mmol, 1 equiv) and PtCl_2 ($2.5 \cdot 10^{-3}$ mmol, 0.05 equiv) was dissolved in toluene (2.5 mL). The solution was warmed under conditions stated in Table 1 and then filtered through a short path of Celite. The solvent was removed under reduced pressure and the residue was purified by silica gel chromatography (100:1 hexane:EtOAc containing 5% Et_3N) to yield the corresponding 3-oxa-bicyclo[4.1.0]hept-4-ene.

Compound **6a** has been previously described.²

6b: White solid: mp 103-105 °C. ^1H NMR (300 MHz, CDCl_3) δ 7.96 (br s, 1H), 7.94-7.89 (m, 3H), 7.64 (dd, J = 8.5, 1.6 Hz, 1H), 7.56-7.50 (m, 2H), 6.25 (d, J = 5.6 Hz, 1H), 5.23 (d, J = 5.6 Hz, 1H), 4.81 (s, 1H), 3.32 (ddd, J = 10.9, 4.4, 2.4 Hz, 1H), 2.28 (ddd, J = 12.1, 10.9, 2.4 Hz, 1H), 1.87-1.80 (m, 2H), 1.46-1.30 (m, 1H), 1.28 (s, 3H), 1.20-1.11 (m, 1H), 0.94-0.87 (m, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 140.41, 138.26, 133.80 (2 x C), 128.78, 128.37, 128.32, 127.25, 126.60, 126.49, 125.99, 113.69, 74.25, 69.94, 64.95, 25.78, 22.85, 21.96, 16.11, 13.03. EI-HMRS Calcd. for $\text{C}_{20}\text{H}_{20}\text{O}_2$: 292.1463. Found: 292.1469. Anal. Calcd. for $\text{C}_{20}\text{H}_{20}\text{O}_2$: C, 82.16; H, 6.89. Found: C, 81.94; H, 7.03.

Structure and relative configuration of compound **6b** were determined by X-Ray diffraction.

6c: Colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.46-7.44 (m, 2H), 7.39-7.31 (m, 3H), 6.23 (d, $J = 5.6$ Hz, 1H), 5.19 (d, $J = 5.6$ Hz, 1H), 4.59 (s, 1H), 3.33-3.29 (m, 1H), 2.32-2.26 (m, 1H), 1.82-1.77 (m, 2H), 1.63-1.57 (m, 2H), 1.46-1.33 (m, 2H), 1.26-1.11 (m, 1H), 1.05 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 140.78, 140.23, 128.03, 127.87, 127.49, 110.41, 73.89, 69.72, 63.95, 26.80, 25.67, 22.48, 19.98, 15.75, 11.35. Anal. Calcd. for $\text{C}_{17}\text{H}_{20}\text{O}_2$: C, 79.65; H, 7.86. Found: C, 79.78; H, 7.92.

6d: White vitreous solid. ^1H NMR (300 MHz, CDCl_3) δ 7.57 (dd, $J = 8.5, 2.0$ Hz, 2H), 7.48-7.37 (m, 7H), 7.32-7.27 (m, 1H), 6.26 (d, $J = 5.7$ Hz, 1H), 5.25 (d, $J = 5.7$ Hz, 1H), 4.84 (s, 1H), 3.23-3.17 (m, 1H), 2.33 (ddd, $J = 12.5, 10.9, 2.0$ Hz, 1H), 1.99 (m, 2H), 1.88 (ddt, $J = 14.1, 12.1, 6.9$ Hz, 1H), 0.86 (m, 1H), 0.48-0.32 (m, 1H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 140.97 (C), 140.77 (CH), 140.02 (C), 131.93 (CH), 129.44 (CH), 129.25 (CH), 129.17 (CH), 128.71 (CH), 127.55 (CH), 114.12 (CH), 75.23 (CH), 71.64 (C), 65.45 (CH_2), 31.34 (C), 29.41 (CH), 22.64 (CH_2), 19.31 (CH_2). EI-HMRS Calcd. for $\text{C}_{21}\text{H}_{20}\text{O}_2$: 304.1463. Found: 304.1466. Anal. Calcd. for $\text{C}_{21}\text{H}_{20}\text{O}_2$: C, 82.86; H, 6.62. Found: C, 82.56; H, 6.66.

6e: Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 6.03 (d, $J = 5.8$ Hz, 1H), 5.95 (dq, $J = 15.4, 6.5, 0.8$ Hz, 1H), 5.64 (ddq, $J = 15.4, 7.7, 1.6$ Hz, 1H), 5.06 (d, $J = 5.8$ Hz, 1H), 4.11 (d, $J = 7.7$ Hz, 1H), 3.73-3.67 (m, 1H), 3.36 (ddd, $J = 11.1, 11.1, 3.2$ Hz, 1H), 2.05-1.81 (m, 2H), 1.83 (dd, $J = 6.5, 1.6$ Hz, 3H), 1.60-1.56 (m, 2H), 1.26 (dd, $J = 7.7, 1.2$ Hz, 1H), 1.20 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 138.88, 130.07, 128.09, 112.42, 72.67, 67.04, 64.64, 24.09, 22.53, 20.18, 17.95, 15.66, 12.40. EI-HMRS Calcd. for $\text{C}_{13}\text{H}_{18}\text{O}_2$: 206.1306. Found: 206.1304. Anal. Calcd. for $\text{C}_{13}\text{H}_{18}\text{O}_2$: C, 75.69; H, 8.80. Found: C, 75.74; H, 8.90.

6f: Yellow oil. ^1H NMR (300 MHz, CDCl_3) δ 7.33-7.14 (m, 5H), 6.53 (d, $J = 16.0$ Hz, 1H), 6.14 (dd, $J = 16.0, 9.7$ Hz, 1H), 5.90 (d, $J = 5.7$ Hz, 1H), 5.07 (d, $J = 5.7$ Hz, 1H), 4.22 (d, $J = 9.7$ Hz, 1H), 3.90 (m, 1H), 3.35 (m, 1H), 2.18 (d, $J = 9.7$ Hz, 1H), 2.00-1.95 (m, 1H), 1.87-1.73 (m, 3H), 1.30 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 137.85, 136.79, 132.21, 128.40, 126.75, 125.88, 123.57, 111.02, 70.75, 67.62, 67.09, 38.73, 26.83, 25.32, 24.82, 12.40. EI-HMRS Calcd. for $\text{C}_{18}\text{H}_{20}\text{O}_2$: 268.1463 Found: 268.1467.

6g: Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 6.00 (d, J = 5.6 Hz, 1H), 4.94 (d, J = 5.6 Hz, 1H), 3.74 (ddd, J = 10.9, 4.0, 2.4 Hz, 1H), 3.49 (d, J = 1.6 Hz, 1H), 3.34 (ddd, J = 12.1, 10.9, 3.2 Hz, 1H), 2.27 (septet d, J = 7.3, 1.6 Hz, 1H), 1.87 (m, 3H), 1.46 (m, 1H), 1.19 (m, 1H), 1.13 (s, 3H), 1.08 (d, J = 7.3 Hz, 3H), 0.96 (d, J = 7.3 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 140.39 (CH), 113.00 (CH), 76.25 (CH), 66.29 (C), 65.03 (CH_2), 28.84 (CH), 26.21 (CH), 23.20 (CH_2), 21.44 (CH_3), 16.33 (CH_2), 16.28 (CH_3), 13.08 (CH_3). EI-HMRS Calcd. for $\text{C}_{12}\text{H}_{20}\text{O}_2$: 208.1463. Found: 208.1459. Anal. Calcd. for $\text{C}_{12}\text{H}_{20}\text{O}_2$: C, 74.96; H, 9.68. Found: C, 74.94; H, 8.70.

7a: White solid. ^1H NMR (300 MHz, CDCl_3) δ 7.45-7.32 (m, 5H), 6.20 (d, J = 5.9 Hz, 1H), 5.07 (dd, J = 5.9, 0.6 Hz, 1H), 4.91 (s, 1H), 3.78-3.71 (m, 1H), 3.59-3.50 (m, 1H), 2.12-2.01 (m, 1H), 1.93-1.82 (m, 2H), 1.18 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 140.85 (CH), 139.78, 128.08 (CH), 127.92 (CH), 127.29 (CH), 109.82 (CH), 78.72, 75.45 (CH_2), 71.28 (CH), 34.37 (CH), 27.48, 26.30 (CH_2), 11.22 (CH_3). EI-HMRS Calcd for $\text{C}_{15}\text{H}_{16}\text{O}_2$: 228.1150. Found: 228.1143. Anal. Calcd for $\text{C}_{15}\text{H}_{16}\text{O}_2$: C, 78.92; H, 7.06. Found: C, 79.05; H, 7.68.

7b: Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.35 (d, J = 8.1 Hz, 2H), 7.22 (d, J = 8.1 Hz, 2H), 6.30 (d, J = 6.1 Hz, 1H), 5.11 (d, J = 6.1 Hz, 1H), 4.91 (s, 1H), 3.85-3.78 (m, 1H), 3.64-3.56 (m, 1H), 2.40 (s, 3H), 2.16-2.06 (m, 1H), 1.98-1.87 (m, 2H), 1.54 (q, J = 7.3 Hz, 2H), 1.09 (t, J = 7.3 Hz, 3H). ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 142.65 (CH), 138.24 (C), 137.57 (C), 129.46 (CH), 127.92 (CH), 108.15 (CH), 79.90 (C), 76.56 (CH_2), 72.35 (CH), 35.61 (CH or CH_3), 33.86 (C), 27.19 (CH_2), 21.86 (CH or CH_3), 19.14 (CH_2), 12.03 (CH or CH_3). EI-HMRS Calcd. for $\text{C}_{17}\text{H}_{20}\text{O}_2$: 256.1463. Found: 256.1462.

7c: Colorless oil; ^1H NMR (300 MHz, CDCl_3) δ 6.11 (d, J = 5.9 Hz, 1H), 4.95 (d, J = 5.9 Hz, 1H), 4.30-4.21 (m, 1H), 3.88 (ddd, J = 13.7, 4.9, 4.9 Hz, 1H), 3.67 (s, 1H), 2.37-2.25 (m, 1H), 2.10-2.00 (m, 1H), 1.71 (d, J = 6.5 Hz, 1H), 1.11 (s, 3H), 1.09 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 141.53 (CH), 110.24 (CH), 77.25 (C), 76.39 (CH), 73.90 (CH_2), 36.23 (C), 35.50 (CH), 27.48 (CH_3), 26.83 (CH_2), 25.12 (C), 12.04 (CH_3). EI-HMRS Calcd. for $\text{C}_{13}\text{H}_{19}\text{O}_2$ ($\text{M}^+ - 1$): 207.1385. Found: 207.1378.

8a: Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.47-7.31 (m, 5H), 6.20 (d, J = 5.7 Hz, 1H), 5.22 (d, J = 5.7 Hz, 1H), 4.69 (br s, 1H), 3.31 (dq, J = 9.3, 7.3 Hz, 1H), 2.91 (dq, J

= 9.3, 7.3 Hz, 1H), 1.38 (dd, J = 6.1, 5.6 Hz, 1H), 1.30 (s, 3H), 0.99 (dd, J = 5.3, 1.6 Hz, 1H), 0.94 (t, J = 7.3 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 141.61, 140.61, 128.98, 128.75, 128.44, 112.88, 74.25, 65.05, 24.92, 17.97, 16.05 (two C signals were not observed due to overlapping). EI-HMRS Calcd. for $\text{C}_{15}\text{H}_{18}\text{O}_2$: 230.1307. Found: 230.1308. Anal. Calcd. for $\text{C}_{15}\text{H}_{18}\text{O}_2$: C, 78.23; H, 7.88. Found: C, 78.32; H, 7.83.

8b: Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 6.07 (d, J = 5.7 Hz, 1H), 5.07 (d, J = 5.7 Hz, 1H), 3.70 (m, 2H), 3.67 (dq, J = 8.9, 7.1 Hz, 1H), 2.24 (septet d, J = 6.9, 2.2 Hz, 1H), 1.27 (s, 3H), 1.18 (t, J = 7.1 Hz, 3H), 1.11 (d, J = 5.3 Hz, 1H), 1.10 (d, J = 6.9 Hz, 3H), 0.98 (d, J = 6.9 Hz, 3H), 0.93 (dd, J = 5.3, 1.4 Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 141.45 (CH), 112.56 (CH), 74.38 (CH), 71.42 (C), 64.31 (CH_2), 29.51 (CH), 26.47 (CH_2), 21.00 (CH_3), 19.19 (C), 17.90 (CH_3), 16.40 (CH_3), 16.12 (CH_3). Anal. Calcd. for $\text{C}_{12}\text{H}_{20}\text{O}_2$: C, 73.43; H, 10.27. Found: C, 73.45; H, 10.05.

Experimental procedure for oxidation of 6a to 12: A mixture of **6a** (50 mg, 0.2 mmol) and MCPBA (46 mg, 0.2 mmol) was cooled at 0°C and dissolved in MeOH (1.5 mL). The mixture was stirred for 7 h and then quenched with a saturated solution of NH_4Cl (pH = 8) and diluted with Et_2O . The aqueous layer was extracted several times with Et_2O and then dried over MgSO_4 . Solvent was removed under reduced pressure and the residue was purified by silica gel chromatography (100:1 hexane:EtOAc). **12**: Yield: 80% (40 mg, 0.14 mmol). Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.58-7.54 (m, 2H), 7.24-7.15 (m, 3H), 4.95 (s, 1H), 4.14 (d, J = 7.3 Hz, 1H), 3.54 (d, J = 7.3 Hz, 1H), 3.27 (ddd, J = 10.6, 3.3, 3.3 Hz, 1H), 3.16 (s, 3H), 2.35 (ddd, J = 12.2, 10.6, 2.6 Hz, 1H), 1.39 (s, 3H), 1.35-1.10 (m, 4H), 0.65-0.57 (m, 1H) (the OH signal was not observed); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 140.56 (C), 128.81 (CH), 128.56 (CH), 127.78 (CH), 105.97 (CH), 78.48 (CH), 73.95 (CH), 64.97 (CH_2), 63.48 (C), 57.36 (CH_3), 27.73 (CH_2), 26.63 (C), 20.17 (CH), 14.89 (CH_3), 8.54 (CH). EI-HRMS Calcd. for $\text{C}_{17}\text{H}_{22}\text{O}_4$: 290.1518. Found: 290.1521. Anal. Calcd. for $\text{C}_{17}\text{H}_{22}\text{O}_4$: C, 70.32; H, 7.64. Found: C, 69.94; H, 7.70.

The relative configuration of compound **13** was determined by a NOESY experiment in C_6D_6 .

Experimental procedure for oxidation of 6a to 13: A mixture of **6a** (100 mg, 0.4 mmol) and $\text{Pb}(\text{OAc})_4$ (85% in acetic acid) (238 mg, 0.54 mmol) was dissolved in acetic

acid (1.5 ml) and stirred 12 h at room temperature. The reaction was quenched with 5% aqueous NaHCO₃ and diluted with Et₂O. The aqueous layer was extracted several times with Et₂O (10 x 3 mL) and the organic residue was washed with brine and then dried over MgSO₄. Solvent was removed under reduced pressure and the residue was purified by silica gel chromatography (100:1 hexane:EtOAc). **13**: Yield: 28% (35 mg, 0.10 mmol). White vitreous solid. ¹H NMR (300 MHz, CDCl₃) δ 7.41-7.31 (m, 5H), 6.44 (s, 1H), 6.07 (d, *J* = 2.4 Hz, 1H), 3.87 (t, *J* = 5.3 Hz, 2H), 2.42-2.32 (m, 1H), 2.25-2.10 (m, 1H), 2.09 (s, 3H), 2.08 (s, 3H), 1.88-1.79 (m, 3H), 1.47 (dd, *J* = 11.3, 6.0 Hz, 1H), 0.79 (dd, *J* = 8.1, 6.0 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 171.26 (C), 170.56 (C), 139.91 (C), 137.77 (C), 128.42 (CH), 128.31 (C), 127.89 (C), 106.56 (C), 95.15 (CH), 74.43 (CH), 66.99 (CH₂), 62.05 (C), 31.07 (CH), 23.09 (CH₂), 22.50 (CH₂), 21.89 (2 x CH₃), 16.37 (CH₂). EI-HRMS Calcd. for C₂₀H₂₂O₆: 358.1416. Found: 358.1424.

The structure and relative configuration of compound **12** were determined by HMQC, HMBC, COSY, and NOESY experiments.

Experimental procedure for the oxidation of compounds 6a-d, 7a-b:

Conditions a: A mixture of tricycle (0.1 mmol, 1.0 equiv) and DDQ (0.2 mmol, 2.0 equiv) was dissolved in 1.5 mL CH₂Cl₂ and 0.15 mL H₂O and heated under refluxing conditions for 1 or 2 h. The reaction was then quenched with 5% aqueous NaHCO₃ and diluted with CH₂Cl₂. The aqueous layer was extracted several times with CH₂Cl₂. The organic phases are dried over Na₂SO₄ and solvent was removed under reduced pressure. The residue was purified by silica gel chromatography (gradient 75:1-50:1 hexane:EtOAc with 1% Et₃N).

Conditions b: A mixture of tricycle (0.1 mmol, 1.0 equiv) and CAN (0.1 mmol, 1.0 equiv) was dissolved in 1.5 mL CH₃CN and 0.15 mL H₂O and warmed at 70 °C 1 h. The reaction was then quenched with a saturated solution of NH₄Cl (pH = 8) and diluted with Et₂O. The aqueous layer was extracted several times with Et₂O and then dried over MgSO₄. Solvent was removed under reduced pressure and the residue was purified by silica gel chromatography (gradient 75:1-50:1 hexane:EtOAc with 1% Et₃N).

14a: Pale yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 7.41-7.28 (m, 5H), 5.75 (d, *J* = 3.2 Hz, 1H), 5.75-5.73 (m, 1H), 4.87 (s, 1H), 4.03-3.94 (m, 2H), 1.94-1.90 (m, 1H), 1.65-1.53 (m, 3H), 1.39 (d, *J* = 1.2 Hz, 3H), 1.37-1.33 (m, 1H); ¹³C NMR (75 MHz, CDCl₃,

DEPT) δ 140.56 (C), 138.83 (C), 128.34 (CH), 127.87 (CH), 126.53 (CH), 123.57 (CH), 106.25 (C), 99.48 (CH), 85.50 (CH), 65.65 (CH₂), 41.97 (CH), 25.80 (CH₂), 24.74 (CH₂), 19.35 (CH₃). EI-HRMS Calcd. for C₁₆H₁₈O₃: 259.1334. Found: 259.1346. Anal. Calcd. for C₁₆H₁₈O₃: C, 74.29; H, 7.02. Found: C, 73.89; H, 7.18.

The structure and relative configuration of compound **14a** were determined by HMQC, HMBC, COSY, and NOESY experiments.

14b: White vitreous solid. ¹H NMR (300 MHz, CDCl₃) δ 7.85-7.76 (m, 4H), 7.50-7.44 (m, 3H), 5.78 (br s, 2H), 4.99 (s, 1H), 4.06-3.90 (m, 2H), 1.88-1.77 (m, 1H), 1.62-1.46 (m, 4H), 1.28 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 140.73, 136.35, 133.67, 133.46, 128.60, 128.26, 127.93, 126.66, 126.37, 125.43, 124.78, 123.60, 106.33, 99.60, 85.73, 65.70, 42.05, 25.78, 24.69, 19.34. EI-HRMS Calcd. for C₂₀H₂₀O₃: 308.1412. Found: 308.1420. Anal. Calcd. for C₂₀H₂₀O₃: C, 77.90; H, 6.54. Found: C, 78.05; H, 6.74.

14c: Colorless oil. ¹H NMR (300MHz, CDCl₃) δ 7.42-7.27 (m, 5H), 5.80 (d, *J* = 3.6 Hz, 1H), 5.72-5.70 (m, 1H), 4.89 (s, 1H), 4.07-3.95 (m, 2H), 1.93-1.84 (m, 2H), 1.65-1.48 (m, 5H), 0.72 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 145.71 (C), 138.88 (C), 127.82 (CH), 127.81 (CH), 126.61 (CH), 121.54 (CH), 106.22 (C), 99.59 (CH), 85.28 (CH), 65.64 (CH₂), 40.61 (CH), 25.95 (CH₂), 25.48 (CH₂), 24.79 (CH₂), 11.52 (CH₃). Anal. Calcd. for C₁₇H₂₀O₃: C, 74.97; H, 7.40. Found: C, 74.86; H, 7.40.

14d: Colorless oil. ¹H NMR (300MHz, CDCl₃) δ 7.47-7.42 (m, 2H), 7.36-7.21 (m, 6H), 6.79-6.76 (m, 2H), 6.13 (dd, *J* = 3.2, 0.8 Hz, 1H), 5.97 (d, *J* = 3.2 Hz, 1H), 4.99 (s, 1H), 4.10-3.98 (m, 2H), 2.14-2.09 (m, 1H), 1.65-1.48 (m, 4H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 144.26 (C), 138.63 (C), 138.49 (C), 129.04 (CH), 128.59 (2 x CH), 128.09 (CH), 126.67 (2 x CH), 125.05 (CH), 106.36 (C), 99.53 (CH), 85.56 (CH), 65.81 (CH₂), 39.66 (CH), 26.24 (CH₂), 24.79 (CH₂). EI-HRMS Calcd. for C₂₁H₂₀O₃: 320.1412. Found: 320.1410. Anal. Calcd. for C₂₁H₂₀O₃: C, 78.63; H, 6.29. Found: C, 78.23; H, 6.43.

15a: Pale yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 7.44-7.33 (m, 5H), 5.91 (dc, *J* = 3.6, 1.6 Hz, 1H), 5.64 (d, *J* = 3.6 Hz, 1H), 5.03 (s, 1H), 4.33 (td, *J* = 8.5, 2.0 Hz, 1H), 4.00-3.92 (m, 1H), 2.14-1.98 (m, 2H), 1.84-1.77 (m, 1H), 1.63 (dd, *J* = 1.6, 0.8 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 138.72, 138.23, 128.83, 128.31, 126.35, 125.67, 111.94,

97.22, 79.40, 70.66, 46.07, 29.01, 20.62. FAB-HRMS Calcd. for $C_{15}H_{17}O_3$ (M+1): 245.1178. Found: 245.1177.

15b: Colorless oil. 1H NMR (300 MHz, $CDCl_3$) δ 7.25 (d, $J = 7.9$ Hz, 2H), 7.12 (d, $J = 7.9$ Hz, 2H), 5.86 (dq, $J = 3.5, 1.6$ Hz, 1H), 5.61 (d, $J = 3.5$ Hz, 1H), 4.96 (s, 1H), 4.28 (td, $J = 8.5, 2.0$ Hz, 1H), 3.94-3.86 (m, 1H), 2.34 (s, 3H), 2.08-1.77 (m, 5H), 0.93 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 144.23, 137.97, 135.04, 129.46, 126.47, 127.87, 112.02, 97.27, 79.45, 70.67, 44.56, 29.18, 27.38, 21.79, 12.16. FAB-HRMS Calcd. for $C_{17}H_{21}O_3$ (M+1): 273.1491. Found: 273.1497.

16a: Colorless oil. 1H NMR (300 MHz, $CDCl_3$) δ 7.50-7.46 (m, 2H), 7.38-7.31 (m, 3H), 5.73 (d, $J = 3.2$ Hz, 1H), 5.56 (dq, $J = 6.1, 2.0$ Hz, 1H), 4.95 (s, 1H), 3.40 (ddt, $J = 12.1, 4.9, 1.6$ Hz, 1H), 2.91 (ddd, $J = 12.1, 12.1, 2.8$ Hz, 1H), 2.83-2.76 (m, 1H), 2.12-2.01 (m, 3H), 1.81-1.65 (m, 1H), 1.58-1.48 (m, 2H), 1.06 (t, $J = 7.3$ Hz, 3H) (the OH signals were not observed); ^{13}C NMR (75 MHz, $CDCl_3$) δ 143.31, 137.02, 130.37, 129.02, 128.49, 118.98, 105.30, 97.91, 64.54, 47.86, 26.36, 24.86, 22.45, 12.15 (one C signal was not observed due to overlapping).

The relative configuration of compound **16a** was determined by differential-NOE experiments.

16b: Colorless oil. 1H NMR (300 MHz, $CDCl_3$) δ 7.53-7.50 (m, 2H), 7.41-7.27 (m, 8H), 5.93 (dd, $J = 3.4, 2.7$ Hz, 1H), 5.86 (dd, $J = 3.4, 0.6$ Hz, 1H), 5.10 (s, 1H), 3.43 (ddt, $J = 12.5, 5.1, 1.7$ Hz, 1H), 3.31 (ddd, $J = 13.4, 3.5, 2.6$ Hz, 1H), 2.93 (td, $J = 12.5, 3.1$ Hz, 1H), 2.07-1.99 (m, 1H), 1.81-1.71 (m, 1H), 1.50-1.31 (m, 2H) (the OH signals were not observed); ^{13}C NMR (75 MHz, $CDCl_3$, DEPT) δ 141.26 (C), 137.32 (C), 136.10 (C), 129.70 (CH), 128.47 (CH), 128.34 (CH), 127.87 (CH), 127.81 (CH), 126.89 (CH), 122.73 (CH), 104.92 (C), 96.83 (CH), 77.58 (CH), 64.06 (CH_2), 46.45 (CH), 25.78 (CH_2), 22.61 (CH_2). EI-MS (m/z): 348.1.

The structure and relative configuration of **16b** were determined by HMQC, HMBC, COSY, and NOESY experiments.

Experimental procedure for the acid-catalyzed reaction of 6a-d and 7a-b:

Conditions a: To a solution of tricycle (0.1 mmol, 1 equiv) in 1.5 mL of THF was added HCl 3M (large excess: 10-15 equiv). The mixture was heated at 100°C for the stated time (see Table 2) and then quenched with 5% aqueous NaHCO₃ and diluted with Et₂O. The aqueous layer was extracted several times with Et₂O. The organic phases were dried over MgSO₄ and solvent was removed under reduced pressure. The residue was purified by silica gel chromatography (100:1 hexane:EtOAc).

Conditions b: A mixture of tricycle (0.1 mmol, 1 equiv) and *p*-TsOH·H₂O (1.0 equiv) was dissolved in 1.5 mL toluene and warmed at 70°C (reaction times are indicated in Table 2). The reaction was then quenched with 5% aqueous NaHCO₃ and diluted with Et₂O. The aqueous layer was extracted several times with Et₂O. The organic phases were dried over MgSO₄ and solvent was removed under reduced pressure. The residue was purified by silica gel chromatography (100:1 hexane:EtOAc).

5-Methyl-8-phenyl-3,4-dihydro-2H-chromene (19a): Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.55 (dd, *J* = 8.4, 1.3 Hz, 2H), 7.43 (t, *J* = 8.4 Hz, 2H), 7.35 (tt, *J* = 8.4, 1.3 Hz, 1H), 7.10 (d, *J* = 7.6 Hz, 1H), 6.85 (d, *J* = 7.6 Hz, 1H), 4.19 (dd, *J* = 5.2, 5.1 Hz, 2H), 2.76 (t, *J* = 6.7 Hz, 2H), 2.30 (s, 3H), 2.13-2.00 (m, 2H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 152.00 (C), 138.90 (C), 137.07 (C), 129.59 (CH), 127.99 (CH), 127.90 (CH), 126.60 (CH), 121.49 (CH), 121.38 (C), 65.91 (CH₂), 23.05 (CH₂), 22.33 (CH₂), 19.18 (CH₃) (one C signal was not observed due to overlapping). EI-HRMS Calcd. for C₁₆H₁₆O: 224.1201. Found: 224.1193.

The structure of **19a** was determined by HMQC, HMBC, COSY, and NOESY experiments.

5-Methyl-8-(1-naphthyl)-3,4-dihydro-2H-chromene (19b): White vitreous solid. ¹H NMR (300 MHz, CDCl₃) δ 7.86 (s, 1H), 7.84-7.76 (m, 3H), 7.61 (dd, *J* = 8.5, 1.7 Hz, 1H), 7.47-7.36 (m, 2H), 7.10 (d, *J* = 7.6 Hz, 1H), 6.78 (d, *J* = 7.6 Hz, 1H), 4.09 (dd, *J* = 5.2, 5.1 Hz, 2H), 2.67 (t, *J* = 6.6 Hz, 2H), 2.21 (s, 3H), 2.16-2.00 (m, 2H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 152.21 (C), 137.25 (C), 136.61 (C), 133.50 (C), 132.35 (C), 128.34 (CH), 128.24 (CH), 128.09 (CH), 127.96 (CH), 127.87 (C), 127.57 (CH), 127.03 (CH), 125.78 (CH), 125.58 (CH), 121.62 (CH), 121.46 (C), 65.98 (CH₂), 23.05 (CH₂), 22.36 (CH₂), 19.21 (CH₃). EI-HRMS Calcd. for C₂₀H₁₈O: 274.1358. Found: 274.1364.

5-Ethyl-8-phenyl-3,4-dihydro-2H-chromene (19c): Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.57 (dd, $J = 8.5, 1.6$ Hz, 2H), 7.43 (t, $J = 8.5$ Hz, 2H), 7.37-7.29 (m, 1H), 7.17 (d, $J = 8.1$ Hz, 1H), 6.89 (d, $J = 8.1$ Hz, 1H), 4.20 (dd, $J = 4.9, 4.9$ Hz, 2H), 2.84 (t, $J = 6.9$ Hz, 2H), 2.67 (q, $J = 7.7$ Hz, 2H), 2.14-2.06 (m, 2H), 1.31 (t, $J = 7.7$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 152.60, 143.40, 139.64, 131.55, 130.29, 128.93, 128.53, 127.25, 121.31, 120.20, 66.57, 26.02, 23.06, 23.01, 14.78.

5,8-Diphenyl-3,4-dihydro-2H-chromene (19d): Pale yellow oil. This compound was obtained along with 7,8-diphenyl-3,4-dihydro-2H-chromene as a 15:1 mixture. ^1H NMR (300 MHz, CDCl_3) δ 7.63 (dd, $J = 8.4, 1.7$ Hz, 2H, major), 7.49-7.45 (m, 4H, major), 7.46-7.36 (m, 4H major and 4H minor), 7.26 (d, $J = 7.6$ Hz, 1H, major), 7.24-7.15 (m, 5H, minor), 7.10-7.08 (m, 2H, minor), 6.99 (d, $J = 7.9$ Hz, 1H, minor), 6.94 (d, $J = 7.6$ Hz, 1H, major), 4.26 (dd, $J = 5.1, 5.1$ Hz, 2H, major), 4.21 (dd, $J = 5.1, 5.1$ Hz, 2H, minor), 2.96 (t, $J = 6.5$ Hz, 2H, minor), 2.76 (t, $J = 6.5$ Hz, 2H, major), 2.09-2.05 (m, 2H, minor), 2.00-1.96 (m, 2H, major); ^{13}C NMR (75 MHz, CDCl_3) δ 152.31, 151.87, 142.38, 141.63, 141.18, 140.66, 138.74, 136.99, 131.37, 129.88, 129.66, 129.18, 128.92, 128.49, 128.34, 128.25, 127.98, 127.53, 127.41, 127.34, 127.02, 126.84, 126.30, 126.07, 121.84, 121.49, 120.43 (2 x C), 66.77, 66.31, 25.34, 24.73, 22.30, 22.21. EI-HRMS Calcd. for $\text{C}_{21}\text{H}_{18}\text{O}$: 286.1358. Found: 258.1361.

The structures of **19d** and 7,8-diphenyl-3,4-dihydro-2H-chromene were confirmed by COSY, HMBC, and HMQC experiments.

(E)-5-Methyl-8-styryl-3,4-dihydro-2H-chromene (19e): Colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.17 (d, $J = 7.7$ Hz, 1H), 6.71 (d, $J = 7.7$ Hz, 1H), 6.67 (d, $J = 15.8$ Hz, 1H), 6.18 (dq, $J = 15.8, 6.5$ Hz, 1H), 4.22 (dd, $J = 5.3, 5.3$ Hz, 2H), 2.68 (t, $J = 6.5$ Hz, 2H), 2.22 (s, 3H), 2.11-2.03 (m, 2H), 1.92 (dd, $J = 6.5, 1.6$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3 , DEPT) δ 152.36 (C), 136.70 (C), 126.41 (CH), 125.99 (CH), 124.90 (C), 124.22 (CH), 122.00 (CH), 121.50 (C), 66.53 (CH_2), 23.41 (CH_2), 22.96 (CH_2), 19.66 (CH_3), 19.51 (CH_3).

4-Methyl-7-phenyl-2,3-dihydrobenzofuran (20a): Yellow vitreous solid. ^1H NMR (300 MHz, CDCl_3) δ 7.71 (dd, $J = 8.5, 1.4$ Hz, 2H), 7.45 (t, $J = 8.5$ Hz, 2H), 7.36-7.32 (m, 1H), 7.25 (d, $J = 7.9$ Hz, 1H), 6.80 (d, $J = 7.9$ Hz, 1H), 4.68 (dd, $J = 8.7, 8.7$ Hz, 2H), 3.22 (t, $J = 8.7$ Hz, 2H), 2.33 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 138.16,

134.55, 128.95, 128.85, 128.42, 127.40, 127.22, 127.17, 122.55, 121.61, 71.60, 29.49, 19.53. EI-HRMS Calcd. for C₁₅H₁₄O: 210.1045. Found: 210.1040.

The structure of compound **20a** was confirmed by a NOESY experiment.

4-Ethyl-7-(4-methylphenyl)-2,3-dihydrobenzofuran (20b): Colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.57 (d, *J* = 8.3 Hz, 2H), 7.24-7.21 (m, 3H), 6.78 (d, *J* = 7.9 Hz, 1H), 4.62 (dd, *J* = 8.7, 8.7 Hz, 2H), 3.20 (t, *J* = 8.7 Hz, 2H), 2.62 (q, *J* = 7.7 Hz, 2H), 2.38 (s, 3H), 1.26 (t, *J* = 7.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃, DEPT) δ 157.51 (C), 140.36 (C), 137.07 (C), 135.26 (C), 129.68 (CH), 128.76 (CH), 128.48 (CH), 126.39 (C), 121.76 (C), 120.75 (CH), 71.59 (CH₂), 29.19 (CH₂), 26.99 (CH₂), 21.81 (CH₃), 14.81 (CH₃). EI-HRMS Calcd. for C₁₇H₁₈O: 238.1358. Found: 238.1355.

X-ray structure of 6b

Figure 1. Chem3D drawing of **6b** based on the X-ray crystallographic data.

Table 1. Summary of crystallographic data and structure refinement for 6b

Empirical formula	$C_{20}H_{20}O_2$	
Formula weight	292.36	
Temperature	100(2) K	
Wavelength	1.54178 Å	
Crystal system	Monoclinic	
Space group	P2(1)	
Unit cell dimensions	$a = 7.37920(10)$ Å	$\alpha = 90^\circ$.
	$b = 6.09700(10)$ Å	$\beta = 94.1530(10)^\circ$.
	$c = 16.8579(2)$ Å	$\gamma = 90^\circ$.
Volume	$756.462(18)$ Å ³	
Z	2	
Density (calculated)	1.284 Mg/m ³	
Absorption coefficient	0.640 mm ⁻¹	
F(000)	312	
Crystal size	0.19 x 0.09 x 0.03 mm ³	
Theta range for data collection	2.63 to 70.54°.	
Index ranges	$-7 \leq h \leq 9$, $-6 \leq k \leq 7$, $-19 \leq l \leq 19$	
Reflections collected	4895	

Independent reflections	2330 [R(int) = 0.0174]
Completeness to theta = 70.54°	94.5 %
Absorption correction	YES, SADABS v. 2.03
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	2330 / 1 / 280
Goodness-of-fit on F ²	1.028
Final R indices [I>2sigma(I)]	R1 = 0.0313, wR2 = 0.0825
R indices (all data)	R1 = 0.0326, wR2 = 0.0839
Absolute structure parameter	0.0(2)
Extinction coefficient	0.0004(7)
Largest diff. peak and hole	0.156 and -0.155 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 6b. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
C(1)	1436(2)	7293(3)	-226(1)	22(1)
C(2)	1221(2)	8379(3)	-938(1)	23(1)
C(3)	1993(2)	10462(3)	-1034(1)	24(1)
C(4)	2960(2)	11457(3)	-411(1)	22(1)
C(5)	3211(2)	10386(3)	338(1)	19(1)
C(6)	2447(2)	8277(3)	431(1)	18(1)
C(7)	2736(2)	7185(3)	1177(1)	19(1)
C(8)	3735(2)	8133(3)	1799(1)	20(1)
C(9)	4491(2)	10255(3)	1705(1)	21(1)
C(10)	4234(2)	11350(3)	994(1)	21(1)
C(11)	3957(2)	7002(3)	2597(1)	22(1)
C(12)	6247(2)	6631(4)	3660(1)	31(1)
C(13)	5101(2)	6855(4)	4223(1)	31(1)
C(14)	3254(2)	7770(3)	4062(1)	25(1)
C(15)	1880(3)	7254(4)	4655(1)	31(1)
C(16)	2699(2)	7960(3)	3184(1)	21(1)
C(17)	3163(2)	10026(3)	3642(1)	25(1)
C(18)	1694(3)	11711(3)	3752(1)	30(1)
C(19)	-188(3)	11088(4)	3369(1)	33(1)
C(20)	-38(2)	9561(3)	2669(1)	28(1)
O(1)	5845(2)	7176(3)	2873(1)	29(1)
O(2)	872(1)	7577(2)	2936(1)	24(1)

Table 3. Bond lengths [Å] and angles [°] for 6b.

C(1)-C(2)	1.370(2)
C(1)-C(6)	1.421(2)
C(1)-H(1)	0.99(2)
C(2)-C(3)	1.406(3)
C(2)-H(2)	1.00(2)
C(3)-C(4)	1.369(2)
C(3)-H(3)	1.016(19)
C(4)-C(5)	1.422(2)
C(4)-H(4)	0.94(2)
C(5)-C(6)	1.417(2)
C(5)-C(10)	1.420(2)
C(6)-C(7)	1.425(2)
C(7)-C(8)	1.366(2)
C(7)-H(7)	0.99(2)
C(8)-C(9)	1.423(3)
C(8)-C(11)	1.509(2)
C(9)-C(10)	1.372(2)
C(9)-H(9)	0.990(18)
C(10)-H(10)	0.97(2)
C(11)-O(1)	1.4404(19)
C(11)-C(16)	1.523(2)
C(11)-H(11)	1.01(2)
C(12)-C(13)	1.323(3)
C(12)-O(1)	1.379(2)
C(12)-H(12)	0.96(2)
C(13)-C(14)	1.479(3)
C(13)-H(13)	0.98(2)
C(14)-C(15)	1.508(2)
C(14)-C(16)	1.511(2)
C(14)-C(17)	1.547(3)
C(15)-H(15A)	0.98(4)
C(15)-H(15B)	1.01(2)
C(15)-H(15C)	1.00(3)
C(16)-O(2)	1.4024(19)
C(16)-C(17)	1.503(2)
C(17)-C(18)	1.514(3)

C(17)-H(17)	0.99(2)
C(18)-C(19)	1.536(3)
C(18)-H(18A)	0.99(2)
C(18)-H(18B)	0.96(3)
C(19)-C(20)	1.514(3)
C(19)-H(19A)	0.97(2)
C(19)-H(19B)	1.01(3)
C(20)-O(2)	1.439(2)
C(20)-H(20A)	0.97(2)
C(20)-H(20B)	1.008(19)

C(2)-C(1)-C(6)	119.91(17)
C(2)-C(1)-H(1)	122.7(10)
C(6)-C(1)-H(1)	117.4(10)
C(1)-C(2)-C(3)	121.07(16)
C(1)-C(2)-H(2)	119.1(13)
C(3)-C(2)-H(2)	119.8(13)
C(4)-C(3)-C(2)	120.30(16)
C(4)-C(3)-H(3)	120.0(11)
C(2)-C(3)-H(3)	119.7(11)
C(3)-C(4)-C(5)	120.37(17)
C(3)-C(4)-H(4)	119.2(11)
C(5)-C(4)-H(4)	120.4(11)
C(6)-C(5)-C(10)	118.91(15)
C(6)-C(5)-C(4)	119.12(15)
C(10)-C(5)-C(4)	121.96(16)
C(5)-C(6)-C(1)	119.22(15)
C(5)-C(6)-C(7)	119.03(14)
C(1)-C(6)-C(7)	121.75(16)
C(8)-C(7)-C(6)	121.22(16)
C(8)-C(7)-H(7)	120.4(11)
C(6)-C(7)-H(7)	118.3(11)
C(7)-C(8)-C(9)	119.51(15)
C(7)-C(8)-C(11)	120.59(16)
C(9)-C(8)-C(11)	119.82(15)
C(10)-C(9)-C(8)	120.67(15)
C(10)-C(9)-H(9)	119.7(11)
C(8)-C(9)-H(9)	119.6(11)

C(9)-C(10)-C(5)	120.67(16)
C(9)-C(10)-H(10)	120.1(10)
C(5)-C(10)-H(10)	119.2(10)
O(1)-C(11)-C(8)	107.16(13)
O(1)-C(11)-C(16)	112.46(13)
C(8)-C(11)-C(16)	111.83(14)
O(1)-C(11)-H(11)	108.8(10)
C(8)-C(11)-H(11)	110.5(11)
C(16)-C(11)-H(11)	106.1(11)
C(13)-C(12)-O(1)	124.09(16)
C(13)-C(12)-H(12)	123.2(11)
O(1)-C(12)-H(12)	112.6(11)
C(12)-C(13)-C(14)	122.42(16)
C(12)-C(13)-H(13)	119.1(12)
C(14)-C(13)-H(13)	118.3(12)
C(13)-C(14)-C(15)	117.05(15)
C(13)-C(14)-C(16)	112.90(15)
C(15)-C(14)-C(16)	121.19(14)
C(13)-C(14)-C(17)	115.36(15)
C(15)-C(14)-C(17)	118.70(16)
C(16)-C(14)-C(17)	58.88(11)
C(14)-C(15)-H(15A)	115.5(17)
C(14)-C(15)-H(15B)	108.5(13)
H(15A)-C(15)-H(15B)	107(2)
C(14)-C(15)-H(15C)	111.5(14)
H(15A)-C(15)-H(15C)	106(3)
H(15B)-C(15)-H(15C)	108(2)
O(2)-C(16)-C(17)	118.09(14)
O(2)-C(16)-C(14)	117.79(13)
C(17)-C(16)-C(14)	61.75(12)
O(2)-C(16)-C(11)	111.12(13)
C(17)-C(16)-C(11)	121.90(14)
C(14)-C(16)-C(11)	118.22(14)
C(16)-C(17)-C(18)	119.55(15)
C(16)-C(17)-C(14)	59.37(11)
C(18)-C(17)-C(14)	123.68(14)
C(16)-C(17)-H(17)	112.8(13)
C(18)-C(17)-H(17)	116.3(13)

C(14)-C(17)-H(17)	112.8(13)
C(17)-C(18)-C(19)	114.44(16)
C(17)-C(18)-H(18A)	110.6(12)
C(19)-C(18)-H(18A)	108.8(11)
C(17)-C(18)-H(18B)	104.2(15)
C(19)-C(18)-H(18B)	109.5(14)
H(18A)-C(18)-H(18B)	109(2)
C(20)-C(19)-C(18)	111.39(15)
C(20)-C(19)-H(19A)	106.2(13)
C(18)-C(19)-H(19A)	113.5(12)
C(20)-C(19)-H(19B)	111.1(13)
C(18)-C(19)-H(19B)	109.2(13)
H(19A)-C(19)-H(19B)	105.3(18)
O(2)-C(20)-C(19)	109.41(15)
O(2)-C(20)-H(20A)	104.0(13)
C(19)-C(20)-H(20A)	110.9(11)
O(2)-C(20)-H(20B)	106.9(11)
C(19)-C(20)-H(20B)	113.5(11)
H(20A)-C(20)-H(20B)	111.6(14)
C(12)-O(1)-C(11)	115.44(13)
C(16)-O(2)-C(20)	111.72(13)

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 6b. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
C(1)	18(1)	22(1)	25(1)	-4(1)	5(1)	1(1)
C(2)	20(1)	29(1)	22(1)	-4(1)	2(1)	4(1)
C(3)	23(1)	28(1)	22(1)	3(1)	4(1)	5(1)
C(4)	20(1)	21(1)	26(1)	4(1)	7(1)	3(1)
C(5)	14(1)	21(1)	23(1)	-1(1)	5(1)	2(1)
C(6)	14(1)	20(1)	21(1)	-1(1)	6(1)	2(1)
C(7)	16(1)	18(1)	23(1)	0(1)	6(1)	0(1)
C(8)	17(1)	23(1)	21(1)	0(1)	7(1)	2(1)
C(9)	18(1)	24(1)	21(1)	-4(1)	4(1)	-2(1)
C(10)	18(1)	19(1)	27(1)	-2(1)	7(1)	0(1)
C(11)	23(1)	23(1)	20(1)	1(1)	2(1)	-1(1)
C(12)	27(1)	36(1)	27(1)	5(1)	-5(1)	2(1)
C(13)	37(1)	33(1)	21(1)	4(1)	-5(1)	-2(1)
C(14)	33(1)	24(1)	18(1)	0(1)	2(1)	-2(1)
C(15)	44(1)	30(1)	20(1)	-1(1)	7(1)	-6(1)
C(16)	24(1)	22(1)	17(1)	2(1)	4(1)	-2(1)
C(17)	29(1)	26(1)	19(1)	0(1)	4(1)	-5(1)
C(18)	42(1)	22(1)	25(1)	-5(1)	9(1)	-3(1)
C(19)	32(1)	35(1)	35(1)	-5(1)	10(1)	5(1)
C(20)	23(1)	33(1)	28(1)	-2(1)	3(1)	5(1)
O(1)	23(1)	40(1)	24(1)	4(1)	0(1)	5(1)
O(2)	22(1)	27(1)	23(1)	0(1)	4(1)	-2(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for 6b.

	x	y	z	U(eq)
H(1)	890(20)	5830(40)	-140(10)	23(5)
H(2)	530(30)	7660(40)	-1395(12)	36(6)
H(3)	1820(20)	11230(30)	-1569(11)	21(5)
H(4)	3490(20)	12830(40)	-490(10)	23(5)
H(7)	2170(30)	5730(40)	1238(11)	24(5)
H(9)	5220(20)	10940(30)	2153(10)	19(4)
H(10)	4770(20)	12800(40)	933(10)	21(4)
H(11)	3630(20)	5400(40)	2542(11)	21(5)
H(12)	7480(30)	6140(40)	3764(11)	27(5)
H(13)	5520(30)	6520(40)	4775(13)	40(6)
H(15A)	1800(40)	5710(60)	4807(17)	70(9)
H(15B)	2210(30)	8110(40)	5158(12)	37(6)
H(15C)	630(30)	7680(50)	4452(14)	53(7)
H(17)	4380(30)	10610(40)	3540(11)	29(5)
H(18A)	1590(20)	12010(40)	4321(12)	27(5)
H(18B)	2120(30)	13010(50)	3502(13)	48(7)
H(19A)	-950(30)	10340(40)	3730(12)	29(5)
H(19B)	-870(30)	12460(50)	3205(12)	40(6)
H(20A)	-1230(30)	9070(40)	2462(11)	24(5)
H(20B)	680(20)	10200(30)	2238(10)	19(4)

Table 6. Torsion angles [°] for 6b.

C(6)-C(1)-C(2)-C(3)	0.2(2)
C(1)-C(2)-C(3)-C(4)	-0.8(2)
C(2)-C(3)-C(4)-C(5)	0.7(2)
C(3)-C(4)-C(5)-C(6)	0.0(2)
C(3)-C(4)-C(5)-C(10)	178.82(14)
C(10)-C(5)-C(6)-C(1)	-179.40(15)
C(4)-C(5)-C(6)-C(1)	-0.5(2)
C(10)-C(5)-C(6)-C(7)	-0.3(2)
C(4)-C(5)-C(6)-C(7)	178.56(15)
C(2)-C(1)-C(6)-C(5)	0.4(2)
C(2)-C(1)-C(6)-C(7)	-178.66(14)
C(5)-C(6)-C(7)-C(8)	-0.1(2)
C(1)-C(6)-C(7)-C(8)	178.94(14)
C(6)-C(7)-C(8)-C(9)	0.4(2)
C(6)-C(7)-C(8)-C(11)	177.20(14)
C(7)-C(8)-C(9)-C(10)	-0.3(2)
C(11)-C(8)-C(9)-C(10)	-177.11(14)
C(8)-C(9)-C(10)-C(5)	-0.1(2)
C(6)-C(5)-C(10)-C(9)	0.4(2)
C(4)-C(5)-C(10)-C(9)	-178.40(15)
C(7)-C(8)-C(11)-O(1)	135.32(16)
C(9)-C(8)-C(11)-O(1)	-47.9(2)
C(7)-C(8)-C(11)-C(16)	-101.01(18)
C(9)-C(8)-C(11)-C(16)	75.73(18)
O(1)-C(12)-C(13)-C(14)	-1.3(3)
C(12)-C(13)-C(14)-C(15)	159.9(2)
C(12)-C(13)-C(14)-C(16)	12.0(3)
C(12)-C(13)-C(14)-C(17)	-53.1(3)
C(13)-C(14)-C(16)-O(2)	144.53(16)
C(15)-C(14)-C(16)-O(2)	-1.9(3)
C(17)-C(14)-C(16)-O(2)	-108.72(17)
C(13)-C(14)-C(16)-C(17)	-106.75(18)
C(15)-C(14)-C(16)-C(17)	106.9(2)
C(13)-C(14)-C(16)-C(11)	6.4(2)
C(15)-C(14)-C(16)-C(11)	-140.03(19)
C(17)-C(14)-C(16)-C(11)	113.10(17)

O(1)-C(11)-C(16)-O(2)	-174.25(15)
C(8)-C(11)-C(16)-O(2)	65.12(19)
O(1)-C(11)-C(16)-C(17)	39.1(2)
C(8)-C(11)-C(16)-C(17)	-81.50(18)
O(1)-C(11)-C(16)-C(14)	-33.5(2)
C(8)-C(11)-C(16)-C(14)	-154.11(15)
O(2)-C(16)-C(17)-C(18)	-5.6(2)
C(14)-C(16)-C(17)-C(18)	-113.90(17)
C(11)-C(16)-C(17)-C(18)	138.77(16)
O(2)-C(16)-C(17)-C(14)	108.25(16)
C(11)-C(16)-C(17)-C(14)	-107.33(17)
C(13)-C(14)-C(17)-C(16)	102.52(17)
C(15)-C(14)-C(17)-C(16)	-111.05(17)
C(13)-C(14)-C(17)-C(18)	-150.37(17)
C(15)-C(14)-C(17)-C(18)	-3.9(2)
C(16)-C(14)-C(17)-C(18)	107.11(18)
C(16)-C(17)-C(18)-C(19)	0.2(2)
C(14)-C(17)-C(18)-C(19)	-70.8(2)
C(17)-C(18)-C(19)-C(20)	-26.7(2)
C(18)-C(19)-C(20)-O(2)	60.6(2)
C(13)-C(12)-O(1)-C(11)	-28.9(3)
C(8)-C(11)-O(1)-C(12)	167.45(16)
C(16)-C(11)-O(1)-C(12)	44.2(2)
C(17)-C(16)-O(2)-C(20)	39.25(19)
C(14)-C(16)-O(2)-C(20)	110.27(17)
C(11)-C(16)-O(2)-C(20)	-108.77(16)
C(19)-C(20)-O(2)-C(16)	-67.51(18)

Symmetry transformations used to generate equivalent atoms:

12.0478
25.1273
26.8563
27.4977
35.5016
36.2267

73.9032
76.3852
77.2498
77.6681
78.0864

110.2411

141.5314

acetone peak: 2.16 ppm

acetone peak: 30.3 ppm