New Bioactive Lupane Triterpene Coumaroyl Esters Isolated from *Buxus cochinchinensis**

Authors

Li Pan¹, Ulyana Muñoz Acuña², Heebyung Chai¹, Hyun-Young Park¹, Tran Ngoc Ninh³, Bui Van Thanh³, Emilio F. Merino⁴, Maria B. Cassera⁴, L. Harinantenaina Rakotondraibe¹, Esperanza J. Carcache de Blanco^{1,2}, Djaja D. Soejarto^{5,6}, A. Douglas Kinghorn¹

Affiliations

The affiliations are listed at the end of the article

Key words

- lupane triterpenes
- coumaroyl esters
- *Buxus cochinchinensis*
- Buxaceae
- cytotoxicity
- antimalarial activity

Abstract

Five new lupane triterpene coumaroyl esters (**1–5**), together with betulin (**6**) and a known *Buxus* alkaloid, *N*-3-benzoyldihydrocyclocymophylline F (**7**), were isolated from a CHCl₃-soluble partition of a methanol extract of *Buxus cochinchinensis* Pierre ex Gagnep. (Buxaceae) collected in Vietnam. Isolation work was monitored using human colon cancer cells (HT-29). The structures of the new compounds (**1–5**) were determined on the basis of spectroscopic data interpretation. In addition to their cytotoxicity against HT-29 cells and nuclear factor-kappa B (p65) inhibitory activity in an enzyme-linked immunosorbent assay, all

isolates as well as two semisynthetic compounds derived from betulin and **5**, respectively, were also evaluated for their *in vitro* antiparasitic activities against the drug-resistant Dd2 strain of *Plasmodium falciparum* and antifungal effects on the growth of the pathogenic yeast *Candida albicans*. The new lupane triterpene coumaroyl esters (**1–5**), along with a betulin derivative and the known *Buxus* alkaloid, were found to show significant *in vitro* antimalarial activities, with IC₅₀ values ranging from 0.26 to 2.07 μM.

Supporting information available online at <http://www.thieme-connect.de/products>

Introduction

Buxus, commonly known as “boxwood”, is the largest genus of the plant family Buxaceae, which comprises over 70 species, and is distributed mainly in southern Europe, southern Asia, Africa, Madagascar, and North and Central America [1]. In addition to being widely used for landscaping purposes, plants of the genus *Buxus* have also been employed in traditional medicine for the treatment of epilepsy, leprosy, malaria, rheumatism, skin infections, toothache, and venereal disease [2]. Phytochemical studies on plants of this genus have resulted in the isolation and structural characterization of over 200 *Buxus* alkaloids, which biosynthetically are considered as degraded triterpenoid alkaloids based on a cycloartenol skeleton. As the major biological active agents from *Buxus* species, these alkaloids have shown a wide variety of activities, including cytotoxicity, antibacterial, antimalarial, and cholinesterase inhibitory properties [3–5].

Buxus cochinchinensis Pierre ex Gagnep., a large shrub, 1–2 m high, with gray-white stems and dark green fruits, is a boxwood species native to southeastern Asia [6]. To the best of our knowledge, no phytochemical investigation has been carried out on this plant to date. As part of our continuing efforts to discover naturally occurring biologically active agents from plants, a chloroform-soluble partition of a methanol extract from a combination of leaves, twigs, and fruits of *B. cochinchinensis* collected in Vietnam was found to show cytotoxicity against the HT-29 human colon cancer line, with an ED₅₀ value of 7.8 μg/mL, and thus was fractionated by bioactivity-guided isolation using this assay. Seven lupane-type triterpenes, including five new lupane triterpene coumaroyl esters (**1–5**), together with two previously known compounds, betulin (**6**) [7] and *N*-benzoyldihydrocyclocymophylline F (**7**) [8,9], were isolated and identified in the present study. The structures of the new compounds were determined as 3-*O*-(*E*)-*p*-coumaroyl-23-hydroxybetulin (**1**), 3-*O*-(*E*)-*p*-coumaroyl-23-hydroxy-3-*epi*-betulin (**2**), 3-*O*-(*Z*)-*p*-coumaroyl-23-hydroxy-3-*epi*-betulin (**3**), 23-*O*-(*E*)-*p*-coumaroyl-23-hy-

received February 24, 2015
revised April 17, 2015
accepted April 20, 2015

Bibliography

DOI <http://dx.doi.org/10.1055/s-0035-1546118>
Published online July 1, 2015
Planta Med 2015; 81: 1133–1140 © Georg Thieme Verlag KG Stuttgart · New York · ISSN 0032-0943

Correspondence

Prof. Dr. A. Douglas Kinghorn
College of Pharmacy
The Ohio State University
Division of Medicinal Chemistry
and Pharmacognosy
500 W. 12th Avenue
Columbus, OH 43210
USA
Phone: +1 614 247 8642
Fax: +1 614 247 8081
kinghorn.4@osu.edu

* Dedicated to Professor Dr. Adolf Nahrstedt on the occasion of his 75th birthday.

Fig. 1 Chemical structures of new compounds 1–9.

droxybetulin (4), and 23-*O*-(*Z*)-*p*-coumaroyl-23-hydroxybetulin (5) on the basis of spectroscopic data interpretation.

As mentioned above, plants of the genus *Buxus* have been reported with antimalarial- and antifungal-related properties. Thus, besides the human colon cancer cell line HT-29 and the NF- κ B inhibitory bioassays, all the isolates obtained in the current study, together with two semisynthetic compounds, 3,28-*O*-diacetylbetulin [10] and 23-hydroxybetulin [11], which were derived from betulin and 5, respectively, were further screened for their antiplasmodial activities against the drug-resistant Dd2 strain of *Plasmodium falciparum* and for inhibitory effects on the growth of the pathogenic yeast *Candida albicans*.

Results and Discussion

Compounds 1–5 were found to share a common molecular formula of $C_{39}H_{56}O_5$ as determined from the sodium adduct ion peak $[M + Na]^+$ in the HRESIMS of each compound. Based on their spectroscopic data interpretation, the structures of compounds 1–5 could be determined as coumaroyl esters of a lupane-type triterpenoid (Fig. 1), being stereoisomers or positional isomers to each other, as discussed in detail below.

The HRESIMS of 1 afforded a sodiated molecular ion peak at m/z 627.4028, corresponding to an elemental formula of $C_{39}H_{56}O_5Na$ (calcd. for m/z 627.4025). The IR spectrum of compound 1 showed characteristic hydroxy group (3400 cm^{-1}) and conjugated ester carbonyl group (1681 cm^{-1}) absorptions as well as those for an alkene and/or a phenyl ring (1604 and 1514 cm^{-1}) [12, 13]. The UV absorption maxima at 312 and 227 nm suggested the presence of a phenolic acid residue [14]. The ^1H NMR spectrum displayed signals attributable to a *trans*-*p*-coumaroyl

group, with an AA'BB' spin system of a 1,4-disubstituted benzene ring at δ_H 7.52 (2H, d, $J=8.5$ Hz, H-5' and 9') and 6.79 (2H, d, $J=8.5$ Hz, H-6' and 8') as well as a double bond with a *trans* configuration at δ_H 6.32 (1H, d, $J=16.0$ Hz, H-2') and 7.49 (1H, d, $J=16.0$ Hz, H-3'). Also observed in the ^1H NMR spectrum were signals of five tertiary methyl groups at δ_H 0.72 (3H, s, H₃-24), 0.85 (3H, s, H₃-25), 0.96 (3H, s, H₃-27), 1.10 (3H, s, H₃-26), and 1.65 (3H, s, H₃-30), while resonances at δ_H 4.84 (1H, dd, $J=11.0$ and 5.0 Hz, H-3), δ_H 2.96 and 3.16 (each 1H, d, $J=10.5$ Hz, H-23a, and H-23b), as well as δ_H 3.09 and 3.53 (each 1H, d, $J=10.1$ Hz, H-28a, and H-28b), were attributed to proton signals attached to an oxygenated methine group and two oxygenated methylene groups, respectively. In addition, the presence of an olefinic methylene was recognized based on the proton resonances at δ_H 4.68 and 4.55 (each 1H, brs, H-29a and H-29b) (Table 1). In the ^{13}C NMR spectrum of 1, besides the nine carbon resonances ascribed to the *p*-coumaroyl moiety at δ_C 166.1 (C, C-1'), 114.9 (CH, C-2'), 144.1 (CH, C-3'), 125.1 (C, C-4'), 130.7 (2 $\times$ CH, C-2' and 9'), 115.7 (2 $\times$ CH, C-6' and 8'), and 159.6 (C, C-7'), the remaining 30 carbon signals were classified from DEPT and HSQC analysis into five methyls, ten alkyl methylenes, five alkyl methines, five alkyl quaternary carbons, three oxygen-bearing carbons (including two primary and one secondary), and two alkene carbons belonging to a disubstituted terminal double bond (Table 2). These NMR data suggested that 1 is an (*E*)-*p*-coumaroyl ester of a pentacyclic lupane-type triterpene, with the skeleton similar to that of known oxygenated betulin analogues [7, 12, 13, 15]. In the HMBC spectrum, besides the correlations with C-4, C-23, and CH₃-24, the low-field oxygenated methine proton at δ_H 4.84 was also found to show a key correlation with C-1', the carbonyl carbon of the coumaroyl moiety. Consequently, it could be deduced that the esterification by a coumaric acid unit occurs at C-

Table 1 ^1H NMR chemical shifts of compounds **1–5**^a.

Position	1	2	3	4	5
1 α	0.91, m	1.21 ^b	1.34 ^b	0.93, m	0.85, m
1 β	1.61 ^b	1.48 ^b	1.44 ^b	1.69 ^b	1.65 ^b
2 α	1.61 ^b	1.92 ^b	1.90 ^b	1.67 ^b	1.63 ^b
2 β	1.68 ^b	1.69 ^b	1.67 ^b	1.67 ^b	1.63 ^b
3	4.84, dd (11.0, 5.0)	4.94, s	4.85, s	3.45, t (8.0)	3.35 ^b
5	1.27 ^b	1.37 ^b	1.29 ^b	1.03 ^b	0.91 ^b
6 α	1.45 ^b	1.46 ^b	1.41 ^b	1.49 ^b	1.39 ^b
6 β	1.31 ^b	1.30 ^b	1.29 ^b	1.43 ^b	1.39 ^b
7a	1.31 ^b	1.39 ^b	1.34 ^b	1.43 ^b	1.31 ^b
7b	1.49 ^b	1.48 ^b	1.44 ^b	1.43 ^b	1.31 ^b
9	1.35 ^b	1.46 ^b	1.38 ^b	1.32 ^b	1.25 ^b
11a	1.38 ^b	1.46 ^b	1.44 ^b	1.42 ^b	1.39 ^b
11b	1.19 ^b	1.22 ^b	1.20 ^b	1.21 ^b	1.20 ^b
12 α	1.04 ^b	1.11 ^b	1.10 ^b	1.05 ^b	1.04 ^b
12 β	1.59 ^b	1.66 ^b	1.65 ^b	1.65 ^b	1.63 ^b
13	1.59 ^b	1.64 ^b	1.63 ^b	1.65 ^b	1.62 ^b
15a	1.69 ^b	1.71 ^b	1.71 ^b	1.70 ^b	1.68 ^b
15b	0.91 ^b	1.08 ^b	1.05 ^b	1.08 ^b	1.04 ^b
16a	1.93 ^b	1.93 ^b	1.94 ^b	1.94 ^b	1.94 ^b
16b	1.10 ^b	1.24 ^b	1.22 ^b	1.22 ^b	1.21 ^b
18	1.50 ^b	1.60 ^b	1.60 ^b	1.57 ^b	1.57 ^b
19	2.39, m	2.40, m	2.39, m	2.38, td (10.0, 5.8)	2.38, td (10.3, 5.7)
21 α	1.28 ^b	1.41 ^b	1.41 ^b	1.41 ^b	1.41 ^b
21 β	1.87 ^b	1.96 ^b	1.94 ^b	1.94 ^b	1.94 ^b
22 α	0.88 ^b	1.06 ^b	1.05 ^b	1.05 ^b	1.05 ^b
22 β	1.87 ^b	1.88 ^b	1.86 ^b	1.86 ^b	1.86 ^b
23a	2.96, d, (10.5)	3.34 ^b	3.35 ^b	3.88, d, (11.6)	3.83, d, (11.4)
23b	3.16, d, (10.5)	3.40, d, (11.2)	3.35 ^b	4.38, d, (11.6)	4.28, d, (11.4)
24	0.72, s	0.95, s	0.92, s	0.81, s	0.76, s
25	0.85, s	0.92, s	0.89, s	0.87, s	0.84, s
26	1.10, s	1.05 ^b	1.02 ^b	1.03, s	1.00, s
27	0.96, s	1.05 ^b	1.03 ^b	0.98, s	0.93, s
28a	3.09, d, (10.1)	3.34 ^b	3.35 ^b	3.34, d, (10.9)	3.34 ^b
28b	3.53, d, (10.1)	3.82, d, (10.3)	3.82, d, (10.3)	3.80, d, (10.9)	3.80, d, (11.2)
29a	4.68, brs	4.70, brs	4.70, brs	4.68, brs	4.67, brs
29b	4.55, brs	4.59, brs	4.59, brs	4.57, brs	4.58, brs
30	1.65, s	1.69, s	1.70, s	1.67, s	1.68, s
Coumaric acid					
1'					
2'	6.32, d, (16.0)	6.33, d, (15.9)	5.88, d, (12.6)	6.27, d, (15.9)	5.84, d, (12.6)
3'	7.49, d, (16.0)	7.66, d, (15.8)	6.89, d, (12.8)	7.63, d, (15.9)	6.89, d, (12.7)
5', 9'	7.52, d, (8.5)	7.44, d, (8.5)	7.64, d, (8.5)	7.41, d, (8.6)	7.61, d, (8.4)
6', 8'	6.79, d, (8.5)	6.84, d, (8.5)	6.79, d, (8.6)	6.83, d, (8.6)	6.80, d, (8.6)

^a Measured at 400 MHz; NMR data of compound **1** were obtained in DMSO- d_6 ; NMR data of compounds **2–5** were obtained in CDCl₃ with TMS as the internal standard; *J* values (Hz) are given in parentheses. Assignments are based on ^1H - ^1H COSY, HSQC, and HMBC spectroscopic data; ^b Multiplicity patterns unclear due to signal overlapping

3. Furthermore, the oxygenated methylene protons at δ_{H} 2.96 and 3.16 were found to show HMBC correlations with C-5 and C-3, respectively, while signals from another set of oxygenated methylene protons at δ_{H} 3.09 and 3.53 displayed long-range correlations with C-16 and C-22, respectively. Therefore, the positions of the free hydroxy groups were concluded as occurring at C-23 and C-28. In addition, HMBC correlations of the terminal methylene protons of H-29 with CH₃-30 and C-19 confirmed the presence of a characteristic isopropene moiety on ring E, which is a typical structural feature of betulin analogues (● Fig. 2). In the NOESY spectrum, a key correlation of H-3 with H-5 indicated the β -orientation of the coumaroyl substituent on C-3. The NOE effects observed from H₃-24 to H₃-25, as well as from H-23 to H-5, suggested the hydroxyethylene group (C-23) on C-4 to be in an α -position. In the same way, the hydroxyethylene group (C-28) on

C-17 was inferred as being β -oriented based on the NOE resonances of H-28a/H-13 and H₃-26. Furthermore, key NOE effects were also observed from H₃-25 to H₃-26, H₃-26 to H-13, H₃-27 to H-9 and H-18 as well as H-19 to H-13, which revealed that the relative configuration of compound **1** is consistent with those of known betulin analogues (● Fig. 2). Accordingly, the structure of **1** was elucidated as 3-*O*-(*E*)-*p*-coumaroyl-23-hydroxybetulin. The HRESIMS of compound **2** showed a sodiated molecular ion peak at *m/z* 627.4048, corresponding to a molecular formula of C₃₉H₅₆O₅Na, the same as that of compound **1**. The NMR spectra of compound **2** were quite comparable with those of compound **1**, with the major differences focused on ring A. In the ^1H NMR spectrum of **1**, H-3 appeared at δ_{H} 4.94 as a broad singlet, with the coupling pattern very different from that of compound **1**, where it was observed as a double doublet with coupling con-

Position	1	2	3	4	5
1	37.6	33.8	33.7	38.6	38.4
2	22.9	23.0	23.0	26.0	26.0
3	73.6	76.1	76.4	72.8	72.7
4	41.3	41.2	41.1	42.5	42.5
5	46.2	46.0	45.8	48.5	48.3
6	17.1	18.2	18.3	18.2	18.2
7	33.2	33.7	33.7	34.0	33.9
8	40.4	40.9	40.9	40.9	40.9
9	49.7	50.3	50.3	50.6	50.5
10	36.2	37.1	37.0	37.1	37.0
11	20.4	20.7	20.7	20.9	20.8
12	24.8	25.1	25.2	25.2	25.2
13	36.7	37.2	37.3	37.3	37.3
14	42.2	42.8	42.8	42.7	42.7
15	26.7	27.0	27.0	27.0	27.0
16	29.0	29.2	29.2	29.2	29.2
17	47.4	47.7	47.8	47.8	47.8
18	48.2	48.8	48.8	48.8	48.8
19	47.3	47.8	47.8	47.9	47.8
20	150.3	150.4	150.5	150.4	150.4
21	29.3	29.7	29.8	29.7	29.8
22	33.8	34.0	34.0	34.0	34.0
23	62.8	68.9	69.0	67.0	67.0
24	13.2	16.8	16.9	11.9	11.9
25	16.2	16.4	16.4	16.7	16.6
26	15.7	16.0	16.0	16.0	16.0
27	14.5	15.0	15.0	14.8	14.7
28	57.9	60.6	60.6	60.6	60.6
29	109.6	109.7	109.7	109.8	109.7
30	18.8	19.1	19.1	19.1	19.1
Coumaric acid					
1'	166.1	167.8	167.0	168.0	167.2
2'	114.9	115.0	116.5	114.7	116.5
3'	144.1	145.7	144.8	145.4	144.3
4'	125.1	126.7	126.8	126.6	127.2
5', 9'	130.7	130.2	132.6	130.1	132.4
6', 8'	115.7	115.9	115.2	116.0	115.1
7'	159.6	158.3	157.6	158.6	157.3

Table 2 ^{13}C NMR chemical shifts of compounds **1–5**^a.

^a Measured at 100 MHz; NMR data of compound **1** were obtained in DMSO- d_6 ; NMR data of compounds **2–5** were obtained in CDCl_3 with TMS as the internal standard. Assignments are based on HSQC and HMBC NMR spectra

Fig. 2 Selected HMBC (→, blue) and NOESY (↔, red) correlations observed for compound **1**. (Color figure available online only.)

stants of 11.0 and 5.0 Hz (Table 1). This implied that rather than an equatorial β -position in compound **1**, the coumaroyl substituent on C-3 in compound **2** adopts an axial α -substitution. Due to

this change, a gauche γ -effect on C-1 of compound **1** was caused by the axially oriented coumaroyl group at C-3 [16], which led to an upfield shift of 3.8 ppm for C-1 in the ^{13}C NMR spectrum when compared with this same signal in compound **1** (Table 2). Furthermore, in the NOESY spectrum, H-3 was observed to show strong correlations with H-24 and H-2 β , as well as relatively weak correlations with H-23 and H-2 α , which was consistent with its presumed equatorial β -configuration. Thus, the structure of compound **2** was elucidated as 3-*O*-(*E*)-*p*-coumaroyl-23-hydroxy-3-*epi*-betulin (**2**), or the C-3 epimer of compound **1**.

The molecular formula of compound **3** was determined to be $\text{C}_{39}\text{H}_{56}\text{O}_5\text{Na}$, the same as that of compounds **1** and **2**, from the sodiated molecular ion peak at m/z 627.4060 in the HRESIMS. A close inspection of the NMR data of compounds **2** and **3** revealed that the major differences were in the coumaroyl residue on C-3. By comparison of the ^1H NMR spectrum of **3** with that of **2**, the olefinic doublets of H-2' and H-3' of the coumaroyl group were upfield shifted from δ_{H} 6.33 and 7.66 to δ_{H} 5.88 and 6.89, respectively, with the coupling constant being ~ 12.7 Hz, smaller than ~ 16.0 Hz, which was observed for that of compound **2** (Table 1). These notable differences suggested that in compound **3**, the

Compound	HT-29 ^c	NF-κB (p65) ^d	<i>P. falciparum</i> ^e
1	> 20	> 20	0.65 ± 0.06
2	13.2	> 20	1.28 ± 0.13
3	3.3	> 20	1.02 ± 0.09
4	> 20	> 20	0.26 ± 0.01
5	18.4	> 20	0.63 ± 0.02
6	> 20	> 20	> 10
7	1.9	> 20	2.07 ± 0.13
8	> 20	> 20	> 10
9	> 20	4.0	1.33 ± 0.37
Paclitaxel ^f	0.001	–	–
Rocaglamide ^g	–	0.08	–
Artemisinin ^h	–	–	0.0007

Table 3 Bioactivity evaluation of compounds isolated from *B. cochinchinensis*^{a,b}.

^a No obvious growth inhibition of *C. albicans* cells were observed for the test compounds at a concentration of 40 mg/mL; ^b Results are expressed as ED₅₀ or IC₅₀ values (μM); ^c Compounds with ED₅₀ values of > 20 μM in HT-29 cells are considered inactive; ^d Compounds with IC₅₀ values of > 20 μM on an enzyme-based ELISA NF-κB (p65) assay are considered inactive; ^e Compounds with IC₅₀ values of > 10 μM in *P. falciparum* assay are considered inactive; ^f Used as a positive control substance for the cytotoxicity assay; ^g Used as a positive control substance for the cytotoxicity assay; ^h Used as a positive control substance for the *P. falciparum* assay

double bond of the *p*-coumaroyl group adopted a *Z* configuration rather than an *E* configuration. Further analysis of the 2D NMR data of **3** revealed the presence of comparable HMBC effects and NOE correlations with that of compound **2**. Accordingly, the structure of compound **3** was determined as 3-*O*-(*Z*)-*p*-coumaroyl-23-hydroxy-3-*epi*-betulin (**3**), a configurational stereoisomer of compound **2**.

Compound **4** gave a molecular formula of C₃₉H₅₆O₅, the same as that of compounds **1–3**, based on the analysis of the HRESIMS. When the ¹H NMR data of compound **4** were compared with those of compound **1**, the oxygenated methine proton signal assigned to H-3 was shifted upfield from δ_H 4.84 ppm to δ_H 3.45 ppm, while the protons of one oxygenated methylene group appeared at a relatively low field region with chemical shifts of δ_H 3.88 (1H, d, *J* = 11.6 Hz) and 4.38 (1H, d, *J* = 11.6 Hz), respectively (• Table 1). In the HMBC spectrum, these two oxygenated methylene protons were observed to have correlations with C-3, C-4, C-24, and C-1', the carbon of the carbonyl group of the coumaroyl moiety. All of this evidence suggested that the esterification with the coumaroyl residue occurred at C-23 in compound **4** instead of on C-3 in compound **1**. In addition, the equatorial β-orientation of the free hydroxy group on C-3 was concluded from the coupling pattern of H-3 (*t*, *J* = 8.0 Hz) and confirmed by the observed NOE correlations of H-3 with H-23 and H-1α as well as H-5. Thus, the structure of compound **4** was elucidated as 23-*O*-(*E*)-*p*-coumaroyl-23-hydroxybetulin (**4**), a positional isomer of compound **1**. Compound **5** showed comparable NMR spectra to those of **4**. The *cis* configuration of the coumaroyl group was recognized based on the notable upfield shifted olefinic protons (H-2' and H-3') at δ_H 5.84 and 6.89 ppm as well as a relatively smaller coupling constant of ~ 12.7 Hz ascribed to these two characteristic doublets when compared with compound **4** (• Table 1). Further HMBC analysis revealed that the (*Z*)-*p*-coumaroyl functionality is also located at C-23, the same as in compound **4**. Moreover, key NOE effects observed for **5** were similar to those observed for **4**. Consequently, compound **5** was designated as 23-*O*-(*Z*)-*p*-coumaroyl-23-hydroxybetulin (**5**), a configurational stereoisomer of compound **4**.

Betulin-derived lupane-type triterpenes are distributed widely throughout the plant kingdom. A variety of biological activities, including cytotoxicity, antiviral, antibacterial, and anti-inflammatory activities, and an *in vitro* antimalarial effect, have been

ascribed to certain betulin derivatives [17]. The natural occurring 3-*epi*-betulin-derived triterpenes are much less common than the normal analogues, and the plant family Celastraceae seems to be prominent in biosynthesizing 3-*epi*-betulin derivatives [18, 19]. When considering the naturally occurring esterification of the hydroxy groups of these betulin-type triterpenes, cinnamic acid-derived substitution is not very common and more often occurs at C-3 and C-28 rather than at other positions on the skeleton. The co-occurrence of *E*- and *Z*-isomers of the coumaroyl moieties in the present study raises the question of if the *E/Z*-isomerization was induced experimentally. The presence of both *E/Z*-coumaroyl isomers in the CHCl₃-soluble extract could be detected by ¹H NMR and HPLC analysis. In addition, for pure compounds **1–5**, when dissolved in DMSO-*d*₆ or CDCl₃ and stored under normal storage conditions at room temperature for over 48 h, no changes were observed between the *E/Z*-coumaroyl isomer profiles. Thus, in the present study, the co-occurrence of the *E/Z*-isomers of the coumaroyl moieties of the triterpenes in *B. cochinchinensis* is more likely derived through a naturally biosynthetic isomerization. However, it should also be noted that according to the literature, certain *Z*-coumaroyl-substituted betulin analogues are not very stable, and can be slowly and partially converted into *E*-configured derivatives when stored with light protection in pyridine-*d*₅ or when dry [20].

All pure compounds obtained in the present investigation were evaluated for their cytotoxic activity against the HT-29 human colon cancer cell line. The known *Buxus* alkaloid *N*-3-benzoyldihydrocyclophyllylline **7** and the new compound 3-*O*-(*Z*)-*p*-coumaroyl-23-hydroxy-3-*epi*-betulin (**3**) were found to be the most cytotoxic agents against HT-29 cells in the current work, with ED₅₀ values of 1.9 μM and 3.3 μM, respectively. In addition, marginal cytotoxic effects were observed for 3-*O*-(*E*)-*p*-coumaroyl-23-hydroxy-3-*epi*-betulin (**2**) and 23-*O*-(*Z*)-*p*-coumaroyl-23-hydroxybetulin (**5**). In this bioassay, triterpenes with a (*Z*)-*p*-coumaroyl group exhibited more potent activity than the (*E*)-*p*-coumaroyl triterpene esters (• Table 3).

An ELISA NF-κB assay was also employed to evaluate the p65 (Rel A) inhibitory activity of all the isolates, of which most were considered inactive with IC₅₀ values of > 20 μM in this assay, except for the semisynthetic compound 23-hydroxybetulin (**9**, IC₅₀ = 4.0 μM). This result implies that the occurrence of free hydroxy groups at C-3, C-23, and C-28 in the molecule might be important

for the observed NF- κ B inhibitory activity ascribed to 23-hydroxybetulin.

Furthermore, the drug-resistant Dd2 strain of *P. falciparum* and a pathogenic strain of *C. albicans* (ATCC® 18804™) were used to evaluate the antiplasmodial and antifungal effects of all the isolates, respectively. No obvious growth inhibition of *C. albicans* cells was observed for the test compounds at a concentration of 40 μ g/mL.

In the *P. falciparum* assay, all the new compounds (1–5), along with the known *Buxus* alkaloid, *N*-benzoyldihydrocyclocimaphylline F (7), were found to show *in vitro* antimalarial activities, with IC₅₀ values ranging from 0.26 to 2.07 μ M (● Table 3). To gain some information on the structure-activity relationships, two semisynthetic compounds, 3,28-*O*-diacetylbetulin and 23-hydroxybetulin, which were derived from betulin and compound 5, respectively, were also subjected to the same bioassay. The test results showed that 3,28-*O*-diacetylbetulin, the diacetate derivative of betulin, was inactive (IC₅₀ > 20 μ M), while 23-hydroxybetulin, the hydrolysis product of compound 5, retained antimalarial potency with an IC₅₀ value of 1.33 $\pm$ 0.37 μ M. Although betulin triterpenes have been reported to possess antimalarial activities, the *in vitro* test results of these analogues against the parasitic strains, to date, are not very promising. According to previous studies, betulinic acid exhibited *in vitro* antiplasmodial activities against *P. falciparum* strains K1, T9–96, and 3D7, with IC₅₀ values ranging from 19.0 μ M to 56.8 μ M, and its methyl derivative showed *in vitro* antimalarial activity against *P. falciparum* strain 3D7, with an IC₅₀ value of 7.0 μ M [21–23]. All of this evidence suggested that the substitution by a hydroxy group or a coumaroyl ester group at C-23 plays an important role for the observed antiplasmodial activities ascribed to these derivatives. Moreover, esterification at C-3 or C-23 by a coumaroyl group seems to slightly increase the resultant activity, while 3-*epi* betulin analogues were found to be less potent than the normal 3 β -hydroxy derivatives in this regard.

Although many *Buxus* plants have documented ethnomedicinal uses for the treatment of malaria [3,4], and some plant extracts in this genus have also been reported to show antimalarial activities in *in vitro* and/or *in vivo* studies [24,25], the chemical types of the active principles that are responsible for the antiplasmodial effects are incompletely understood. As the major constituents of boxwood species, *Buxus* alkaloids have been proposed as the most likely active constituents for the antimalarial activity ascribed to these plants. However, to the best of our knowledge, thus far only the known cycloartane alkaloid *O*-tigloylcyclovibuxine-B, isolated from *Buxus sempervirens*, has been reported to show *in vitro* antiplasmodial activity against *P. falciparum*, with an IC₅₀ value of 0.92 μ M (0.46 μ g/mL) [26]. The present investigation has revealed that besides the well-known *Buxus* alkaloids, lupane triterpenoids based on 23-hydroxybetulin are another class of constituents potentially responsible for the antimalarial properties exhibited by plants of the genus *Buxus*.

Materials and Methods

General experimental procedures

Optical rotations were recorded on a Perkin-Elmer 343 automatic polarimeter (PerkinElmer). UV spectra were measured on a Hitachi U-2910 spectrophotometer (Hitachi). NMR spectroscopic data were obtained on a Bruker Avance DRX-400 spectrometer (Bruker) using standard Bruker pulse sequences (room tempera-

ture), and were processed with Topspin™ 3.1 (Bruker BioSpin). High-resolution electrospray ionization mass spectra (HRESIMS) were performed on a Micromass Q-ToF™ (Micromass) mass spectrometer (calibration with sodium iodide). Analytical TLC was carried out with precoated 250 μ m thickness silica gel UV₂₅₄ aluminum-backed plates (Sorbent Technologies), and preparative TLC was carried out with precoated 500 μ m thickness silica gel UV₂₅₄ glass-backed plates (Sorbent Technologies). Column chromatography was conducted on silica gel (230–400 mesh; Sorbent Technologies). HPLC was carried out on a Waters system comprised of a 600 controller, a 717 Plus autosampler, and a 2487 dual wavelength absorbance detector. Waters Xbridge® (4.6 $\times$ 150 mm), semipreparative (10 $\times$ 150 mm), and preparative (19 $\times$ 150 mm) C₁₈ (5 μ m) columns were used for analytical, semipreparative, and preparative HPLC, respectively.

Plant material

A combination of the leaves, twigs, and fruits of *B. cochinchinensis* Pierre ex Gagnep. (Buxaceae) was collected in Nui Chua National Park, Ninh Thuan Province, Vietnam (11°41.128' N; 109°09.694' E) by D.D.S., T.N.N., and Bui Van Thanh, on 24 July, 2011, who also identified this plant. A voucher specimen (original collection Soejarto et al. 14857) has been deposited in the John G. Searle Herbarium of the Field Museum of Natural History (under accession number FM-2300811), Chicago, Illinois, USA.

Extraction and isolation

A combination of dried and milled leaves, twigs, and fruits of *B. cochinchinensis* (5 kg) was macerated overnight using 95% MeOH at room temperature (3 $\times$ 6 L). This crude extract was concentrated under reduced pressure to yield 750 g of thick dark brown syrup, which was partitioned sequentially with hexane (3 $\times$ 2 L) and chloroform (3 $\times$ 2 L). The chloroform partition was washed with 1% saline solution (3 L) to yield 150 g of a partially detannified chloroform-soluble extract, which was found to be active against the HT-29 cell line (ED₅₀ = 12.0 μ g/mL). Accordingly, part of this fraction (140 g) was subjected to separation over a Si gel column (7 $\times$ 50 cm), with a gradient elution system of CH₂Cl₂-acetone (20:1, 15:1, 10:1, 8:1, 6:1, 4:1, 3:1, 2:1, 1:1, and pure acetone, 4 L per gradient) to yield 17 fractions (F01–F17). A substantial amount of a white precipitate formed from a nontoxic subfraction F03 (15:1, 23.0 g) when it was dissolved in a mixture of CHCl₃-methanol (3:1). Betulin (6) (1.0 g, purity > 95%) was obtained after the recrystallization of this precipitate from the same solvent. Fractions F06 (10:1) and F12 (3:1) were found to be the most active against HT-29 cells (ED₅₀ = 8.0 and 6.3 μ g/mL, respectively) among these subfractions obtained. Fraction F06 (2.5 g) was chromatographed over an open C₁₈ reversed-phase column (2 $\times$ 20 cm) with an MeOH-H₂O gradient solvent system (50:50, 60:40, 70:30, 80:20, and 90:10, 200 mL per gradient) to yield 17 pooled subfractions (F601–F617). Subfractions F610 (80:20, 100 mg) and F611 (80:20, 110 mg) were detected by TLC as terpene-rich subfractions, and were combined and subjected to separation by preparative TLC (20 $\times$ 20 cm, 500 μ m $\times$ 2), developed by CHCl₃-acetone (5:1, 200 mL for each plate) to yield compound 4 (*R*_f = 0.7, 12.0 mg, purity > 98%), and three subfractions designated as F6(10–11)A–D. Subfraction F6(10–11)C (30 mg) was chromatographed on a semipreparative RP-18 column (150 mm $\times$ 10 mm i.d.) by HPLC using a gradient CH₃CN-H₂O solvent system [50:50 to 100% CH₃CN (both solvents with 0.1 formic acid) in 20 min; flow rate 6.0 mL/min] to afford compound 5 (15 mg, *t*_R = 11.9 min, purity > 95%). Subfraction F6

(10–11)D (33 mg) was purified on the same semipreparative RP-18 column by HPLC using a gradient $\text{CH}_3\text{CN-H}_2\text{O}$ solvent system (60:40 to 100% CH_3CN in 17 min; flow rate 5.0 mL/min) to afford compounds **2** (6.0 mg, $t_R = 15.2$ min, purity > 95%) and **3** (7.0 mg; $t_R = 15.8$ min, purity > 95%), respectively. Compound **1** (15 mg, purity > 97%) was precipitated from the parent solution of subfraction F615 (90:10, 35 mg) and recrystallized from a mixture of $\text{MeOH-H}_2\text{O}$.

Fraction F12 (1.2 g) was subjected to passage over a Sephadex LH-20 column (2×35 cm) and eluted with 100% MeOH (600 mL) to furnish five pooled subfractions (F1201–F1205). Subfraction F1204 (200 mg), which gave a positive test for Dragendorff's reagent, was subjected to separation over a Si gel column (1×40 cm) with a gradient elution system of $\text{CHCl}_3\text{-MeOH}$ (20:1, 10:1, 5:1, and 2:1, 200 mL per gradient) to afford the known alkaloid *N*-benzoyldihydrocyclocymophylline F (**7**) (2:1, 5.0 mg).

3-O-(E)-p-coumaroyl-23-hydroxybetulin (1): white powder; m.p. 199–200 °C; $[\alpha]_D^{20} + 32.0$ (c 0.06, MeOH); UV (MeOH) λ_{max} (log ϵ) 203 (4.23), 227 (4.08), 312 (4.34) nm; IR (film) ν_{max} 3400, 2942, 2859, 1681, 1632, 1604, 1514, 1450, 1369, 1168, 1012, 831, 756 cm^{-1} ; ^1H NMR (400 MHz, $\text{DMSO-}d_6$) and ^{13}C NMR (150 MHz, $\text{DMSO-}d_6$) data, see **Tables 1 and 2**; HRESIMS m/z 627.4028 $[\text{M} + \text{Na}]^+$ (calcd. for $\text{C}_{39}\text{H}_{56}\text{O}_5\text{Na}$, 627.4025).

3-O-(E)-p-coumaroyl-23-hydroxy-3-epi-betulin (2): white powder; m.p. 180–182 °C; $[\alpha]_D^{20} + 5.0$ (c 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 205 (4.23), 226 (4.10), 312 (4.34) nm; IR (film) ν_{max} 3350, 2945, 2854, 1681, 1631, 1604, 1514, 1450, 1368, 1167, 1025, 831, 756 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (150 MHz, CDCl_3) data, see **Tables 1 and 2**; HRESIMS m/z 627.4048 $[\text{M} + \text{Na}]^+$ (calcd. for $\text{C}_{39}\text{H}_{56}\text{O}_5\text{Na}$, 627.4025).

3-O-(Z)-p-coumaroyl-23-hydroxy-3-epi-betulin (3): white powder; m.p. 165–167 °C; $[\alpha]_D^{20} - 26.5$ (c 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 204 (4.22), 310 (4.19) nm; IR (film) ν_{max} 3352, 2944, 2859, 1688, 1604, 1514, 1451, 1376, 1167, 1026, 831, 756 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (150 MHz, CDCl_3) data, see **Tables 1 and 2**; HRESIMS m/z 627.4060 $[\text{M} + \text{Na}]^+$ (calcd. for $\text{C}_{39}\text{H}_{56}\text{O}_5\text{Na}$, 627.4025).

23-O-(E)-p-coumaroyl-23-hydroxybetulin (4): white powder; m.p. 165–167 °C; $[\alpha]_D^{20} + 3.0$ (c 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 204 (4.07), 228 (3.95), 313 (4.24) nm; IR (film) ν_{max} 3370, 2940, 2864, 1687, 1632, 1604, 1514, 1450, 1381, 1167, 1022, 831, 755 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (150 MHz, CDCl_3) data, see **Tables 1 and 2**; HRESIMS m/z 627.4023 $[\text{M} + \text{Na}]^+$ (calcd. for $\text{C}_{39}\text{H}_{56}\text{O}_5\text{Na}$, 627.4025).

23-O-(Z)-p-coumaroyl-23-hydroxybetulin (5): white powder; m.p. 162–163 °C; $[\alpha]_D^{20} + 51$ (c 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 204 (4.10), 311 (4.08) nm; IR (film) ν_{max} 3374, 2942, 2870, 1691, 1604, 1514, 1451, 1376, 1166, 1023, 832, 756 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (150 MHz, CDCl_3) data, see **Tables 1 and 2**; HRESIMS m/z 627.4054 $[\text{M} + \text{Na}]^+$ (calcd. for $\text{C}_{39}\text{H}_{56}\text{O}_5\text{Na}$, 627.4025).

Preparation of 3,28-O-diacetylbetulin (8) from betulin (6)

Betulin (**6**) (18 mg) was dissolved in a mixture of pyridine (1.0 mL) and acetic anhydride (1.0 mL). After being stirred overnight at room temperature, the excess pyridine and acetic anhydride were evaporated to yield 19 mg of 3,28-O-diacetylbetulin (**8**) (purity > 98%). This compound was obtained as a white powder [m.p. 222–223 °C (lit. m.p. 223–224 °C)], and its spectroscopic data (NMR and MS) were consistent with published values [10].

Preparation of 23-hydroxybetulin (9) from 23-O-(Z)-p-coumaroyl-23-hydroxybetulin (5)

23-O-(Z)-p-coumaroyl-23-hydroxybetulin (**5**) (2.0 mg) was dissolved in 0.3 mL MeOH , and then 0.3 mL of $\text{LiOH}\cdot\text{H}_2\text{O}$ (5 mg LiOH dissolved in 0.3 mL of H_2O) was added. After being stirred overnight at room temperature, 3 mL of saturated aqueous solution of NaCl were added to this mixture, which was further partitioned with CHCl_3 . The organic phase was evaporated under reduced pressure after washing with water to give 23-hydroxybetulin (**9**) (1.4 mg, purity > 95%). This compound was obtained as a white powder [m.p. 258–260 °C (lit. m.p. 260–262 °C)], and its spectroscopic data (NMR and MS) were consistent with published values [11].

Cytotoxicity assay

The cytotoxic activity of extracts, chromatographic fractions of extracts, and all pure compounds were evaluated against a human colon cancer (HT-29) cell line, according to a previously described protocol [27]. Paclitaxel (Sigma-Aldrich, purity $\geq 97\%$) was used as the positive control substance.

Enzyme-linked immunosorbent nuclear factor-kappa B assay

The NF- κB p65 subunit inhibitory activity of pure compounds was tested in an ELISA NF- κB assay, which was carried out according to a published protocol [28,29]. Rocaglamide (Enzo Life Sciences, purity $\geq 97\%$) was used as the positive control substance.

Antimalarial bioassay

The effects of compounds **1–6** on *P. falciparum* Dd2 parasite growth was measured in a 72-h growth assay, as described previously, with minor modifications [30–32]. Artemisinin (Sigma-Aldrich, purity $\sim 98\%$) was used as the positive control substance.

Antifungal bioassay

A pathogenic strain of *C. albicans* (ATCC® 18804™) was used to evaluate the antifungal effects of all the isolates, according to a published protocol [33]. The antibiotic amphotericin B (Sigma-Aldrich, purity $\sim 80\%$) was used as the positive control substance in this bioassay, which showed 100% growth inhibition of *C. albicans* cells at a concentration of 0.2 mg/mL.

Supporting information

The ^1H , ^{13}C , and 2D NMR spectra of the new compounds **1–5**, as well as the ^1H NMR spectra of two semisynthetic compounds, are available in the Supporting Information.

Acknowledgements

▼ This study was supported by grant P01 CA125066 (awarded to A.D. Kinghorn) from NCI, NIH, Bethesda, MD, USA. We are grateful to Dr. Craig McElroy, College of Pharmacy, The Ohio State University, for facilitating the acquisition of the NMR spectroscopic and mass spectrometric data measurements. The plant material was collected under the terms and conditions of a Memorandum of Agreement between the University of Illinois at Chicago and the Institute of Ecology and Biological Resources (IEBR) of the Vietnam Academy of Science and Technology, Hanoi, Vietnam. Thanks are expressed to the Director of Nui Chua National Park

for permission, and to the Director of IEBR for overseeing the field operation in the collection of the plant.

Conflict of Interest

The authors declare no conflict of interest.

Affiliations

- ¹ Division of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, The Ohio State University, Columbus, OH, USA
- ² Division of Pharmacy Practice and Administration, College of Pharmacy, The Ohio State University, Columbus, OH, USA
- ³ Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology, Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam
- ⁴ Department of Biochemistry and the Virginia Tech Center for Drug Discovery, M/C 0308, Virginia Tech, Blacksburg, VA, USA
- ⁵ Department of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, University of Illinois at Chicago, Chicago, IL, USA
- ⁶ Science and Education, Field Museum of Natural History, Chicago, IL, USA

References

- 1 Batdorf LR. Boxwood: an illustrated encyclopedia. Boyce, VA: American Boxwood Society; 2004: 1–343. Hawaii Book Library website is available at <http://www.hawaiilib.org/article/when0000334850/buxus>. Accessed February 19, 2015
- 2 Grieve MM. A modern herbal. Available at <https://www.botanical.com/botanical/mgmh/b/box-67.html>. Accessed July 23, 2014
- 3 Atta-ur-Rahman, Choudhary MI. Chemistry and biology of steroidal alkaloids. In: Cordell GA, editor. The alkaloids: chemistry and pharmacology. San Diego: Academic Press; 1998: 61–108
- 4 Ata A, Andersh BJ. Buxus steroidal alkaloids: chemistry and biology. Alkaloids Chem Biol 2008; 66: 191–213
- 5 Yan YX, Sun Y, Li ZR, Zhou L, Qiu MH. Chemistry and biological activities of Buxus alkaloids. Curr Bioact Compd 2011; 7: 47–64
- 6 Julius A. Buxus holtzmaniana of Peninsular Malaysia and Thailand is a variety of B. cochinchinensis (Buxaceae). Phytotaxa 2014; 167: 201–204
- 7 Mahato SB, Kundu AP. Review article number 98: ¹³C NMR spectra of pentacyclic triterpenoids – a compilation and some salient features. Phytochemistry 1994; 37: 1517–1575
- 8 Kupchan SM, Kennedy RM, Schleigh WR, Ohta G. Buxus alkaloids. XII. Benzamide alkaloids from Buxus sempervirens. Tetrahedron 1967; 23: 4563–4586
- 9 Sangare M, Khuong-Huu F, Herlem D, Millie A, Septe B, Berenger G, Lukacs G. Revision of the configuration of the C-4 hydroxymethylene group in Buxus alkaloids by ¹³C NMR spectroscopy. Tetrahedron Lett 1975; 22/23: 1791–1794
- 10 Symon AV, Veselova NN, Kaplun AP, Vlasenkova NK, Fedorova GA, Lyutik AI, Gerasimova GK, Shvets VI. Synthesis and antitumor activity of cyclopropane derivatives of betulinic and betulonic acids. Russ J Bioorg Chem 2005; 31: 286–291
- 11 Dracinsky M, Richtr V, Krecek V, Sejbál J, Klinot J, Budesinsky M. Triterpenes. Part CXIII. Preparation and conformational study of 19 β ,28-epoxy-18 α -olean-5-ene derivatives. Collect Czech Chem Commun 2006; 71: 387–410
- 12 Chang CI, Kou YH. Two new lupane-type triterpenes from Diospyros maritima. J Nat Prod 1999; 62: 309–310
- 13 Chumkaew P, Kato S, Chantrapromma K. A new triterpenoid ester from the fruits of Brugiera parviflora. Chem Pharm Bull 2005; 53: 95–96
- 14 Robbins RJ. Phenolic acids in foods: an overview of analytical methodology. J Agric Food Chem 2003; 51: 2866–2887
- 15 Guerrero-Analco JA, Martineau L, Saleem A, Madiraju P, Muhammad A, Durst T, Haddad P, Arnason JT. Bioassay-guided isolation of the antidiabetic principle from Sorbus decora (Rosaceae) used traditionally by the Eeyou Istchee Cree First Nations. J Nat Prod 2010; 73: 1519–1523
- 16 Schneider HJ, Volker H. Carbon-13 nuclear magnetic resonance substituent-induced shieldings and conformational equilibria in cyclohexanes. J Org Chem 1978; 43: 3866–3873
- 17 Alakurtti S, Maekelae T, Koskimies S, Yli-Kauhaluoma J. Pharmacological properties of the ubiquitous natural product betulin. Eur J Pharm Sci 2006; 29: 1–13
- 18 Du YC, Lin AS, Wu CC, Hsieh PW, Chen YH, Chen IH, Chen SL, Yen HF, Lubken T, Chang FR, Wu YC. New cytotoxic lupane triterpenes from Perrottetia arisanensis. Planta Med 2009; 75: 848–855
- 19 Zhou J, Li CJ, Yang JZ, Ma J, Li Y, Bao XQ, Chen XG, Zhang D, Zhang DM. Lupane triterpenoids from the stems of Euonymus carnosus. J Nat Prod 2014; 77: 276–284
- 20 Chien NQ, Hung NV, Santarsiero BD, Mesecar AD, Cuong NM, Soejarto DD, Pezzuto JM, Fong HHS, Tan GT. New 3-O-acyl betulinic acids from Strychnos vanprukii Craib. J Nat Prod 2004; 67: 994–998
- 21 Steele JCP, Warhurst DC, Kirby GC, Simmonds MSJ. In vitro and in vivo evaluation of betulinic acid as an antimalarial. Phytother Res 1999; 13: 115–119
- 22 Ziegler HL, Franzyk H, Sairafianpour M, Tabatabai M, Tehrani MD, Bagherzadeh K, Hagerstrand H, Staerk D, Jaroszewski JW. Erythrocyte membrane modifying agents and the inhibition of Plasmodium falciparum growth: structure-activity relationships for betulinic acid analogues. Bioorg Med Chem 2004; 12: 119–127
- 23 Duker-Eshun G, Jaroszewski JW, Asomaning WA, Oppong-Boachie F, Christensen SB. Antiplasmodial constituents of Cajanus cajan. Phytother Res 2004; 18: 128–130
- 24 Esmaeili S, Naghibi F, Mosaddegh M, Sahranavard S, Ghafari S, Abdullah NR. Screening of antiplasmodial properties among some traditionally used Iranian plants. J Ethnopharmacol 2009; 121: 400–404
- 25 Orhan IE, Erdem SA, Senol FS, Kartal M, Sener B. Exploration of cholinesterase and tyrosinase inhibitory, antiprotazoal and antioxidant effects of Buxus sempervirens L. (boxwood). Ind Crops Prod 2012; 40: 116–121
- 26 Althaus J, Jerz G, Winterhalter P, Kaiser M, Brun R, Schmidt TJ. Antiprotazoal activity of Buxus sempervirens and activity-guided isolation of O-tigloylcyclovirobuxine-B as the main constituent active against Plasmodium falciparum. Molecules 2014; 19: 6184–6201
- 27 Pan L, Kardono LBS, Riswan S, Chai HB, Carcache de Blanco EJ, Pannell CM, Soejarto DD, McCloud TG, Newman DJ, Kinghorn AD. Isolation and characterization of minor analogues of silvestrol and other constituents from a large-scale recollection of Aglaia foveolata. J Nat Prod 2010; 73: 1873–1878
- 28 Renard P, Ernest I, Houbion A, Art M, Le Calvez H, Raes M, Remacle J. Development of a sensitive multi-well colorimetric assay for active NFkB. Nucleic Acids Res 2001; 29: e21
- 29 Deng Y, Balunas MJ, Kim JA, Lantvit DD, Chin YW, Chai HB, Sugiarso S, Kardono LBS, Fong HHS, Pezzuto JM, Swanson SM, Carcache-Blanco EJ, Kinghorn AD. Bioactive 5,6-dihydro-alpha-pyrone derivatives from Hyptis brevipes. J Nat Prod 2009; 72: 1165–1169
- 30 Smilkstein M, Sriwilaijaroen N, Kelly JX, Wilairat P, Riscoe M. Simple and inexpensive fluorescence-based technique for high-throughput antimalarial drug screening. Antimicrob Agents Chemother 2004; 48: 1803–1806
- 31 Bennett TN, Paguio M, Gligorijevic B, Seudieu C, Kosar AD, Davidson E, Roepe PD. Novel, rapid, and inexpensive cell-based quantification of antimalarial drug efficacy. Antimicrob Agents Chemother 2004; 48: 1807–1810
- 32 Dai Y, Harinantenaina L, Bowman JD, Da Fonseca IO, Brodie PJ, Goetz M, Cassera MB, Kingston DG. Isolation of antiplasmodial anthraquinones from Kniphofia ensifolia and synthesis and structure-activity relationships of related compounds. Bioorg Med Chem 2014; 22: 269–276
- 33 National Committee for Clinical Laboratory Standards. Reference method for broth dilution antifungal susceptibility testing of yeasts: approved standard. Wayne, PA: NCCLS Document M27-A 1997; 17: 1–29