Supporting Information

New Entries in Lewis Acid–Lewis Base Bifunctional Asymmetric Catalyst: Catalytic Enantioselective Reissert Reaction of Pyridine Derivatives

Eiko Ichikawa,^{†,‡} Masato Suzuki,[†] Kazuo Yabu,[†] Matthias Albert,[†] Motomu Kanai,^{*,†,‡} and Masakatsu Shibasaki^{*}

General: NMR spectra were recorded on a JEOL JNM-LA500 spectrometer, operating at 500 MHz for ¹H NMR, 125.65 MHz for ¹³C NMR, and 202 MHz for ³¹P NMR. Chemical shifts were reported downfield from TMS (= 0) for ¹H NMR. For ¹³C NMR, chemical shifts were reported in the scale relative to the solvent used as an internal reference. ³¹P NMR was carried out with phosphoric acid (85%) as an external standard. Optical rotations were measured on a JASCO P-1010 polarimeter. Column chromatographies were performed with silica gel Merck 60 (230-400 mesh ASTM). The enantiomeric excess (ee) was determined by HPLC analysis. HPLC analysis was performed on JASCO HPLC systems consisting of the following: pump, 880-PU or PU-980; detector, 875-UV or UV-970; mobile phase, hexane-2-propanol. EI mass spectra were measured on JEOL JMS-BU20 GCmate. FAB mass spectra were measured on JEOL MStation JMS-700. ESI-mass spectra were measured on Waters micromass ZQ. In general, reactions were carried out in dry solvents under an argon atmosphere, unless noted otherwise. Normal glassware can be used to conduct the reactions in this manuscript. Et₂AlCl was purchased from Kanto Kagaku.

Substrate screening for Reissert reaction of pyridine derivatives using Et₂AlCl as a catalyst:

Table S-1

entry	substrate	temp/°C	A/%	B/%
1	pyridine	rt	0	0
2		−78 to 0	40	—
3		−78 to 0	53	—
4		−40	82 (7:1) ^a	
5		−40	96 (3:1) ^a	
6		−78	45	46

^a The structure of regioisomers was not determined.

Synthesis of chiral ligands 4–6:

(*S,S*)-2,2'-Bis(methoxymethoxy)-3,3'-bis(phenylsulfanylmethyl)[1,1']binaphthalenyl (**D**):

To a solution of alcohol **C** (2.79 g, 6.44 mmol)¹ in pyridine (40 ml), diphenyldisulfide (3.51 g, 16.1 mmol), and *n*-tributylphosphine (4.01 ml, 16.1 mmol) were added, and the mixture was stirred at room temperature. After 1 h, the reaction was quenched by addition of water, and products were extracted with ethyl acetate for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (10% ethyl acetate in hexane) to give the sulfide **D** (3.75 g, 6.06 mmol, 94%) as colorless oil. ¹H NMR (CDCl₃) δ 2.90 (s, 6H), 4.34 (d, *J* = 7.0 Hz, 2H), 4.51 (m, 4H), 4.59 (d, *J* = 5.8 Hz, 2H), 7.03-7.45 (m, 16H), 7.76 (d, *J* = 8.3 Hz, 2H), 7.92 (s, 2H); ¹³C-NMR (CDCl₃) δ 33.9, 56.7, 99.3, 125.1, 125.4, 125.9, 126.1, 126.4, 127.7, 128.8, 129.4, 130.0, 130.5, 130.8, 133.4, 136.5, 152.6; [α]_D²⁵ = +59.17 (c 1.45, CHCl₃); IR (neat) ν 1157, 970 cm⁻¹; ESI-MS *m/z* 641 (M+Na)⁺; HRMS (FAB) Calcd for C₃₈H₃₅O₄S₂ 619.1977, Found 619.1967.

(*S,S,S,S*)-3,3'-Bis(benzenesulfinylmethyl)-2,2'-bis(methoxymethoxy)[1,1']binaphthalenyl (**E-4**):

(*S,R,S,R*)-3,3'-Bis(benzenesulfinylmethyl)-2,2'-bis(methoxymethoxy)[1,1']binaphthalenyl (**E-5**):

(*S,R,S,S*)-3,3'-Bis(benzenesulfinylmethyl)-2,2'-bis(methoxymethoxy)[1,1']binaphthalenyl (**E-6**):

To a solution of sulfide **D** (3.75 g, 6.06 mmol) in methylene chloride (60 ml), aqueous saturated NaHCO₃ (60 ml) was added, and the mixture was stirred at 0 °C. *m*-chloroperbenzoic acid (2.72 g, 12.12 mmol) was added slowly then stirred for 40 minutes. The reaction was quenched by addition of aqueous Na₂S₂O₃, and the products were extracted with methylene chloride for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (50% ethyl acetate in hexane) to give a mixture of sulfoxide **E-4**, **E-5** and **E-6** (3.95 g, 6.06 mmol, 100%) as colorless oil.

(*S,S,S,S*)-3,3'-Bis(benzenesulfinylmethyl)[1,1']binaphthalenyl-2,2'-diol (**4**):

(*S,R,S,R*)-3,3'-Bis(benzenesulfinylmethyl)[1,1']binaphthalenyl-2,2'-diol (**5**):

(*S,R,S,S*)-3,3'-Bis(benzenesulfinylmethyl)[1,1']binaphthalenyl-2,2'-diol (**6**):

The mixture of sulfoxides **E-4**, **E-5** and **E-6** (3.95 g, 6.06 mmol) was dissolved in methylene chloride (40 ml), and methanol (40 ml) and *p*-toluenesulfonic acid monohydrate (60.60 mmol, 11.53 g) were added. The reaction mixture was stirred at room temperature for 16 h, and quenched by addition of water. Then the products were extracted with methylene chloride for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate/acetic acid = 25/25/1) to give sulfoxide **4** (0.85 g, 1.51 mmol, 25%), **5** (0.85 g, 1.51 mmol, 25%), and **6** (1.59 g, 3.00 mmol, 50%) as colorless solids. **4**: ¹H NMR (CDCl₃) δ 4.15 (d, *J* = 12.5 Hz, 2H), 4.55 (d, *J* = 12.5 Hz, 2H), 7.00 (d, *J* = 8.6 Hz, 2H), 7.24 (t like, *J* = 7.6 Hz, 2H), 7.31 (t like, *J* = 7.6 Hz, 2H), 7.49 (m, 6H), 7.64 (m, 4H), 7.73 (s, 2H), 7.78 (d, *J* = 8.3 Hz, 2H); R_f = 0.38 (hexane/ethyl acetate/acetic acid = 25/25/1); **5**: ¹H NMR (CDCl₃) δ 4.25 (d, *J* = 13.5 Hz, 2H), 4.48 (d, *J* = 13.5 Hz, 2H), 6.92 (d, *J* = 8.5 Hz, 2H), 7.18-7.25 (m, 4H), 7.25-

7.48 (m, 6H), 7.63 (d, $J = 8.2$ Hz, 2H); $R_f = 0.17$ (hexane/ethyl acetate/acetic acid = 25/25/1); **6**: $^1\text{H-NMR}$ (CDCl_3) δ 4.30 (t, brd, $J = 13.1$ Hz, 2H), 4.48-4.56 (d x 2, $J = 13.1$ Hz, 2H), 7.0 (d, $J = 8.6$ Hz, 2H), 7.23-7.33 (m, 4H), 7.42-7.46 (m, 6H), 7.52-7.56 (m, 6H), 7.71 (d, $J = 8.3$ Hz, 2H); $^{13}\text{C-NMR}$ (CDCl_3) δ 59.32, 59.46, 100.53, 114.99, 115.16, 119.73, 119.85, 124.07, 124.11, 124.19, 124.22, 124.55, 127.34, 128.12, 128.70, 128.76, 128.99, 129.09, 131.34, 132.72, 132.76, 133.76, 141.94, 142.09, 151.23, 151.25; $[\alpha]_D^{21} = -63.07$ (c 1.01, CHCl_3); IR (neat) ν 3517, 3059, 3003, 1625, 1600, 1443, 1216, 1013, 995, 747 cm^{-1} ; ESI-MS m/z 563 ($\text{M}+\text{H}$) $^+$, 585 ($\text{M}+\text{Na}$) $^+$; HRMS (FAB) Calcd for $\text{C}_{34}\text{H}_{27}\text{O}_4\text{S}_2$ 563.1351, Found 563.1350; $R_f = 0.23$ (hexane/ethyl acetate/acetic acid = 25/25/1).

The non C_2 -symmetric structure of **6** was assigned based on ^1H and ^{13}C NMR. The structure of C_2 -symmetric ligand **5** was determined by X-ray crystallography, which led also to unequivocal structure determination of the other C_2 -symmetric ligand **4**.

Synthesis of chiral ligand **3**:

(*S*)-3,3'-Bis[bis(2-ethylphenyl)phosphinothiomethyl]-2,2'-bis(methoxymethoxy)[1,1']binaphthalenyl (**G**):

To a solution of 1-bromo-2-ethylbenzene (5.16 g, 0.028 mol) in ether (100 ml) at 0 °C, $t\text{-BuLi}$ (1.45 M in pentane, 23.4 ml, 0.034 mol) was added, and the mixture was stirred for 25 minutes. Then PCl_3 (0.81 ml, 9.24 mmol) was added, and the reaction mixture was stirred for 10 h. The reaction was quenched by addition of aqueous saturated ammonium chloride, and products were extracted with ethyl acetate for 3 times. The organic phase was washed with brine, dried over Na_2SO_4 and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (hexane) to give tris(2-ethylphenyl)phosphine (0.95 g, 2.75 mmol, 30%) as a colorless solid.

Tris(2-ethylphenyl)phosphine (0.57 g, 1.65 mmol) was carefully purged with argon, and degassed THF (5 ml) and lithium (wire, 57.3 mg, 8.25 mmol) were added. The reaction mixture was stirred at room temperature vigorously. After 11 hours, t -butyl chloride (0.18 ml, 1.65 mmol) was added to the reaction mixture and stirred for additional 1 h. In another vessel, compound **F** (0.28 g, 0.59 mmol)¹ was dissolved in degassed THF (4 ml), and cooled at 0 °C. The lithium phosphide solution prepared as described above was added to a solution of **F** via cannula, and the mixture was stirred for 30 minutes. Finally, sulfur (0.53 g, 16.50 mmol) was added in one portion, stirred for additional 30 minutes and quenched by addition of water. The reaction mixture was extracted with ethyl acetate for 3 times and the organic phase was washed with brine, dried over Na_2SO_4 and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (15% ethyl acetate in hexane) to give compound **G** (0.37 g, 0.39 mmol, 66%) as a colorless solid. ^1H NMR (CDCl_3) δ 0.80 (d x 2, $J = 7$ Hz, 12H), 2.59-2.70 (m, 8H), 2.96 (s, 6H), 4.29 (t, $J = 13.5$ Hz, 2H), 4.43 (m, 2H), 4.59 (t, $J = 14.1$ Hz, 2H), 6.85 (d, $J = 8.3$ Hz, 2H), 7.12 (t, $J = 7.9$ Hz, 2H), 7.20-7.33 (m, 10H), 7.44 (m, 4H), 7.55 (d, $J = 8.0$ Hz, 2H), 7.76 (s, 2H), 8.15-8.26 (m, 4H); ^{31}P NMR (CDCl_3) δ 39.80.

(*S*)-3,3'-Bis[bis(2-ethylphenyl)phosphinothiomethyl][1,1']binaphthalenyl-2,2'-diol (**3**):

To a solution of compound **G** (0.37 g, 0.39 mmol) in methylene chloride (2 ml) and methanol (2 ml), p -toluenesulfonic acid monohydrate (0.37 g, 1.95 mmol) was added, and the mixture was stirred at room temperature for 16 h. The reaction was quenched by addition of water, and extracted with ethyl acetate for 3 times. The organic phase was washed with brine, dried over Na_2SO_4 and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (25% ethyl acetate in hexane) to give the compound **3** (0.30 g, 0.35 mmol, 90%) as a colorless solid. $^1\text{H-NMR}$ (CDCl_3) δ 0.81 (t, $J = 7.3$ Hz, 6H), 0.84

(t, $J = 7.3$ Hz, 6H), 2.58-2.70 (m, 8H), 4.33 (t, $J = 13.7$ Hz, 2H), 4.39 (t, $J = 13.7$ Hz, 2H), 5.71 (s, 2H), 6.74 (d, $J = 4.8$ Hz, 2H), 7.16 (t, $J = 8.2$ Hz, 2H), 7.20-7.30 (m, 12H), 7.41 (t, $J = 7.6$ Hz, 2H), 7.45 (t, $J = 7.6$ Hz, 2H), 7.59 (m, 4H), 8.02-8.09 (m, 4H); ^{13}C -NMR (CDCl_3) δ 14.22, 14.29, 26.72, 26.76, 26.79, 26.83, 34.46, 34.88, 114.17, 120.28, 120.34, 123.78, 124.48, 125.89, 125.97, 125.99, 126.72, 126.99, 127.95, 128.66, 128.69, 129.67, 129.76, 130.01, 130.10, 131.80, 131.82, 132.54, 132.59, 132.79, 132.82, 146.86, 147.08, 150.84, 150.88; ^{31}P NMR (CDCl_3) δ 38.97; $[\alpha]_D^{25} = -88.00$ (c 1.03, CHCl_3); IR (neat) δ 3520, 3176, 3059, 2970, 2931, 2872, 1625, 1592, 1435, 1216, 1134, 1089, 748 cm^{-1} ; ESI-MS m/z 859 ($\text{M}+\text{H}^+$), 881 ($\text{M}+\text{Na}^+$); HRMS (FAB) Calcd for $\text{C}_{54}\text{H}_{52}\text{O}_2\text{P}_2\text{S}_2$ 859.2962, Found 859.2941.

In the similar manner, ligand **2** was synthesized.

3,3'-Bis(diphenylphosphinothiomethyl)[1,1']binaphthalenyl-2,2'-diol (2): ^1H -NMR (CDCl_3) δ 4.22 (quintet like, $J = 13.0$ Hz, 4H), 5.41 (brd, 2H), 6.81 (d, $J = 8.3$ Hz, 2H), 7.18 (t, $J = 7.5$ Hz, 2H), 7.29 (t, $J = 7.5$ Hz, 2H), 7.41-7.49 (m, 12H), 7.70 (d, $J = 8.3$ Hz, 2H), 7.85-7.91 (m, 10H); ^{13}C -NMR (CDCl_3) δ 34.7, 35.1, 113.1, 120.5, 124.1, 124.4, 127.0, 128.2, 128.4, 128.5, 128.6, 128.8, 131.4, 131.6, 132.2, 132.7, 151.7; ^{31}P NMR (CDCl_3) δ 43.80; $[\alpha]_D^{23} = -104.5$ (c 1.00, CHCl_3); IR (neat) δ 3520, 3179, 3056, 1436, 1103, 744, 706, 690 cm^{-1} ; MS (FAB ($m\text{NBA}$)) m/z 747 ($\text{M}+\text{H}^+$); (FAB ($m\text{NBA} + \text{aq } 1 \text{ N KI}$)) m/z 785 ($\text{M}+\text{K}^+$); HRMS (FAB) Calcd for $\text{C}_{46}\text{H}_{37}\text{O}_2\text{P}_2\text{S}_2$ ($\text{M}+\text{H}^+$) 747.1710, Found 747.1704.

Screening of the chiral ligands (Table S-2):

Results of ligand screening with regards to the Lewis base moiety is summarized in Table S-2. The catalyst prepared from Et_2AlCl and **6** in a 1:2 ratio gave higher regio- and enantioselectivities than catalysts prepared from other ligands. Chirality on the sulfur atom appeared to play a predominant role for enantioselectivity than the axial chirality of BINOL (entries 8 and 9 vs entries 10 and 11). Another catalyst that gave meaningful selectivity is a phosphine sulfide-containing catalyst (entry 4).

Table S-2

entry	ligand (Al:ligand)	time/h	8a yield/% (ee/%)	9a yield/% (ee/%)
1	none (1:0)	6	45 (–)	46 (–)
2	1 (1:1)	3	63 (2)	28 (9)
3	1 (1:2)	15	62 (2)	26 (16)
4	2 (1:1)	3	80 (38)	20 (10)
5	2 (1:2)	9	48 (2)	30 (11)
6	H (1:1)	3	44 (1)	46 (10)
7	I (1:1)	9	32 (0)	46 (4)
8	4 (1:1)	5	38 (8)	38 (9)
9	4 (1:2)	5	44 (15)	34 (7)
10	5 (1:1)	12	37 (11) ^a	40 (6) ^a
11	5 (1:2)	20	28 (14) ^a	28 (4) ^a
12	6 (1:1)	9	50 (7)	30 (14) ^a
13	6 (1:2)	5	90 (75)	8 (8)

^a The enantiomer was obtained as a major isomer.

1: X = P(O)Ph₂
 2: X = P(S)Ph₂
 3: X = P(S)(C₆H₄-o-Et)₂
 H: X = (CH₂)₄NO
 I: X = SO₂Ph

4: X, X' = O[−]; Y, Y' = Ph
 5: X, X' = Ph; Y, Y' = O[−]
 6: X, Y' = Ph; X', Y = O[−]

Screening of chloroformate using catalyst Al-6 (Table S-3):

Table S-3

entry	chloroformate	time/h	J yield/% (ee/%)	K yield/% (ee/%)
1	MeOCOCi	4	98 (91) ^d	2 (13)
2	EtOCOCi	5	90 (75)	8 (8)
3	AllocCl	3	77 (78)	11 (8)
4	ⁱ BuOCOCi	3	78 (75)	13 (9)
5	neopentylOCOCi	3	70 (70)	14 (14)
6	CbzCl	—	n.r. ^e	n.r. ^e
7 ^a	PhOCOCi	3	47 (42)	13 (8)
8	FmocCl	4	89 (93)	5 (9)
9 ^b	FmocCl	6	81 (89)	6 (8)
10 ^c	FmocCl	6	78 (87)	9 (0)

^a Reaction temperature = -40 °C. ^b Catalyst = 2.5 mol %. ^c Catalyst = 1 mol %.

^d Absolute configuration was determined as (S). ^e No reaction.

Excellent enantioselectivity was obtained using MeOCOCi (entry 1) and FmocCl (entry 8). Enantioselectivity was only slightly decreased using 2.5 (entry 9) or 1 mol % (entry 10) of catalyst.

Optimization using phosphine sulfide-type catalysts Al-2 and 3 (Table S-4):

Table S-4

entry	ligand	substrate (R')	chloroformate	time/h	J yield/% (ee/%)	K yield/% (ee/%)
1	2	Me	MeOCOCi	15	n.r. ^a	n.r. ^a
2	2	Me	EtOCOCi	3	71 (36)	23 (13)
3	2	Me	AllocCl	30	70 (40)	25 (11)
4	2	Me	ⁱ BuOCOCi	3.5	83 (37)	19 (55)
5	2	Me	neopentylOCOCi	3	61 (52)	16 (15)
6	2	Me	FmocCl	26	37 (13)	11 (49)
7	2	Bn	neopentylOCOCi	3	73 (48)	17 (15)
8	2	ⁱ Pr	neopentylOCOCi	3	68 (68)	32 (n.d.) ^b
9	L	ⁱ Pr	neopentylOCOCi	3	62 (47)	17 (0)
10	3	ⁱ Pr	neopentylOCOCi	3	56 (80)	15 (0)

^a No reaction. ^b Not determined.

Neopentyl chloroformate is optimum for the reaction using phosphine sulfide containing catalyst 2-Al (entries 1–6). Bulky diisopropylamide-containing substrate gave higher enantioselectivity than dimethyl or dibenzylamide-containing substrate (entry 5, 7, and 8). For the aryl part in the ligand, bis(*o*-ethylphenyl)phosphine sulfide-containing catalyst gave the best results (entry 5, 9, and 10).

Catalytic enantioselective Reissert reaction of pyridine derivatives using Al-6 (General Procedure):

Ligand 6 (0.02 mmol) was dried under reduced pressure for 1 h, and dissolved in methylene chloride (0.5 ml). Et₂AlCl (0.98 M in *n*-hexane, 10.2 μ l, 0.01 mmol) was added and stirred at room temperature for 1 h. The resulting mixture was cooled to -60 °C, and a starting material (0.4 M in methylene chloride, 0.5 ml, 0.2 mmol), a chloroformate (2.8 mmol) and TMSCN (53.3 μ l, 4.0 mmol) were added successively. After the reaction was completed, water was added. The products were extracted with methylene chloride for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting

residue was purified by silica gel column chromatography to give the Reissert product.

2-Cyano-5-dimethylcarbamoyl-2H-pyridine-1-carboxylic acid 9H-fluoren-9-ylmethyl ester

$^1\text{H-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 2.92 (s, 6H), 4.71 (t, J = 6.2 Hz, 1H), 4.88-5.08 (m, 2H), 6.06 (dd, J = 5.7, 9.1 Hz, 1H), 6.29 (d, J = 5.7 Hz, 1H), 6.65 (d, J = 9.1 Hz, 1H), 7.10 (brd, 1H), 7.64 (t, J = 7.4 Hz, 2H), 7.73 (t, J = 7.4 Hz, 2H), 7.97 (t, J = 6.8 Hz, 2H), 8.19 (d, J = 7.4 Hz, 2H); $^{13}\text{C-NMR}$ (CDCl_3) δ 37.4, 42.8, 46.6, 69.4, 112.1, 114.2, 115.5, 120.0, 124.6, 124.7, 126.0, 127.2, 127.9, 141.1, 142.7, 142.8, 151.7, 167.4; $[\alpha]_D^{21}$ = -205.13 (c 1.50, CHCl_3 , 93% ee); IR (neat) δ 3460, 2953, 1957, 1920, 1732, 1652, 1624, 1388, 1294, 1254, 1207, 759, 741 cm^{-1} ; ESI-MS m/z 400 ($\text{M}+\text{H}$) $^+$, 422 ($\text{M}+\text{Na}$) $^+$; HRMS (FAB) Calcd for $\text{C}_{24}\text{H}_{22}\text{N}_3\text{O}_3$ 400.1661, Found 400.1695; HPLC DAICEL CHIRALPAK OD-H, hexane/2-propanol = 4/1, flow rate: 1.5 ml/min, detector: 254 nm, t_R (major) = 40.1 min, t_R (minor) = 30.0 min.

2-Cyano-5-diisopropylcarbamoyl-2H-pyridine-1-carboxylic acid 9H-fluoren-9-ylmethyl ester

$^1\text{H-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 1.21 (brd, 12H), 3.71 (m, 2H), 4.40 (t, J = 5.8 Hz, 1H), 4.64 (brd, 1H), 4.74 (brd, 1H), 5.77 (t, J = 8.9 Hz, 1H), 5.98 (d, J = 5.8 Hz, 1H), 6.26 (d, J = 8.9 Hz, 1H), 6.53 (brd, 1H), 7.33 (t, J = 7.3 Hz, 2H), 7.41 (t, J = 7.3 Hz, 2H), 7.66 (d, J = 6.4 Hz, 2H), 7.86 (d, J = 7.6 Hz, 2H); $^{13}\text{C-NMR}$ (CDCl_3) δ 20.8, 21.0, 42.8, 46.8, 69.3, 115.7, 120.1, 120.2, 124.6, 124.7, 127.3, 128.0, 141.3, 142.8, 142.9, 152.0, 166.7; $[\alpha]_D^{20}$ = -153.33 (c 0.93, CHCl_3 , 90% ee); IR (neat) δ 3462, 3092, 2958, 1736, 1653, 1624, 1442, 1371, 1297, 1255, 1211 cm^{-1} ; ESI-MS m/z 456 ($\text{M}+\text{H}$) $^+$, 478 ($\text{M}+\text{Na}$) $^+$; HRMS (FAB) Calcd for $\text{C}_{28}\text{H}_{30}\text{N}_3\text{O}_3$ ($\text{M}+\text{H}$) $^+$ 456.2287, Found 456.2285; HPLC DAICEL CHIRALPAK AD-H, hexane/2-propanol = 9/1, flow rate: 1.0 ml/min, detector: 254 nm, t_R (major) = 38.5 min, t_R (minor) = 42.1 min.

$^1\text{H-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 1.03 (s, 9H), 1.32 (d, J = 6.4 Hz, 12H), 3.84 (m, 2H), 4.02 (m, 2H), 5.86 (m, J = 9.5 Hz, 1H), 6.14 (d, J = 5.8 Hz, 1H), 6.36 (d, J = 9.5 Hz, 1H), 6.97 (s, 1H); $^{13}\text{C-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 20.4, 25.8, 31.0, 42.7, 47.5, 76.2, 114.2, 116.3, 116.4, 123.9, 124.8, 151.7, 165.9; $[\alpha]_D^{25}$ = -146.3 (c 1.88, CHCl_3 , 94% ee); IR (neat) δ 2966, 2234(w), 1733, 1623, 1279, 1257 cm^{-1} ; MS(FAB(m -NBA)) m/z 348 ($\text{M}+\text{H}$) $^+$, 321 ($\text{M}-\text{CN}$) $^+$, (FAB (m -NBA + aq 1 N KI)) m/z 386 ($\text{M}+\text{K}$) $^+$; HRMS (FAB) Calcd for $\text{C}_{19}\text{H}_{30}\text{N}_3\text{O}_3$ ($\text{M}+\text{H}$) $^+$ 348.2287, Found 348.2281; HPLC DAICEL CHIRALPAK AS, hexane/2-propanol = 20/1, flow rate: 1.0 ml/min, detector: 254 nm, t_R (major) = 13.0 min, t_R (minor) = 16.3 min.

2-Cyano-5-diisopropylcarbamoyl-2H-pyridine-1-carboxylic acid 2,2-dimethylpropyl ester

$^1\text{H-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 1.03 (s, 9H), 1.32 (d, J = 6.4 Hz, 12H), 3.84 (m, 2H), 4.02 (m, 2H), 5.86 (m, J = 9.5 Hz, 1H), 6.14 (d, J = 5.8 Hz, 1H), 6.36 (d, J = 9.5 Hz, 1H), 6.97 (s, 1H); $^{13}\text{C-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 20.4, 25.8, 31.0, 42.7, 47.5, 76.2, 114.2, 116.3, 116.4, 123.9, 124.8, 151.7, 165.9; $[\alpha]_D^{25}$ = -146.3 (c 1.88, CHCl_3 , 94% ee); IR (neat) δ 2966, 2234(w), 1733, 1623, 1279, 1257 cm^{-1} ; MS(FAB(m -NBA)) m/z 348 ($\text{M}+\text{H}$) $^+$, 321 ($\text{M}-\text{CN}$) $^+$, (FAB (m -NBA + aq 1 N KI)) m/z 386 ($\text{M}+\text{K}$) $^+$; HRMS (FAB) Calcd for $\text{C}_{19}\text{H}_{30}\text{N}_3\text{O}_3$ ($\text{M}+\text{H}$) $^+$ 348.2287, Found 348.2281; HPLC DAICEL CHIRALPAK AS, hexane/2-propanol = 20/1, flow rate: 1.0 ml/min, detector: 254 nm, t_R (major) = 13.0 min, t_R (minor) = 16.3 min.

$^1\text{H-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 3.73 (s, 3H), 5.75 (t, J = 5.7 Hz, 1H), 4.65-4.78 (m, 2H), 5.76 (dd, J = 6.3, 9.7 Hz, 1H), 6.00 (d, J = 5.2 Hz, 1H), 6.53 (dd, J = 1.1, 9.7 Hz, 1H), 7.34 (dt, J = 1.1, 7.4 Hz, 2H), 7.43 (t, J = 7.4 Hz, 2H), 7.67 (t, J = 5.7 Hz, 2H), 7.88 (d, J = 7.4 Hz, 2H), 8.51 (d, J = 1.7 Hz, 1H); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 43.4, 46.4, 51.8, 69.3, 108.4, 114.3, 115.7, 116.3, 120.3, 122.7, 125.0, 127.3, 127.9, 133.2, 140.86, 140.88, 143.1, 143.2, 149.8, 151.4, 164.2; $[\alpha]_D^{21}$ = -300.69 (c 0.87, CHCl_3 , 99% ee); IR (neat) δ 3423, 2952, 1967, 1915, 1739, 1712, 1650, 1592, 1449, 1392, 1335, 1234, 1133, 1090, 759, 742 cm^{-1} ; ESI-MS m/z 409 ($\text{M}+\text{Na}$) $^+$; HRMS (FAB) Calcd for $\text{C}_{23}\text{H}_{19}\text{N}_2\text{O}_4$ ($\text{M}+\text{H}$) $^+$ 387.1345 Found 387.1362; HPLC DAICEL CHIRALPAK AS-H, hexane/2-propanol = 9/1, flow rate: 1.0 ml/min, detector: 254 nm, t_R (major) = 44.1 min, t_R (minor) = 31.8 min.

6-Cyano-6H-pyridine-1,3-dicarboxylic acid 1-(9H-fluoren-9-ylmethyl) ester 3-methyl ester

$^1\text{H-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 3.73 (s, 3H), 5.75 (t, J = 5.7 Hz, 1H), 4.65-4.78 (m, 2H), 5.76 (dd, J = 6.3, 9.7 Hz, 1H), 6.00 (d, J = 5.2 Hz, 1H), 6.53 (dd, J = 1.1, 9.7 Hz, 1H), 7.34 (dt, J = 1.1, 7.4 Hz, 2H), 7.43 (t, J = 7.4 Hz, 2H), 7.67 (t, J = 5.7 Hz, 2H), 7.88 (d, J = 7.4 Hz, 2H), 8.51 (d, J = 1.7 Hz, 1H); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 43.4, 46.4, 51.8, 69.3, 108.4, 114.3, 115.7, 116.3, 120.3, 122.7, 125.0, 127.3, 127.9, 133.2, 140.86, 140.88, 143.1, 143.2, 149.8, 151.4, 164.2; $[\alpha]_D^{21}$ = -300.69 (c 0.87, CHCl_3 , 99% ee); IR (neat) δ 3423, 2952, 1967, 1915, 1739, 1712, 1650, 1592, 1449, 1392, 1335, 1234, 1133, 1090, 759, 742 cm^{-1} ; ESI-MS m/z 409 ($\text{M}+\text{Na}$) $^+$; HRMS (FAB) Calcd for $\text{C}_{23}\text{H}_{19}\text{N}_2\text{O}_4$ ($\text{M}+\text{H}$) $^+$ 387.1345 Found 387.1362; HPLC DAICEL CHIRALPAK AS-H, hexane/2-propanol = 9/1, flow rate: 1.0 ml/min, detector: 254 nm, t_R (major) = 44.1 min, t_R (minor) = 31.8 min.

Catalytic enantioselective Reissert reaction of pyridine derivatives using Al-3 (General Procedure):

Ligand **3** (0.02 mmol) was dried under reduced pressure for 1 h, and dissolved in methylene chloride (1.5ml). Et_2AlCl (0.98M in *n*-hexane, 20.4 μl , 0.02 mmol) was added and stirred at room temperature for 1 h. The resulting mixture was cooled to -60 $^\circ\text{C}$, and a starting material (0.4 M in methylene chloride, 0.5 ml, 0.2

mmol), a chloroformate (2.8 mmol) and TMSCN (53.3 μ l, 4.0 mmol) were added successively. After the reaction was completed, water was added. The products were extracted with methylene chloride for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography to give the Reissert product.

4-Chloro-2-cyano-5-diisopropylcarbamoyl-2H-pyridine-1-carboxylic acid 2,2-dimethylpropyl ester

¹H-NMR (DMSO, 50 °C) δ 0.96 (s, 9H), 1.27 (s, 12H), 3.5–4.1 (brd, 2H), 3.9 (d, *J* = 10.3 Hz, 1H), 4.0 (d, *J* = 10.3 Hz, 1H), 5.97 (d, *J* = 6.9 Hz, 1H), 6.15 (d, *J* = 6.9 Hz, 1H), 6.92 (s, 1H); ¹³C-NMR (DMSO) δ 20.0 (brd), 25.9, 31.2, 44.4, 45.2, 50.8, 76.5, 112.0 (brd), 116.1, 123.5 (brd), 126.0 (brd), 128.8, 151.5, 162.7; $[\alpha]^{23}_D = -175.0$ (c 0.68, CHCl₃, 91% ee); IR (neat) ν 3452, 2076, 1634, 759 cm⁻¹; ESI-MS *m/z* 304 (M+Na)⁺; HRMS (FAB) Calcd for C₁₉H₂₉ClN₃O₃ (M+H)⁺, 382.1897 Found 382.1893; HPLC

DAICEL CHIRALPAK OD-H, hexane/2-propanol = 20/1, flow rate: 1.0 ml/min, detector: 254 nm, *t_R*(major) = 24.4 min, *t_R*(minor) = 17.3 min.

4-Bromo-2-cyano-5-diisopropylcarbamoyl-2H-pyridine-1-carboxylic acid 2,2-dimethylpropyl ester

¹H-NMR (C₆D₆, 50 °C) δ 0.69 (s, 9H), 0.9–1.6 (brd, 12H), 3.0–4.2 (brd, 2H), 3.58 (d, *J* = 10.4 Hz, 1H), 3.78 (brd, 1H), 4.86 (dd, *J* = 1.7, 6.9 Hz, 1H), 5.28 (brd, 1H), 7.05 (brd, 1H); ¹³C-NMR (C₆D₆, 50 °C) δ 19.9, 20.7, 26.1, 31.3, 44.5 (brd), 46.0, 51.1, 77.1, 110.0, 119.0 (brd), 115.7, 126.2, 131.2, 151.8, 163.2; $[\alpha]^{22}_D = -24.6$ (c 0.2, CHCl₃, 86% ee); IR (neat) ν 3435, 2084, 1635, 761 cm⁻¹; MS (EI) *m/z* 381 (M-C₃H₈)⁺; HPLC

DAICEL CHIRALPAK AD-H, hexane/2-propanol = 9/1, flow rate: 1.0 ml/min,

detector: 254 nm, *t_R*(major) = 60.7 min, *t_R*(minor) = 47.2 min.

Application to catalytic enantioselective synthesis of CP-293,019:

2-Cyano-5-dimethylcarbamoyl-2H-pyridine-1-carboxylic acid methyl ester

300 mg-scale catalytic enantioselective Reissert reaction using Al-6 system: Ligand **6** (112.6 mg, 0.2 mmol) was dried under reduced pressure for 1 h, and dissolved in methylene chloride (2.5 ml). Et₂AlCl (0.98M in *n*-hexane, 105.3 μ l, 0.1 mmol) was added, and the mixture was stirred at room temperature for 1 h. The resulting mixture was cooled to -60 °C, and *N,N*-dimethylnicotinamide (0.4 M in methylene chloride, 2.5 ml, 2.0 mmol), methyl chloroformate (0.22 ml, 2.8 mmol) and TMSCN (0.53 ml, 4.0 mmol) were added successively. After 14 hours, the reaction was quenched with water. The resulting mixture was extracted with methylene chloride for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (25% acetone in hexane) to give the product **8b** (438.9 mg, 1.87 mmol, 94%, 86% ee) as a colorless solid. ¹H-NMR (DMSO-*d*₆, 50 °C) δ 3.00 (s, 6H), 3.90 (s, 3H), 5.81 (brd, *J* = 5.8 Hz, 1H), 6.11 (brd, *J* = 6.0 Hz, 1H), 6.43 (brd, *J* = 8.8 Hz, 1H), 7.15 (s, 1H); ¹³C-NMR (DMSO-*d*₆, 50 °C) δ 42.7, 54.4, 79.0, 113.4, 114.1, 116.4, 125.0, 127.0, 152.2, 166.5; $[\alpha]^{25}_D = -328.9$ (c 1.65, CHCl₃, 82% ee); IR (neat) ν 3462, 3092, 2958, 1736, 1653, 1624, 1442, 1371, 1297, 1255, 1211 cm⁻¹; MS (EI) *m/z* 235 (M⁺); HRMS (EI) Calcd for C₁₁H₁₃N₃O₃ 235.0957, Found 235.0959; HPLC DAICEL CHIRALPAK AD-H, hexane/2-propanol = 4/1, 1.0 ml/min, detector 254 nm, *t_R*(major) = 22.7 min, *t_R*(minor) = 18.3 min.

2-Cyano-5-dimethylcarbamoyl-3,4-dihydro-2H-pyridine-1-carboxylic acid methyl ester

Reissert product **8b** (0.91 g, 3.87 mmol) was purged with argon. Then 10% palladium on charcoal (15 mol%, 0.61 g) and methanol (40 ml) were added, and the reaction vessel was purged with hydrogen. The reaction mixture was stirred at room temperature for 15 minutes, filtered through celite pad, washed with methanol, and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (50% acetone in hexane) to give product **L** (0.81 g, 3.42 mmol, 88%) as colorless oil. $^1\text{H-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 2.00 (m, 1H), 2.32 (m, 3H), 2.98 (s, 6H), 3.84 (s, 3H), 5.43 (s, 1H), 6.98 (s, 1H); $^{13}\text{C-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 19.3, 23.5, 36.5, 42.0, 53.8, 79.0, 113.7, 117.8, 124.1, 152.1, 168.8; $[\alpha]_D^{22} = -20.53$ (c 0.76, CHCl_3 , 86% ee); IR (neat) 3446, 2958, 2243, 1727, 1656, 1617, 1443, 1356, 1314, 1193, 1083 cm^{-1} ; MS (EI) m/z 237 (M^+); HRMS (EI) Calcd for $\text{C}_{11}\text{H}_{15}\text{N}_3\text{O}_3$ 237.1113, Found 237.1094; HPLC DAICEL CHIRALPAK AD-H, hexane/2-propanol = 4/1, flow rate: 1.0 ml/min, detector: 254 nm, $t_{\text{R}}(\text{major}) = 17.5$ min, $t_{\text{R}}(\text{minor}) = 14.0$ min.

5-Dimethylcarbamoyl-2-(toluene-4-sulfonylamino)methyl-3,4-dihydro-2H-pyridine-1-carboxylic acid methyl ester

Compound **L** (0.81 g, 3.42 mmol) was purged with argon. Then 10% palladium on charcoal (20 mol %, 0.72 g), ethanol (27 ml), conc. HCl (3 ml) were added, and the vessel was purged with hydrogen. The reaction mixture was stirred at room temperature for 1 h, filtered through celite pad, washed with ethanol, and concentrated *in vacuo*. To the resulting residue in methylene chloride (30 ml) at 0 $^\circ\text{C}$, Et_3N (1.43 ml, 10.26 mmol) and *p*-toluenesulfonyl chloride (0.78 g, 4.10 mmol) were added. The reaction mixture was stirred for 1 h. Water was added, and the organic phase was separated. The water phase was extracted with methylene chloride twice, and the combined organic phase was washed with brine, dried over Na_2SO_4 and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (33% hexane in ethyl acetate) to give the tosyl amide **M** (1.11 g, 2.81 mmol, 82%) as colorless oil. $^1\text{H-NMR}$ (DMSO- d_6 , 50 $^\circ\text{C}$) δ 1.66 (m, 1H), 2.15-2.22 (m, 3H), 2.49 (s, 3H), 2.87 (m, 1H), 3.02 (s, 6H), 3.79 (s, 3H), 4.32 (m, 1H), 7.00 (brd, 1H), 7.49 (d, $J = 8.1$ Hz, 2H), 7.78 (d, $J = 8.1$ Hz, 2H), 7.81 (brd, 1H); $^{13}\text{C-NMR}$ (CDCl_3) δ 18.6, 21.5, 22.0, 44.1, 49.4, 53.6, 77.2, 112.5, 126.0, 126.9, 129.7, 136.9, 143.4, 154.3, 170.9; $[\alpha]_D^{23} = -36.94$ (c 1.64, CHCl_3 , 87% ee); IR (neat) 3442, 3171, 2952, 2930, 2866, 1717, 1654, 1601, 1442, 1397, 1308, 1261, 1195, 1159 cm^{-1} ; MS (EI) m/z 395 (M^+); HRMS (EI) Calcd for $\text{C}_{18}\text{H}_{25}\text{N}_3\text{O}_5\text{S}$ 395.1515, Found 395.1523; HPLC DAICEL CHIRALPAK OD-H, hexane/2-propanol = 6/1, flow rate: 0.85 ml/min, detector: 254 nm, $t_{\text{R}}(\text{major}) = 50.0$ min, $t_{\text{R}}(\text{minor}) = 63.0$ min.

6-(Toluene-4-sulfonylamino)methyl-1,4,5,6-tetrahydropyridine-3-carboxylic acid dimethylamide

Tosyl amide **M** (1.11 g, 2.81 mmol) was dissolved in methanol (20 ml), and aqueous 3 M NaOH (10 ml) was

added. The reaction mixture was refluxed for 1.5 h, then cooled to room temperature. The reaction mixture was acidified with aqueous 1 N HCl and extracted with ethyl acetate. Then the water phase was basified with 25% ammonia solution, and extracted with ethyl acetate for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (33% hexane in acetone) to give **19** (0.64 g, 1.90 mmol, 68%) as colorless oil. ¹H-NMR (CDCl₃) δ 1.54 (m, 1H), 1.74 (m, 1H), 2.20-2.38 (m, 2H), 2.43 (s, 3H), 2.85-3.20 (m, 2H), 2.97 (s, 6H), 3.36 (m, 1H), 4.57 (d, *J* = 4.9 Hz, 1H), 5.74 (t, *J* = 6.1 Hz, 1H), 6.71 (d, *J* = 5.5 Hz, 1H), 7.30 (d, *J* = 8.4 Hz, 2H), 7.75 (d, *J* = 8.4 Hz, 2H); ¹³C-NMR (CDCl₃) δ 37.8, 37.9, 47.5, 49.8, 49.9, 98.6, 126.8, 126.9, 129.5, 129.6, 137.1, 138.3, 138.4, 143.1, 173.5; [α]_D²³ = +43.34 (c 1.86, CHCl₃, 89% ee); IR (neat) 3401, 3296, 3136, 2926, 2856, 1631, 1566, 1494, 1447, 1392, 1327, 1195, 1158, 1092 cm⁻¹; MS (EI) *m/z* 337 (M⁺); HRMS (EI) Calcd for C₁₆H₂₃N₃O₃S 337.1460, Found 337.1437; HPLC DAICEL CHIRALPAK AD-H, hexane/2-propanol = 4/1, flow rate: 1.0 ml/min, detector: 254 nm, *t*_R(major) = 21.5 min, *t*_R(minor) = 31.7 min.

2-(Toluene-4-sulfonyl)octahydropyrido[1,2-*a*]pyrazine-7-carboxylic acid dimethylamide

Compound **19** (0.29 g, 0.86 mmol) was dissolved in methanol (4 ml) and acetic acid (4 ml) at -20 °C. Sodium cyanoborohydride (81.1 mg, 1.29 mmol) was added. After 2.5 h the reaction mixture was slowly poured into aqueous saturated NaHCO₃, and the products were extracted with methylene chloride for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (20% methanol in chloroform) to give amine (*trans*: 176.3 mg, 0.52 mmol, 60%, *cis*: 44.1 mg, 0.13 mmol, 15%). To the *trans*-amine (176.3 mg, 0.52 mmol) in acetonitrile (8 ml), cesium carbonate (1.40 g, 4.3 mmol) and 1,2-dibromoethane (1.5 ml, 3.66 mmol) were added. The reaction mixture was refluxed for 1 h, then cooled to room temperature and quenched with water. The reaction mixture was extracted with ethyl acetate for 3 times. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (33% hexane in acetone) to give **20** (0.17 g, 0.48 mmol, 92%) as colorless solid. ¹H-NMR (CDCl₃) δ 1.13 (m, 1H), 1.50-1.70 (m, 2H), 1.77 (m, 1H), 1.97 (t, *J* = 10.6 Hz, 1H), 2.08 (m, 1H), 2.28 (t, *J* = 11.3 Hz, 1H), 2.33-2.42 (m, 2H), 2.38 (s, 3H), 2.75 (m, 3H), 2.87 (s, 3H), 2.97 (s, 3H), 3.52 (dt, *J* = 2.4, 10.9 Hz, 1H), 3.61 (m, 1H), 7.27 (d, *J* = 8.1 Hz, 2H), 7.58 (d, *J* = 8.1 Hz, 2H); ¹³C-NMR (CDCl₃) δ 21.4, 26.5, 28.5, 35.4, 36.9, 39.2, 45.7, 51.0, 53.7, 56.7, 59.3, 127.8, 129.5, 132.1, 143.6, 173.1; [α]_D²² = +8.17 (c 1.03, CHCl₃, 89% ee); IR (neat) 2935, 2851, 1638, 1347, 1165 cm⁻¹; ESI-MS 366 (M+H)⁺, 388 (M+Na)⁺; Anal. Calcd for C₁₈H₂₇N₃O₃S C: 59.15 H: 7.45 N: 11.50, Found C: 59.18 H: 7.35 N: 11.41; HPLC DAICEL CHIRALPAK AD-H, hexane/2-propanol = 4/1, flow rate: 1.0 ml/min, detector: 254 nm, *t*_R(major) = 18.1 min, *t*_R(minor) = 25.9 min.

[2-(Toluene-4-sulfonyl)octahydropyrido[1,2-*a*]pyrazin-7-yl] methanol

To borane-ammonia complex (64.5 mg, 1.88 mmol) in THF (3 ml) at 0 °C, *n*-BuLi (1.58 M in *n*-hexane, 1.19 ml, 1.88 mmol) was added, and the mixture was stirred for 5 minutes. Then the reaction mixture was stirred at room temperature for 5 minutes. After cooling to 0 °C, amide **20** (0.17 g, 0.47 mmol) in THF (3 ml) was

added, and the mixture was stirred for 4 h. The reaction was quenched by the addition of water, and the products were extracted with ethyl acetate for 3 times. The organic phase was washed with brine, dried over Na_2SO_4 and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (10% methanol in chloroform) to give alcohol **N** (102.4 mg, 0.32 mmol, 68%) as a colorless solid. ^1H -NMR (CDCl_3) δ 1.01 (m, 1H), 1.15 (m, 1H), 1.63 (d, $J = 12.0$, 1H), 1.77 (m, 3H), 2.01 (m, 2H), 2.35 (dt, $J = 2.9$, 11.5 Hz, 1H), 2.42 (s, 3H), 2.43 (m, 1H), 2.76 (brd d, $J = 11.5$ Hz, 1H), 2.91 (brd d, $J = 9.8$ Hz, 1H), 3.40 (dd, $J = 6.3$, 10.9 Hz, 1H), 3.49 (dd, $J = 5.2$, 10.9 Hz, 1H), 3.55 (m, 1H), 3.65 (m, 1H), 7.32 (d, $J = 8.3$ Hz, 2H), 7.62 (d, $J = 8.3$ Hz, 2H); ^{13}C -NMR (CDCl_3) δ 21.5, 26.5, 28.8, 38.9, 45.8, 51.1, 54.0, 58.1, 60.0, 66.0, 127.8, 129.6, 132.1, 143.7; $[\alpha]_{\text{D}}^{25} +23.9$ (c 1.16, CHCl_3 , 79% ee); IR (neat) 3265, 2926, 2852, 1456, 1347, 1332, 1166 cm^{-1} ; MS (EI) m/z 324 (M^+), 169 (M-Ts^+); HRMS (EI) Calcd for $\text{C}_9\text{H}_{17}\text{N}_2\text{O}$ (M-Ts) 169.1341, Found 169.1248; HPLC DAICEL CHIRALPAK OD-H, hexane/2-propanol = 9/1, flow rate: 0.85 ml/min, detector: 254 nm, t_{R} (major) = 22.5 min, t_{R} (minor) = 26.2 min.

7-Hydroxymethyloctahydropyrido[1,2-a]pyrazine-2-carboxylic acid *tert*-butyl ester

Naphthalene (1.05 g, 8.19 mmol) was dissolved in DME (5 ml) at room temperature, and sodium (0.15 g, 1.17 mmol) was added. The naphthalenide solution was stirred vigorously for 1 hour. Alcohol **N** (102.4 mg, 0.32 mmol) was dissolved in DME (3 ml), stirred at -60°C and added slowly to the naphthalenide solution until the reaction mixture shows pale green (total ca. 1 ml). The reaction was quenched with water (3 ml), and the water phase was washed with ethyl acetate. Then the water phase was concentrated *in vacuo*. The resulting aminoalcohol was treated with Boc_2O (104.8 mg, 0.48 mmol), DMAP (58.6 mg, 0.48 mmol) and acetonitrile (3 ml) without purification. The reaction mixture was stirred at 70°C for 30 minutes, then quenched with water. The reaction mixture was extracted with ethyl acetate for 3 times. The organic phase was washed with brine, dried over Na_2SO_4 and concentrated *in vacuo*. The resulting residue was purified by silica gel column chromatography (10% methanol in chloroform) to give **21** (37.6 mg, 0.14 mmol, 44% in 2 steps) as a colorless solid. ^1H -NMR (CDCl_3) δ 1.03 (dd, $J = 3.5$, 12.8 Hz, 1H), 1.24 (dd, $J = 4.0$, 12.8 Hz, 1H), 1.46 (s, 9H), 1.63 (m, 2H), 1.77-1.89 (m, 4H), 2.13 (dt, $J = 2.9$, 12.0 Hz, 1H), 2.57 (brd, 1H), 2.74 (d, $J = 10.9$ Hz, 1H), 2.97 (d, $J = 10.9$ Hz, 2H), 3.43 (dd, $J = 7.0$, 11.0 Hz, 1H), 3.51 (dd, $J = 5.2$, 11.0 Hz, 1H), 3.95 (brd 2H); ^{13}C -NMR (CDCl_3) δ 26.8, 28.5, 28.8, 39.1, 43.5 (brd), 49.0 (brd), 54.9, 58.7, 60.9, 66.1, 79.9, 154.7; $[\alpha]_{\text{D}}^{28} -0.76$ (c 0.93, CHCl_3 , 84% ee); IR (neat) 3400, 2925, 1697, 1424, 1272, 1243, 1272, 1242, 1171, 1126 cm^{-1} ; ESI-MS m/z 271 ($\text{M}+\text{H}^+$), 293 ($\text{M}+\text{Na}^+$); HRMS (FAB) Calcd for $\text{C}_{14}\text{H}_{27}\text{N}_2\text{O}_3$ ($\text{M}+\text{H}^+$) 271.2022, Found 271.2026; HPLC DAICEL CHIRALPAK AS-H, hexane/2-propanol = 20/1, flow rate: 1.0 ml/min, detector: 210 nm, t_{R} (major) = 14.9 min, t_{R} (minor) = 18.5 min. The absolute configuration was determined comparing the sign of optical rotation.³

Other useful conversions:

3-Oxo-1,2,3,7,8,8a-hexahydroimidazo[1,5-a]pyridine-6-carboxylic acid dimethylamide

To a solution of starting material **L** (30.6 mg, 0.13 mmol) in methanol (1.3 ml), Mg (powder, 25.3 mg, 1.04 mmol) was added and the mixture was stirred at room temperature. After 12 hours, the reaction mixture was added to aqueous 1N HCl and concentrated *in vacuo*. The resulting residue was purified by silica gel column

chromatography (10% methanol in chloroform) to give **O** (16.4 mg, 0.078 mmol, 60%) as colorless oil. $^1\text{H-NMR}$ (CDCl_3) δ 1.61 (m, 1H), 2.15 (m, 1H), 2.36 (m, 1H), 2.55 (dd, $J = 5.2, 17.7$ Hz, 1H), 3.02 (s, 6H), 3.20 (t, $J = 8.9$ Hz, 1H), 3.67 (t, $J = 8.6$ Hz, 1H), 3.97 (m, 1H), 5.29 (s, 1H), 6.94 (d, $J = 1.9$ Hz, 1H); $^{13}\text{C-NMR}$ (CDCl_3) δ 22.8, 26.3, 37.6, 45.1, 52.7, 110.0, 125.0, 157.5, 171.2; $[\alpha]_D^{24} +117.85$ (c 1.21, CHCl_3); IR (neat) 3412, 2925, 1704, 1640, 1592, 1430, 1390, 1327, 1257, 1192, 737 cm^{-1} ; ESI-MS m/z 210 ($\text{M}+\text{H}$) $^+$, 232 ($\text{M}+\text{Na}$) $^+$; HRMS (FAB) Calcd for $\text{C}_{10}\text{H}_{16}\text{N}_3\text{O}_2$ ($\text{M}+\text{H}$) $^+$ 210.1243, Found 210.1249.

2-Diisopropylcarbamoyl-8-azabicyclo[3.2.1]octane-8-carboxylic acid 2,2-dimethylpropyl ester

P (22.9 mg, 0.065 mmol) was dissolved in THF (0.7 ml), and stirred at room temperature. Boranetrifluoride diethylether complex (0.05 ml, 0.39 mmol) was added and after 2 h the reaction was quenched by the addition of methanol. The reaction mixture was concentrated *in vacuo* and purified by silica gel column chromatography (25% methanol in chloroform) to give **Q** (20.0 mg, 0.057 mmol, 88%) as a colorless solid. $^1\text{H-NMR}$ (CDCl_3) δ 0.96 (s, 9H), 1.31 (d, $J = 6.4$ Hz, 12H), 1.87 (m, 1H), 2.30-2.70 (m, 2H), 2.96 (m, 1H), 3.07 (m, 1H), 3.74 (brd, 2H), 3.86 (d, $J = 10.1$ Hz, 1H), 3.90 (d, $J = 10.1$ Hz, 1H), 4.39 (brd, 1H), 4.96 (brd, 1H), 5.15 (brd, 1H), 6.87 (s, 1H); ESI-MS m/z 354 ($\text{M}+\text{H}$) $^+$, 376 ($\text{M}+\text{Na}$) $^+$.

3-Bromo-2-cyano-5-dimethylcarbamoyl-4-hydroxy-3,4-dihydro-2H-pyridine-1-carboxylic acid methyl ester

To a solution of **8b** (58.7 mg, 0.25 mmol, 86% ee) in DMSO (2 ml) and H_2O (1 ml), NBS (44.5 mg, 0.25 mmol) was added at room temperature. After 4.5 h, the reaction mixture was added to water, and the products were extracted with ethyl acetate for 3 times. The organic phase was washed with brine, dried over Na_2SO_4 and concentrated *in vacuo*. NMR analysis of the crude oil indicated that the reaction is completely stereoselective, and none of minor stereoisomers were observed. The resulting residue was purified by silica gel column chromatography (10% methanol in chloroform) to give **R** (43.0 mg, 0.12 mmol, 48%) as a colorless solid. $^1\text{H-NMR}$ (DMSO-d_6 , 50 $^\circ\text{C}$) δ 2.91 (s, 3H), 3.02 (s, 3H), 3.79 (s, 3H), 5.28 (d, $J = 5.5$ Hz, 1H), 5.60 (brd, 1H), 5.97 (brd, 1H), 6.24 (d, $J = 5.5$ Hz, 1H); $^{13}\text{C-NMR}$ (DMSO) δ 34.4, 38.2, 41.0, 44.0, 53.7, 71.0, 116.0, 124.3, 135.8, 155.0, 167.1; $[\alpha]_D^{25} +225.74$ (c 0.95, $\text{CHCl}_3/\text{MeOH} = 1/1$, 87% ee); IR (neat) 3352, 2969, 1716, 1623, 1446, 1330, 1310, 1266, 1203, 1048 cm^{-1} ; ESI-MS 354, 356 ($\text{M}+\text{Na}$) $^+$; MS (EI) m/z 314 ($\text{M}-\text{OH}$) $^+$, 252 ($\text{M}-\text{Br}$) $^+$; HRMS (EI) Calcd for $\text{C}_9\text{H}_8\text{N}_3\text{O}$ ($\text{M}-\text{BrOH}-\text{HCOOMe}$) 174.0662, Found 174.0648; HPLC (DAICEL CHIRALPAK AD-H, hexane/2-propanol = 9/1, flow rate: 1.0 ml/min, detector: 254 nm, t_R (major) = 51.1 min, t_R (minor) = 45.0 min. The stereochemistry of **R** was temporarily assigned based on NOESY and coupling constant.

2-Cyano-5-dimethylcarbamoyl-3-hydroxy-4-methoxy-3,4-dihydro-2H-pyridine-1-carboxylic acid methyl ester

3,3-Dimethyldioxirane in acetone solution was prepared following a general procedure, using oxone (10 g, 16.3 mmol), acetone (6 ml), and aq NaHCO₃ (4.8 g in 8 ml water).⁴ To a solution of 3,3-dimethyldioxirane in acetone at -78 °C, a solution of **8b** (90 mg, 0.436 mmol) in methylene chloride (0.5 ml) was added. The mixture was stirred at -30 °C for 15 h. After the starting material disappeared on TLC, argon gas was bubbled to evaporate 3,3-dimethyldioxirane in an ice bath. Methanol (0.5 ml) was added, and the mixture was stirred in an ice bath for 3 h. The volatiles were evaporated, and the residue oil was purified through silica gel chromatography to give the product **S** in 49% yield (60 mg). NMR analysis of the crude oil indicated that the reaction is completely stereoselective, and none of minor stereoisomers were observed. The stereochemistry of **S** was temporarily assigned based on NOESY. ¹H NMR (C₆D₆) δ 2.14 (brd, 3H), 2.26 (brd, 3H), 2.82 (s, 3H), 3.02 (s, 3H), 3.89 (d, *J* = 8.6 Hz, 1H), 4.07 (brd, 1H), 4.25 (d, *J* = 6.1 Hz, 1H), 5.49 (s, 1H), 5.67 (s, 1H); ¹³C NMR (C₆D₆) δ 34.8, 38.2, 48.9, 53.3, 56.7, 64.4, 82.1, 116.4, 132.0, 155.4, 167.0; [α]_D²⁶ -24.80 (c 0.29, CHCl₃) IR (neat) 3446, 1634 cm⁻¹; ESI-MS *m/z* 306 (M+Na)⁺; HRMS (FAB) Calcd for C₁₁H₁₄O₄N₃ (M-CH₃OH+H)⁺ 252.0984, Found 252.0984.

ESI-MS studies of sulfoxide catalysts (Fig. 1–6):

A solution of ligand **6** (0.02 mmol) and Et₂AlCl (0.98 M in *n*-hexane, 0.02 or 0.01 mmol) in methylene chloride (0.5 ml) was stirred at room temperature for 1 h, and part of the mixture (50 μ l) was diluted with methylene chloride (5 ml). This solution was analyzed by ESI-mass (Fig. 1,2). The reaction mixture was cooled to -60 °C and *N,N*-diisopropylnicotinamide (0.4 M in methylene chloride, 0.5 ml, 0.20 mmol) was added. Part of this mixture (50 μ l) was diluted with methylene chloride (5 ml) and this solution was analyzed by ESI-mass (Fig. 3,4). Fig. 5 and 6 were measured as the same method using the ligand **4** and **5**.

Fig. 3 **Fig. 4** **Fig. 1** **Fig. 5** **Fig. 6**

The ESI-MS studies indicated that the distribution of the components is dependent on the ratio of Et_2AlCl and the ligand when the catalyst was prepared, and on the stereochemistry of the sulfoxide. Based on the comparison of Fig. 1 and 2, and Fig. 3 and 4, a catalyst solution prepared from Et_2AlCl and ligand **6** in a 1:2

ratio contained more amounts of the 2:3 complexes (complex **W** and **X**) of aluminum and **6**. Based on the comparison of Fig. 1, 5, and 6, the amounts of the 2:3 complexes are significantly higher in the case of catalyst prepared from ligand **6** than those from **4** and **5**. Thus, the enantioselectivity of the Reissert reaction seemed to be higher when the amounts of the 2:3 complexes are higher. These results suggested that the active enantioselective catalyst is a 2:3 complex of aluminum and the ligand **6**.

Confirmation of the ability of a sulfoxide and a phosphine sulfide as a Lewis base to activate TMSCN:

	Control	+DMSO	+Bu ₃ PS
7.5 h	1%	14%	9%
10 h	1.6%	18%	11%

For the initial kinetics, DMSO and Bu₃PS accelerated a cyanosilylation of hydrocinnamaldehyde in ca. 14 and 9 times. Using Bu₃PO (40 mol %), the reaction proceeded to 81% yield of the product at room temperature for 7.5 h, which indicated the higher ability of a phosphine oxide for activation of TMSCN as a Lewis base.

Lewis base catalyzed Reissert reaction of **10:**

The Lewis base itself (**Y** and **Z**) can promote the Reissert reaction of **10**, however with completely different regioselectivity (1,2-selective). These results indicated that the bifunctional system is important for the high regio- (1,6-selectivity) and enantioselectivity. 1,2-products obtained using **Y** and **Z** were almost racemic.

¹ Hamashima, Y.; Sawada, D.; Nogami, H.; Kanai, M.; Shibasaki, M. *Tetrahedron* **2001**, 57, 805.

² Meyers, A. G.; Yang, B. H.; Kopecky D. J. *Tetrahedron Lett.*, **1996**, 37, 3623.

³ Urban, F. J. PCT Int. Appl. WO9325552

⁴ Murray, R. W.; Jeyaraman, R. *J. Org. Chem.* **1985**, 50, 2847.