SUPPLEMENTARY DATA 1

Journal of Vertebrate Paleontology

New Middle Triassic tetrapods from the upper Fremouw Formation of Antarctica and their depositional setting

CHRISTIAN A. SIDOR,^{*,1} ROGER M. H. SMITH,² ADAM K. HUTTENLOCKER,¹ and
BRANDON R. PEECOOK¹

¹Department of Biology and Burke Museum, University of Washington, Seattle, Washington
98195, U.S.A., casidor@uw.edu, bpeecook@uw.edu, huttenla@uw.edu;

²Karoo Palaeontology, Iziko: South African Museum, Cape Town 8000, South Africa,
rsmith@iziko.org.za

EXPLANATION

The figures included below correspond to those in the print version of the journal article.
They are included here as additional information.

FIGURE S1. Measured sections at Gordon Valley and Fremouw Peak vertebrate sites.

FIGURE S2. Isolated left squamosal attributed to *Angonisaurus* sp. (UWBM 95538) from the upper Fremouw Formation of Antarctica.

FIGURE S3. Left ilium attributed to *Therocephalia* indet. (UWBM 96187) from the upper Fremouw Formation of Antarctica.

FIGURE S4. Proximal right archosaur femur (UWBM 95548) from the upper Fremouw Formation of Antarctica.