New Role of CO₂ as a Selective Agent in Palladium-Catalyzed Reductive Ullmann Coupling with Zinc in Water

Jin-Heng Li* Ye-Xiang Xie and Du-Lin Yin

*Institute of Fine Catalysis and Synthesis, College of Chemistry and Chemical
Engineering, Hunan Normal University, Changsha 410081, China*

Supporting Information

List of Contents

(A) General	S2
(B) Materials	S2
(C) A typical experimental procedure for the Ullmann coupling	S2
(D) Analytical data for 2	S2-4
(E) References	S4
(F) NMR spectra	S5-26

(A) General:

^1H and ^{13}C NMR spectra were recorded on a Bruker AMX-400 or 300 spectrometers

(B) Materials:

All reagents were directly used as obtained commercially.

(C) Typical Experimental Procedure for the Ullmann Coupling:

Iodobenzene (2 mmol) was added to a mixture of Pd/C (5% w/w, 300 mg), zinc powder (170 mg, 2.6 mmol), and H_2O (5 mL) in HF-25 autoclave. CO_2 was then transferred into the autoclave to 1.0 MPa. The reaction mixture was stirred at room temperature for desired time. After the gas was vented, the residue was extracted with ethyl ether and purified by flash column chromatography to give **2a** (hexane/ethyl acetate).

(D) Analytical data for 2:

Biphenyl (2a)

A white solid; ^1H NMR: δ (400 MHz) 7.59 (d, J = 8.4 Hz, 4H), 7.44 (t, J = 8.0 Hz, 4H), 7.39 (t, J = 7.6 Hz, 2H); ^{13}C NMR (100 MHz) 141.2, 129.9, 128.7, 127.2.

4,4'-Dimethyl-biphenyl (2b)

A white solid; ^1H NMR: δ (400 MHz) 7.46 (d, J = 8.0 Hz, 4H), 7.22 (d, J = 7.6 Hz, 4H), 2.37 (s, 6H); ^{13}C NMR (100 MHz) 138.3, 136.7, 129.4, 126.8, 21.0.

2,2'-Dimethyl-biphenyl (2c)

A colorless oil; ^1H NMR: δ (400 MHz) 7.25–7.20 (m, 6H), 7.09 (d, J = 7.6 Hz, 2H), 2.04 (s, 6H);

^{13}C NMR (100 MHz) 141.6, 135.8, 129.8, 129.3, 127.1, 125.5, 19.8.

4,4'-Dimethoxy-biphenyl (2d)

A white solid; ^1H NMR: δ (400 MHz) 7.46 (d, J = 8.8 Hz, 4H), 7.22 (d, J = 8.8 Hz, 4H), 2.37 (s,

6H); ^{13}C NMR (100 MHz) 138.3, 136.7, 129.4, 126.8, 21.0.

4,4'-Bis-trifluoromethyl-biphenyl (2e)

A white solid; ^1H NMR: δ (300 MHz) 7.75–7.57 (m, 8H); ^{13}C NMR (75 MHz) 158.7, 133.5, 127.7,

114.4, 55.3.

[1,1']Binaphthalenyl (2f)

A white solid; ^1H NMR: δ (400 MHz) 7.86–7.83 (m, 7H), 7.49–7.46 (m, 7H); ^{13}C NMR (100

MHz) 133.4, 127.9, 125.8, 125.4.

 $N^4, N^4, N^{4'}, N^{4'}$ -Tetramethyl-biphenyl-4,4'-diamine (2i)

A white solid; ^1H NMR: δ (300 MHz) 7.05 (d, J = 8.0 Hz, 4H), 6.63 (d, J = 8.0 Hz, 4H), 2.94 (s,

12H); ^{13}C NMR (75 MHz) 129.7, 113.4, 41.3, 40.2.

2,6,2',6'-Tetramethoxy-biphenyl (2j)

A white solid; ^1H NMR: δ (300 MHz) 6.93–6.83 (m, 6H), 3.79 (s, 6H), 3.72 (s, 6H); ^{13}C NMR (75

MHz) 153.7, 151.7, 129.0, 117.5, 113.8, 112.9, 56.9, 56.1.

[1,1']Binaphthalenyl-2,2'-diol (2k)

A white solid; ^1H NMR: δ (300 MHz) 7.98 (d, J = 9.0 Hz, 2H), 7.89 (d, J = 8.7 Hz, 2H),

7.40–7.26 (m, 6H), 7.15 (d, $J = 6.6$ Hz, 2H), 1.54 (s, 2H); ^{13}C NMR (75 MHz) 153.1, 133.7, 131.8, 129.8, 128.8, 124.5, 124.4, 118.1, 111.2.

3,5,3',5'-Tetramethoxy-biphenyl (2m)

A white solid; ^1H NMR: δ (300 MHz) 6.71 (s, 4H), 6.47 (s, 2H), 5.84 (s, 12H); ^{13}C NMR (75 MHz) 161.3, 106.5, 105.9, 99.8, 55.8, 55.6.

Biphenyl-3,3'-dicarboxylic acid dimethyl ester (2o)

A white solid; ^1H NMR: δ (300 MHz) 8.30 (s, 2H), 8.05 (d, $J = 7.8$ Hz, 2H), 7.82 (d, $J = 6.9$ Hz, 2H), 7.56 (t, $J = 8.0$ Hz, 2H), 4.46–4.39 (m, 4H), 1.43 (t, $J = 7.2$ Hz, 6H); ^{13}C NMR (75 MHz) 166.8, 140.8, 131.8, 131.5, 129.3, 129.1, 128.6, 61.5, 14.7.

(E) References:

- (1) Kuroboshi, M.; Waki, Y.; Tanaka, H. *J. Org. Chem.* **2003**, *68*, 3938 and references cited therein.
- (2) Mukhopadhyay, S.; Rothenberg, G.; Wiener, H.; Sasson, Y. *Tetrahedron* **1999**, *55*, 14763

(F) NMR spectra:

Biphenyl (2a)

Biphenyl (2a)

2,2'-Dimethyl-biphenyl (2c)

2,2'-Dimethyl-biphenyl (2c)

4,4'-Dimethoxy-biphenyl (2d)

[1,1']Binaphthalenyl (2f)

[1,1']Binaphthalenyl (2f)

N,N',N',N'-Tetramethyl-biphenyl-4,4'-diamine (2i)

2,6,2',6'-Tetramethoxy-biphenyl (2j)

3,5,3',5'-Tetramethoxy-biphenyl (2m)

Biphenyl-3,3'-dicarboxylic acid dimethyl ester (2o)