Supporting Information

for communication entitled:

“New Structures from Multiple Rearrangements of Propargylic Dialkoxy Disulfides”

by

Samuel Braverman,* Tatiana Pechenick, Hugo E. Gottlieb and Milon Sprecher

Department of Chemistry, Bar-Ilan University, Ramat-Gan 52900, Israel

*Corresponding author: Tel. 972-3-5318322; Fax: 972-3-5351250; e-mail: bravers@mail.biu.ac.il

Contents:

- 1. General procedure for preparation of 10/11 and 14/15.**
- 2. Full spectral data for 10a/11a; 10b/11b; 10c; 10d/11d.**
- 3. Full spectral data for 14f/15f; 14g/15g.**

1. General procedure for the preparation of 10/11 and 14/15.

A chloroform solution of the appropriate propargyloxy disulfide (**1a-g**) was heated under reflux for the time indicated in the text. After cooling to room temperature the solvent was evaporated under reduced pressure and the residue was purified by silica gel chromatography. For compounds **10a/11a** and **10b/11b** CHCl₃ was used as eluent. Compound **10c** was isolated by using first: hex/CHCl₃ 1:2 followed by CHCl₃. **10d/11d** were purified by using an eluent with gradient of polarities as follows: hex/CHCl₃ 4:1→2:1→1:1→1:2→100% CHCl₃. **14f/15f** were isolated and separated by using a mixture of hex/EtAc 4:1 as eluent. **14g/15g** were purified and separated by using an eluent with gradient of polarities as follows: hex/EtAc 100:1→95:5→8:1→4:1→1:1.

2. Full spectral data for 10a/11a; 10b/11b; 10c; 10d/11d.

NMR spectra were taken at 298 K in CDCl₃ with TMS as internal reference and run at 600 (¹H) and 75 (¹³C) MHz. Table 1 and 2 were unambiguously determined with the aid of two-dimensional techniques such as COSY, HMQC and HMBC.

Table 1. ^1H NMR chemical shifts

H	10a	11a	10b	11b	10c	10d	11d
1	7.30	7.10					
3	2.85	2.81	2.53	2.77	2.91	2.83	2.86
4	2.61	2.59	2.48	2.45	2.44	2.51	2.54
6	2.09	2.14					
α			2.70	2.43		0.34	0.46
β			1.24	1.27	1.39		
α'			2.16 ^a	2.16 ^a		0.15	0.15 ^b
β'			1.11	1.12	1.20		

^a ca. 2.16, ^b ca.0.15

Table 2. ¹³C NMR chemical shifts

C	10a	11a	10b	11b	10c	10d
1	125.92	124.22	153.42	147.57 ^a	154.28	156.89
2	140.03	149.43	143.01	147.32 ^a	146.89	162.93
3	25.79	31.79	26.94	28.20	29.08	31.23
4	18.50	18.22	17.48	17.76	18.21	19.48
5	81.40	80.64	76.77	76.23	75.19	102.64
6	70.93	71.55	83.77	85.14	92.17	88.63
α			22.33	20.68	37.84	-0.75
β			13.02	12.85	29.58	
α'			12.37	12.37	27.39	-0.17
β'			14.33	14.03	30.98	

^a values with this superscript may be interchanged

IR (neat), MS and HRMS (elemental composition) data:**10a: 4-but-3-ynyl-1,2-dithiete 1,1-dioxide and 11a: 3-but-3-ynyl-1,2-dithiete 1,1-**

dioxide: IR (neat): 1130, 1319, 2256 cm⁻¹, MS (CI/CH₄): *m/z* 174 (M⁺, 11.68%), 110 (MH⁺-“SO₂”, 21.93%), HRMS (elemental composition): calc. (C₆H₆O₂S₂) 173.980923; found 173.978803.

10b: 3-ethyl-4-hex-3-ynyl-1,2-dithiete 1,1-dioxide and 11b: 4-ethyl-3-hex-3-ynyl-1,2-dithiete 1,1-dioxide: IR (neat): 1153, 1323, 2368, 2975 cm^{-1} , MS (CI/ CH_4): m/z 231 (MH^+ , 65.98%), 167 ($\text{MH}^+ - \text{SO}_2$, 91.91%), 166 ($\text{M}^+ - \text{SO}_2$, 99.32%), 151 (100%), HRMS (elemental composition): calc. ($\text{C}_{10}\text{H}_{15}\text{O}_2\text{S}_2$) 231.051348; found 231.055073.

10c: 3-tert-butyl-4-(5,5-dimethylhex-3-ynyl)-1,2-dithiete 1,1-dioxide: IR (neat): 1155, 1323, 1611, 1646, 2239, 2969 cm^{-1} , MS (CI/ CH_4): m/z 287 (MH^+ , 9.31%), 222 ($\text{M}^+ - \text{SO}_2$, 59.28%), 207 (100%), HRMS (elemental composition): calc. ($\text{C}_{14}\text{H}_{23}\text{O}_2\text{S}_2$) 287.113949; found 287.115349.

10d: 3-trimethylsilanyl-4-(4-trimethylsilanyl-but-3-ynyl)-1,2-dithiete 1,1-dioxide: IR (neat): 1155, 1325, 2179, 2961 cm^{-1} , MS (EI): m/z 319 (MH^+ , 0.92%), 254 ($\text{M}^+ - \text{SO}_2$, 100%), HRMS (elemental composition): calc. ($\text{C}_{12}\text{H}_{23}\text{O}_2\text{Si}_2\text{S}_2$) 319.067806; found 319.068886.

3. Full spectral data for 14f/15f; 14g/15g.

NMR spectra were taken at 298 K in CDCl_3 with TMS as internal reference and run at 600 (^1H) and 75 (^{13}C) MHz. Table 1 and 2 were unambiguously determined with the aid of two-dimensional techniques such as NOESY, COSY, HMQC and HMBC. As yet, it has not been established which of the isomers (**14** or **15**) has the $\text{S}=\text{O}$ *endo* configuration and which the *exo*.

14f/15f: 3,6-Bis-(2,2-dimethyl-propylidene)-2-oxa-5,7-dithia-bicyclo[2.2.1]heptane 5-oxide (overall yield for two isomers endo+exo 57%):

for 14f: ^1H NMR (300 MHz, CDCl_3): δ 6.76 (br. t, $J = 0.8$ Hz, 1H (CH–O)), 6.23 (br. d, $J = 0.6$ Hz, 1H (=CH of “SO” side)), 5.77 (br. d, $J = 0.8$ Hz, 1H (CH–SO)), 5.26 (s, 1H (=CH of “O” side)), 1.25 (s, 9H ((CH_3)₃ of “SO” side)), 1.20 (s, 9H ((CH_3)₃ of “O” side)), ^{13}C NMR (75 MHz, CDCl_3): δ 149.60 (=C–SO), 147.04 (=CH– of “SO” side), 140.57 (=C–O), 119.22 (=CH– of “O” side), 81.03 (–CH–O), 69.18 (–CH–SO), 35.14 (–C(CH_3)₃ of “SO” side), 31.41 (–C(CH_3)₃ of “O” side), 31.41 (–C(CH_3)₃ of “O” side), 30.45 (–C(CH_3)₃ of “SO” side), IR (neat): 1062 (S=O), 1365, 1465, 1669, 2237, 2961 cm^{-1} , MS (CI/ CH_4): m/z 286 (M^+ , 71.77%), 238 (M^+ –“SO”, 20.45%), 175 (77.36%), HRMS (elemental composition): calc. ($\text{C}_{14}\text{H}_{22}\text{O}_2\text{S}_2$) 286.106124; found 286.103421.

for 15f: ^1H NMR (300 MHz, CDCl_3): δ 6.58 (d, $J = 0.8$ Hz, 1H (CH–O)), 5.91 (br. s, 1H (=CH of “SO” side)), 5.83 (t, $J = 0.8$ Hz, 1H (CH–SO)), 5.56 (br. s, 1H (=CH of “O” side)), 1.21 (s, 9H ((CH_3)₃ of “SO” side)), 1.19 (s, 9H ((CH_3)₃ of “O” side)), ^{13}C NMR (75 MHz, CDCl_3): δ 143.75 (=C–SO), 142.09 (=CH– of “SO” side), 140.86 (=C–O), 120.29 (=CH– of “O” side), 82.64 (–CH–O), 63.84 (–CH–SO), 35.03 (–C(CH_3)₃ of “SO” side), 31.95 (–C(CH_3)₃ of “O” side), 31.85 (–C(CH_3)₃ of “O” side), 30.57 (–C(CH_3)₃ of “SO” side), IR (neat): 1082 (S=O), 1365, 1464, 1673, 2240, 2960 cm^{-1} , MS (CI/ CH_4): m/z 286 (M^+ , 50.86%), 239 (MH^+ –“SO”, 66.41%), 209 (100%), HRMS (elemental composition): calc. ($\text{C}_{14}\text{H}_{22}\text{O}_2\text{S}_2$) 286.106124; found 286.108552.

14g/15g: 3,6-Bis-(1-adamantyl-methylidene)-2-oxa-5,7-dithia-bicyclo[2.2.1]heptane 5-oxide (overall yield for two isomers endo+exo 62%):

for 14g: ^1H NMR (600 MHz, CDCl_3): δ 6.78 (s, 1H (CH–O)), 5.94 (s, 1H (=CH of “SO” side)), 5.79 (s, 1H (CH–SO)), 5.04 (s, 1H (=CH of “O” side)), 2.03 (m, 3H (CH₂–CH–CH₂ of “SO” side)), 2.00 (m, 3H (CH₂–CH–CH₂ of “O” side)), 1.81 (m, 6H (–C–CH₂–CH of “SO” side)), 1.75 (m, 6H (–C–CH₂–CH of “O” side)), 1.73 (m, 6H (CH–CH₂–CH of “O” side)), 1.67 (m, 6H (CH–CH₂–CH of “SO” side)), ^{13}C NMR (75 MHz, CDCl_3): δ 149.43 (=C–SO), 146.69 (=CH– of “SO” side), 140.45 (=C–O), 119.52 (=CH– of “O” side), 81.91 (–CH–O), 69.49 (–CH–SO), 43.46 (–C–CH₂–CH of “O” side), 42.15 (–C–CH₂–CH of “SO” side), 38.32 (–C– of “SO” side), 36.43 (CH–CH₂–CH of “O” side), 36.21 (CH–CH₂–CH of “SO” side), 33.66 (–C– of “O” side), 28.40 (CH₂–CH–CH₂ of “O” side), 28.09 (CH₂–CH–CH₂ of “SO” side), IR (neat): 1053 (S=O), 1446, 2231, 2910 cm^{-1} , MS (CI/i-Bu): m/z 442 (M^+ , 3.10%), 135 (22.08%), HRMS (elemental composition): calc. ($\text{C}_{26}\text{H}_{34}\text{O}_2\text{S}_2$) 442.200024; found 442.203926.

for 15g: ^1H NMR (600 MHz, CDCl_3): δ 6.61 (s, 1H (CH–O)), 5.84 (s, 1H (CH–SO)), 5.66 (s, 1H (=CH of “SO” side)), 5.36 (s, 1H (=CH of “O” side)), 2.02 (m, 3H (CH₂–CH–CH₂ of “O” side)), 1.99 (m, 3H (CH₂–CH–CH₂ of “SO” side)), 1.79 (m, 6H (–C–CH₂–CH of “SO” side)), 1.74 (m, 6H (CH–CH₂–CH of “SO” side)) + m, 6H (–C–CH₂–CH of “O” side)), 1.68 (m, 6H (CH–CH₂–CH of “O” side)), ^{13}C NMR (75 MHz, CDCl_3): δ 143.54 (=C–SO), 141.78 (=CH– of “SO” side), 140.64 (=C–O),

120.76 (=CH- of "O" side), 82.87 (-CH-O), 64.25 (-CH-SO), 43.96 (-C-CH₂-CH of "O" side), 42.30 (-C-CH₂-CH of "SO" side), 37.53 (-C- of "SO" side), 36.51 (CH-CH₂-CH of "SO" side), 36.30 (CH-CH₂-CH of "O" side), 33.95 (-C- of "O" side), 28.39 (CH₂-CH-CH₂ of "SO" side), 27.17 (CH₂-CH-CH₂ of "O" side), IR (neat): 1079 (S=O), 1450, 2903 cm⁻¹, MS (CI/i-Bu): *m/z* 442 (M⁺, 10.70%), 135 (86.34%), HRMS (elemental composition): calc. (C₂₆H₃₄O₂S₂) 442.200024; found 442.203509.

ppm

¹H NMR chemical shifts

Current Data Parameters
NAME 79-2-95
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 500000
Time 20.48
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 8
DS 0
SWH 4194.631 Hz
FIDRES 0.128010 Hz
AQ 3.905956 sec
RG 256
DM 119.200 usec
DE 4.50 usec
TE 300.0 K
D1 0.00100000 sec
P1 8.50 usec
DE 4.50 usec
SF01 300.1319000 MHz
NUC1 1H
PL1 -3.00 dB

F2 - Processing parameters
SI 32768
SF 300.1300043 MHz
WDW EM
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 22.50 cm
F1P 10.750 ppm
F1 3226.40 Hz
F2P -0.500 ppm
F2 -150.07 Hz
PRGCM 0 50000 ppm/cm
HZCM 150.06500 Hz/cm

ppm

¹H NMR chemical shifts

2.716
2.691
2.533
2.515
2.478
2.470
2.447
2.443
2.418
2.176
2.150
2.146
1.291
1.266
1.264
1.239
1.214
1.144
1.140
1.119
1.115
1.094
1.090

22.820
40.646
20.296

30.912
30.000

10
9
8
7
6
5
4
3
2
1
0

Current Data Parameters
NAME 104-s11c
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20000404
Time 22.17

INSTRUM 5 mm Multinu
PROBHD zg
PULPROG 32786
TD 32786
SOLVENT CDCl3
NS 8
DS 0

SWH 4194.631 Hz
FIDRES 0.128010 Hz
AQ 3.905956 sec
RG 36.9
DE 119.200 usec
TE 300.0 K

D1 0.00100000 sec
P1 8.50 usec
DE 4.50 usec
SF01 300.1315000 MHz
NUC1 1H
PL1 -3.00 dB

F2 - Processing parameters
SI 32766
SF 300.1299825 MHz
MDN EN
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

10 NMR plot parameters
CX 22.50 cm
F1P 10.750 ppm
F1 3226.40 Hz
F2P -0.500 ppm
F2 -150.07 Hz
FREQH 0.50000 ppm/cm
HZCM 150.06500 Hz/cm

ppm

153.42
147.57
147.32
143.01

85.14
83.77
77.59
77.16
76.77
76.74
76.23

28.20
26.93
26.24
22.33
20.68
18.26
17.89
17.76
14.33
14.03
13.81
13.02
12.85
12.41
12.37

¹³C NMR chemical shifts

Current Data Parameters
NAME 104a11.a
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20060404
Time 22.32
INSTRUM spect
PROBHD 5 mm AULI11V
PULPROG zgpgc
TD 32768
SOLVENT CDCl3
NS 1362
DS 0
SWH 19723.865 Hz
FIDRES 0.801525 Hz
AQ 0.8307186 sec
RG 5792.6
DM 25.350 usec
DE 9.46 usec
TE 300.0 K
D11 0.1000000 sec
R12 10.00 dB
CPRPG2 waltz16
PCPR2 100.00 usec
PCPR2 300.1312000 MHz
NUC2 1H
PL2 -8.00 dB
D1 0.1000000 sec
P1 3.00 usec
DE 9.46 usec
SF01 75.4759867 MHz
NUC1 13C
PL1 -5.00 dB

F2 - Processing parameters
SI 32768
SF 75.4677459 MHz
WDW EM
SSB 0
LB 2.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 22.50 cm
F1P 220.000 ppm
F1 16502.90 Hz
F2P -377.34 Hz
F2 10.00000 ppm/cm
H2CH 754.67745 Hz/cm

ppm

154.28

146.89

92.17

77.57
77.15
76.73
75.1937.83
33.09
31.12
30.98
30.77
30.67
30.50
30.18
29.58
29.08
27.39
18.99
18.21¹³C NMR chemical shifts

Current Data Parameters
NAME 142-811c
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20000804
Time 18.03
INSTRUM spect
PROBHD 5 mm BBL1nu
PULPROG zgpg
TD 32768
SOLVENT CDCl₃
NS 408
DS 0
SWH 19723.885 Hz
FIDRES 0.801585 Hz
AQ 0.6307188 sec
RG 9782.6
DB 25.350 usec
DE 9.46 usec
TE 300.0 K
D1 0.10000000 sec
PL12 10.00 dB
CHRG22 waltz16
PCPD2 100.00 usec
SFO2 300.1312000 MHz
NUC2 1H
PL2 -9.00 dB
D1 0.10000000 sec
P1 3.00 usec
DE 9.46 usec
SFO1 75.4756807 MHz
NUC1 13C
PL1 -9.00 dB

F2 - Processing Parameters
SI 32768
SF 75.4877488 MHz
WDW EM
SSB 0
LB 2.00 Hz
GB 0
PC 1.00

10 NMR plot parameters
CX 22.50 cm
F1P 220.000 ppm
F1 16002.80 Hz
F2P -9.000 ppm
F2 -977.34 Hz
FREQM 10.00000 ppm/cm
HZCM 754.87748 Hz/cm

Current Data Parameters
NAME 194-15
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20010120
Time 19.41
INSTRUM spect
PROBHD 5 mm N/11nu
PULPROG zgpg
TD 32768
SOLVENT CDCl3
NS 3130
DS 0
SIH 19723.865 Hz
FIDRES 0.601563 Hz
AQ 0.6307186 sec
RG 5782.0
DM 25.350 usec
DE 9.46 usec
TE 300.0 K
D11 0.10000000 sec
PL12 10.00 dB
CHPRG2 waltz16
PCPD2 100.00 usec
SFO2 300.1312000 MHz
NUC2 1H
PL2 -6.00 dB
D1 0.10000000 sec
P1 3.00 usec
DE 9.46 usec
SFO1 75.475967 MHz
NUC1 13C
PL1 -5.00 dB

F2 - Processing parameters
SI 32768
SF 75.467467 MHz
WDW EM
SSB 0
LB 2.00 Hz
GB 0
PC 1.00

1D FID plot parameters
CX 22.50 cm
FAP 220.000 ppm
F1 16602.90 Hz
F2 -5.000 ppm
F2 -377.34 Hz
PCPDH 10.00000 ppm/cm
H2DM 754.67748 Hz/cm

ppm

7.27
6.76
6.75
6.75
6.23
6.23
5.77
5.77
5.26

1.28
1.27
1.27
1.26
1.26
1.25
1.24
1.24
1.20
1.19

0.00

¹H NMR chemical shifts

Integral

10.000
10.023
10.056
10.045

90.761
90.127

10
9
8
7
6
5
4
3
2
1
0

Current Data Parameters
NAME 178-2a11
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20001106

Time 19.44
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl₃

NS 0
DS 0
SWH 4194.631 Hz
FIDRES 0.128010 Hz
AQ 3.9058956 sec

RG 90.5
DM 119.200 usec
DE 4.50 usec
TE 300.0 K
D1 0.00100000 sec
P1 8.50 usec
DE 4.50 usec
SF01 300.1315000 MHz
NUC1 ¹H
PL1 -3.00 dB

F2 - Processing parameters
SI 32768
SF 300.1300023 MHz
WDW EM
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

10 peak list parameters
CX 22.50 cm
FIP 10.750 ppm
F1 3226.40 Hz
F2P -0.500 ppm
F2 -150.07 Hz
PPMACH 0.50000 ppm/cm
HZCM 150.06500 Hz/cm

Current Data Parameters
NAME 178-n1
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20001114
Time 5.43
INSTRUM spect
PROBHD 5 mm Naltru
PULPROG zgpgc
TD 32768
SOLVENT CDCl3
NS 38000
DS 0
SWH 19723.868 Hz
FIDRES 0.601565 Hz
AQ 0.6307188 sec
RG 5792.6
DB 25.350 usec
DE 9.46 usec
TE 300.0 K
D11 0.10000000 sec
PL12 10.00 dB
CHPRG2 waltz16
PCPD2 100.00 usec
SFO2 300.1312000 MHz
NUC2 1H
PL2 -8.00 dB
D1 0.10000000 sec
P1 3.00 usec
DE 9.46 usec
SFO1 75.4759867 MHz
NUC1 13C
PL1 -5.00 dB

F2 - Processing parameters
SI 32768
SF 75.4677461 MHz
WDW EM
SSB 0
LB 2.00 Hz
GB 0
PC 1.00

10 MHz plot parameters
CX 22.50 cm
FAP 220.00 ppm
F1 16802.80 Hz
F2 -5.000 ppm
F2 -377.34 Hz
FREQCH 10.00000 ppm/cm
HZCH 754.67746 Hz/cm

ppm

7.260
6.774
5.968
5.783
5.042

2.040
2.034
2.007
1.822
1.813
1.762
1.752
1.709
1.705
1.662
1.651

¹H NMR chemical shifts

14g R=adamantyl-

Current Data Parameters
NAME 367-811
EXPNO 5
PROCNO 1

F2 - Acquisition Parameters
Date_ 20021016
Time 15.58
INSTRUM spect
PROBHD 5 mm Naltru
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 16
DS 0
SH1 4194.631 Hz
FIDRES 0.128010 Hz
AQ 3.9059556 sec
RG 181
DM 119.200 usec
DE 4.50 usec
TE 300.0 K
D1 0.00100000 sec
P1 8.50 usec
DE 4.50 usec
SF01 300.1318000 MHz
NUC1 1H
RL1 -3.00 dB

F2 - Processing parameters
SI 32768
SF 300.130063 MHz
MCM EM
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 22.50 cm
F1P 10.750 ppm
F1 3226.40 Hz
F2P -0.500 ppm
F2 -150.07 Hz
PPH4M 0.50000 ppm/cm
HZCM 150.06500 Hz/cm

ppm

¹³C NMR chemical shifts

Current Data Parameters
NAME 367-s114
EXPNO 8
PROCNO 1

F2 - Acquisition Parameters

Date_ 20021018
Time 8.18
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 38504
DS 0
SWH 19723.865 Hz
FIDRES 0.601925 Hz
AQ 0.6307188 sec
RG 5792.6
DM 25.350 usac
DE 9.46 usac
TE 300.0 K
D11 0.1000000 sec
PL12 10.00 dB
CPOPRG2 waltz16
PCPD2 100.00 usac
SF02 300.1312000 MHz
NUC2 1H
PL2 -8.00 dB
D1 0.1000000 sec
P1 3.00 usac
DE 9.46 usac
SF01 75.475867 MHz
NUC1 13C
PL1 -5.00 dB

F2 - Processing parameters

SI 32768
SF 75.467511 MHz
WDW EM
SSB 0
LB 2.00 Hz
GB 0
PC 1.00

1D NMR plot parameters

CH 22.50 cm
F1P 220.000 ppm
F1 16602.50 Hz
F2P -377.34 Hz
F2 10.00000 ppm/cm
HZCM 754.67748 Hz/cm