A new three-step domino reaction “thiophilic addition – β -elimination of fluoride – [3,3] sigmatropic rearrangement”: Synthesis of α -allylic and α,α -bis(allylic) α -trifluoromethyl dithioesters.

Fabienne Grellepois, Vadim M. Timoshenko, Yuriy G. Shermolovich, Charles Portella.*

General experimental:

THF and monoglyme (dimethoxyethane, DME) were distilled from sodium-benzophenone and all reactions were carried out with dry, freshly distilled solvents, under anhydrous conditions (argon atmosphere).

All Grignard reagents were purchased from Aldrich Chemical Company and were titrated through the combined use of (+)-menthol and (1,10)-phenanthroline as indicator.¹

Pentafluoroethyl(trimethyl)silane was given by Bayer Chemicals. It can be ordered (ABCR, Fluorochem) or prepared according reported procedure.²

¹H, ¹³C and ¹⁹F NMR spectra were recorded on a Bruker AC-250 spectrometer in CDCl₃. Chemical shift are reported in ppm related to Me₄Si (for ¹H and ¹³C NMR) and CFCl₃ (for ¹⁹F NMR) as internal standards. In the ¹³C NMR data, reported signal multiplicities are related to C-F coupling. The following abbreviations are used to indicate the multiplicities: s, singlet; bs, broad singlet; d, doublet; t, triplet; q, quartet; quint, quintet; sext, sextet; m, multiplet.

All reactions were monitored by ¹⁹F NMR Spectrometry.

Silicagel Merck 9385 (40-63 μ m) was used for flash chromatography. Thin-layer chromatography using precoated aluminium backed plates (Merck Kieselgel 60 F254) were visualised by UV light and/or by a solution of KMnO₄.

¹ For the titration of Grignard reagents, see: Lin, H.-S.; Paquette, L. A. *Synth. Comm.* **1994**, 24 (17), 2503-2506.

² (a) Petrov, V.A. *Tetrahedron Lett.* **2001**, 42, 3267-3269. (b) Marhold, A.; Wiedermann, J.; Bach, S. *Eur. Pat. Appl.* 2002, EP2001-114733; US2001-895774; CAN 136:85940.

- Preparation and characterization of pentafluoro dithioesters **1a-b**. S3-S4
- Procedure A: reaction of Grignard reagents with pentafluoro dithioesters **1a-b** for the preparation of unsaturated dithioesters **3-7**. S4
- Preparations and characterizations of monounsaturated dithioesters 3-7.* S4-S7
- Procedure B: reaction of Grignard reagents with tetrafluoro monounsaturated dithioesters **3-5** or **7** for the preparation of trifluoromethyl bis(unsaturated) dithioesters **9-14**. S7
- Procedure C: “one-pot” preparation of trifluoromethyl bis(unsaturated) dithioesters **9,12** starting from pentafluoro dithioesters **1a-b**. S7
- Preparations and characterizations of bis(unsaturated) dithioesters 9-14.* S7-S11
- ^{19}F , ^1H and ^{13}C spectra of all compounds. S12-S50

General procedure for the preparation of pentafluoro dithioesters 1a-b:³ To dried tetramethyl ammonium fluoride (TMAF)⁴ (1 equiv) was added monoglyme at 0°C and under Ar. The solution was cooled to -10 °C for the addition of carbon disulfide (1 equiv) and then at -60 °C for the addition of pentafluoroethyl(trimethyl)silane (1.05 equiv). The reaction mixture was stirred at -40 °C for 1 h. At this temperature, tetramethylammonium iodide (0.1 equiv) and *n*-butyl bromide or allyl bromide (1.05 equiv) were added. The reaction was slowly warmed to room temperature. The reaction was monitored by ¹⁹F NMR until completion. The precipitate was then filtered and the filtrate was diluted with diethyl ether and washed several time with brine. The organic layer was dried (MgSO₄) and concentrated under reduced pressure to give pentafluoro dithioester **1** as a deep red liquid.

Butyl pentafluorodithiopropionate (1a). According to the general procedure, the reaction of TMAF (2.54 g, 27.3 mmol, 1 equiv), CS₂ (1.64 mL, 27.3 mmol, 1 equiv), CF₃CF₂SiMe₃ (5.51 g, 28.7 mmol, 1.05 equiv), TMAI (579 mg, 2.87 mmol, 0.1 equiv) and *n*-butyl bromide (3.08 mL, 28.7 mmol, 1.05 equiv) in monoglyme (40 mL) yielded butyl pentafluorodithiopropionate **1a** (4.96 g, 72%) as a red liquid; ¹⁹F NMR: δ -82.0 (t, ³J_{F-F} = 2.0 Hz, 3F, CF₃), -107.2 (q, ³J_{F-F} = 2.0 Hz, 2F, CF₂); ¹H NMR: δ 0.90 (t, ³J_{H-H} = 7.0 Hz, 3H, CH₃), 1.41 (sext, ³J_{H-H} = 7.0 Hz, 2H, CH₂-CH₃), 1.64 (quint, ³J_{H-H} = 7.0 Hz, 2H, SCH₂-CH₂), 3.18 (t, ³J_{H-H} = 7.0 Hz, 2H, SCH₂); ¹³C NMR: δ 11.6 (CH₃), 20.5 (CH₂), 26.7 (CH₂), 34.6 (t, ⁴J_{C-F} = 2.0 Hz, SCH₂), 109.1 (tq, ¹J_{C-F} = 261.0 Hz, ²J_{C-F} = 37.5 Hz, CF₂), 115.9 (qt, ¹J_{C-F} = 288.0 Hz, ²J_{C-F} = 37.0 Hz, CF₃), 210.8 (tq, ²J_{C-F} = 24.5 Hz, ³J_{C-F} = 0.5 Hz, CS₂).

Allyl pentafluorodithiopropionate (1b). According to the general procedure, the reaction of TMAF (2.32 g, 24.9 mmol, 1 equiv), CS₂ (1.50 mL, 24.9 mmol, 1 equiv), CF₃CF₂SiMe₃ (5.03 g, 26.2 mmol, 1.05 equiv), TMAI (500 mg, 2.49 mmol, 0.1 equiv) and allyl bromide (2.28 mL, 26.2 mmol, 1.05 equiv) in monoglyme (45 mL) yielded allyl pentafluorodithiopropionate **1b** (5.28 g, 84%) as a red liquid; ¹⁹F NMR: δ -81.9 (t, ³J_{F-F} = 2.0 Hz, 3F, CF₃), -107.2 (q, ³J_{F-F} = 2.0 Hz, 2F, CF₂); ¹H NMR: δ 3.94 (d, ³J_{H-H} = 7.0 Hz, 2H, SCH₂), 5.30 (dt, ³J_{H-H} = 10.0 Hz, ⁴J_{H-H} = 1.0 Hz, 1H,

³ Modified procedure from: Badadzhanova, L.A.; Kirij, N.V.; Yagupolskii, Yu. L. *J. Fluorine Chem.* **2004**, *125*, 1095-1098.

⁴ For drying TMAF, 4H₂O, see: Christe, K.O.; Wilson, W.W.; Wilson, R.D.; Bau, R. Feng, J. *J. Am. Chem. Soc.* **1990**, *112*, 7619-7625.

CH=CHH), 5.42 (dq, $^3J_{\text{H-H}} = 17 \text{ Hz}$, $^4J_{\text{H-H}} = 1.0 \text{ Hz}$, 1H, CH=CHH), 5.83 (tdd, $^3J_{\text{H-H}} = 17.0 \text{ Hz}$, $^3J_{\text{H-H}} = 10.0 \text{ Hz}$, $^3J_{\text{H-H}} = 7.0 \text{ Hz}$, 1H, CH=CH₂); ^{13}C NMR: δ 39.5 (t, $^4J_{\text{C-F}} = 2.0 \text{ Hz}$, SCH₂), 111.1 (tq, $^1J_{\text{C-F}} = 261.5 \text{ Hz}$, $^2J_{\text{C-F}} = 37.5 \text{ Hz}$, CF₂), 118.3 (qt, $^1J_{\text{C-F}} = 288.0 \text{ Hz}$, $^2J_{\text{C-F}} = 37.0 \text{ Hz}$, CF₃), 121.6 (CH=CH₂), 128.7 (CH=CH₂), 211.9 (tq, $^2J_{\text{C-F}} = 24.5 \text{ Hz}$, $^3J_{\text{C-F}} = 0.5 \text{ Hz}$, CS₂).

Reaction of Grignard reagents with pentafluoro dithioesters 1a-b for the preparation of monounsaturated dithioesters 3-7 (Procedure A). To a solution of pentafluorodithioester **1a-b** (1 equiv) in THF was added, at -78 °C and under Ar, Grignard reagent (1 equiv).¹ The reaction was then slowly warmed to room temperature and after 15 h of stirring the completion of the reaction was checked by ^{19}F NMR. The reaction mixture was diluted with diethyl ether, washed with sat. aq. sol. NH₄Cl and brine. The organic layer was dried (MgSO₄) and concentrated under reduced pressure. Chromatography on silica gel (pentane) of the residue gave unsaturated dithioesters **3-7**.⁵

Butyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate (3).

According to procedure A, reaction of allyl magnesium bromide with butyl pentafluorodithiopropanoate **1a** (633 mg, 2.51 mmol) in THF (10 mL) yielded butyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **3** (523 mg, 76%) as an orange liquid.

According to procedure A, reaction of *n*-butyl magnesium chloride with allyl pentafluorodithiopropanoate **1b** (700 mg, 2.77 mmol) in THF (10 mL) yielded butyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **3** (562 mg, 74%) as an orange liquid.

^{19}F NMR: δ -78.0 (s, 3F, CF₃), -161.7 (ddq, $^3J_{\text{F-F}} = 6.5 \text{ Hz}$, $^3J_{\text{F-H}} = 12.5 \text{ Hz}$, $^3J_{\text{F-H}} = 35.5 \text{ Hz}$, 1F, CF); ^1H NMR: δ 0.95 (t, $^3J_{\text{H-H}} = 7.0 \text{ Hz}$, 3H, CH₃), 1.45 (sext, $^3J_{\text{H-H}} = 7.0 \text{ Hz}$, 2H, CH₂-CH₃), 1.68 (quint, $^3J_{\text{H-H}} = 7.0 \text{ Hz}$, 2H, CH₂-CH₂-CH₃), 2.96 (ddd, $^2J_{\text{AB}} = 14.5 \text{ Hz}$, $^3J_{\text{H-F}} = 12.5 \text{ Hz}$, $^3J_{\text{H-H}} = 6.5 \text{ Hz}$, 1H, CH_AH_B-CF), 3.21 (t, $^3J_{\text{H-H}} = 7.0 \text{ Hz}$, 2H, SCH₂), 3.28 (dddt, $^2J_{\text{AB}} = 14.5 \text{ Hz}$, $^3J_{\text{H-F}} = 35.5 \text{ Hz}$, $^3J_{\text{H-H}} = 7.0 \text{ Hz}$, $^4J_{\text{H-F}} = 1.0 \text{ Hz}$, 1H, CH_AH_B-CF), 5.18 (m, 2H, CH=CH₂), 5.67 (ddt, $^3J_{\text{H-H}} = 17.0 \text{ Hz}$, $^3J_{\text{H-H}} = 10.0 \text{ Hz}$, $^3J_{\text{H-H}} = 7.0 \text{ Hz}$, 1H, CH=CH₂); ^{13}C

⁵ This reaction sequence resembles a titration of dithioesters **1a-b** with Grignard reagents: the red solution of starting dithioester turned pale-yellow after the addition of 1 equivalent of Grignard reagent indicating the complete formation of the intermediate ketene dithioacetal of type **2** and finally turned orange in the course of the rearrangement.

NMR: δ 13.7 (CH₃), 22.4 (CH₂), 28.7 (CH₂), 35.7 (dq, $^4J_{C-F}$ = 5.5 Hz, $^5J_{C-F}$ = 1.0 Hz, SCH₂), 39.5 (d, $^2J_{C-F}$ = 21.0 Hz, CF-CH₂), 103.0 (qd, $^1J_{C-F}$ = 200.5 Hz, $^2J_{C-F}$ = 29.5 Hz, CF), 121.3 (qd, $^1J_{C-F}$ = 287.0 Hz, $^2J_{C-F}$ = 29.5 Hz, CF₃), 121.7 (CH=CH₂), 127.5 (d, $^3J_{C-F}$ = 3.0 Hz, CH=CH₂), 223.9 (d, $^2J_{C-F}$ = 19.5 Hz, CS₂).

Butyl 2-(but-2'-enyl)-2,3,3,3-tetrafluorodithiopropanoate (4)

According to procedure A, reaction of 2-butenyl magnesium chloride with butyl pentafluorodithiopropanoate **1a** (730 mg, 2.89 mmol) in THF (10 mL) yielded butyl 2-(but-2'-enyl)-2,3,3,3-tetrafluorodithiopropanoate **4** (635 mg, 76%) as an orange liquid.

According to procedure A, reaction of 1-methyl-2-propenyl magnesium chloride with butyl pentafluorodithiopropanoate **1a** (240 mg, 0.95 mmol) in THF (5 mL) yielded butyl 2-(but-2'-enyl)-2,3,3,3-tetrafluorodithiopropanoate **4** (227 mg, 83%) as an orange liquid.

¹⁹F NMR: δ -77.8 (d, $^3J_{F-F}$ = 6.5 Hz, 3F, CF₃), -161.7 (ddt, $^3J_{F-F}$ = 6.5 Hz, $^3J_{F-H}$ = 12.5 Hz, $^3J_{F-H}$ = 35.5 Hz, 1F, CF); ¹H NMR: δ 0.86 (t, $^3J_{H-H}$ = 7.0 Hz, 3H, CH₂-CH₃), 1.35 (sext, $^3J_{H-H}$ = 7.0 Hz, 2H, CH₂-CH₃), 1.54 (d, $^3J_{H-H}$ = 5.0 Hz, 3H, CH=CH-CH₃), 1.58 (quint, $^3J_{H-H}$ = 7.0 Hz, 2H, CH₂-CH₂-CH₃), 2.56-3.21 (m, 4H, SCH₂, CH₂-CF), 5.17 (td, $^3J_{H-H}$ = 7.0 Hz, $^3J_{H-H}$ = 15.0 Hz, 1H, CH₂-CH=CH), 5.50 (td, $^3J_{H-H}$ = 5.0 Hz, $^3J_{H-H}$ = 15.0 Hz, 1H, CH=CH-CH₃); ¹³C NMR: δ 13.7 (CH₃), 18.4 (CH₃), 22.3 (CH₂), 29.0 (CH₂), 35.7 (d, $^4J_{C-F}$ = 5.5 Hz, SCH₂), 38.8 (d, $^2J_{C-F}$ = 21.0 Hz, CH₂-CF), 103.5 (qd, $^1J_{C-F}$ = 199.5 Hz, $^2J_{C-F}$ = 29.0 Hz, CF), 120.0 (CH=CH), 121.3 (qd, $^1J_{C-F}$ = 286.0 Hz, $^2J_{C-F}$ = 29.5 Hz, CF₃), 132.7 (CH=CH), 224.6 (d, $^2J_{C-F}$ = 19.0 Hz, CS₂).

Butyl 2-(2'-methylallyl)-2,3,3,3-tetrafluorodithiopropanoate (5)

According to procedure A, reaction of 2-methylallyl magnesium chloride with butyl pentafluorodithiopropanoate **1a** (507 mg, 2.01 mmol) in THF (10 mL) yielded butyl 2-(2'-methylallyl)-2,3,3,3-tetrafluorodithiopropanoate **5** (481 mg, 83%) as an orange liquid.

¹⁹F NMR: δ -78.1 (s, 3F, CF₃), -160.2 (ddt, $^3J_{F-F}$ = 6.5 Hz, $^3J_{F-H}$ = 12.0 Hz, $^3J_{F-H}$ = 38.0 Hz, 1F, CF); ¹H NMR: δ 0.94 (t, $^3J_{H-H}$ = 7.0 Hz, 3H, CH₂-CH₃), 1.45 (sext, $^3J_{H-H}$ = 7.0 Hz, 2H, CH₂-

CH₃), 1.67 (quint, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, CH₂-CH₂-CH₃), 1.77 (s, 3H, CH₃), 2.91 (dd, $^2J_{\text{AB}} = 14.0$ Hz, $^3J_{\text{H-F}} = 12.0$ Hz, 1H, CH_AH_B-CF), 3.20 (t, 2H, SCH₂), 3.28 (dd, $^2J_{\text{AB}} = 14.0$ Hz, $^3J_{\text{H-F}} = 38.0$ Hz, 1H, CH_AH_B-CF), 4.79 (s, 1H, =CHH), 4.92 (quint, $J_{\text{H-H}} = 1.5$ Hz, 1H, =CHH); ¹³C NMR: δ 13.7 (CH₃), 22.4 (CH₂), 24.0 (d, $^4J_{\text{C-F}} = 2.5$ Hz, CH₃), 28.7 (CH₂), 35.8 (d, $^4J_{\text{C-F}} = 5.5$ Hz, SCH₂), 42.3 (d, $^2J_{\text{C-F}} = 19.5$ Hz, CH₂-CF), 104.2 (qd, $^1J_{\text{C-F}} = 202.5$ Hz, $^2J_{\text{C-F}} = 28.5$ Hz, CF), 118.1 (s, C=CH₂), 121.1 (qd, $^1J_{\text{C-F}} = 288.0$ Hz, $^2J_{\text{C-F}} = 29.0$ Hz, CF₃), 136.8 (s, C=CH₂), 224.0 (d, $^2J_{\text{C-F}} = 20.0$ Hz, CS₂).

Ethyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate (6).

According to procedure A, reaction of ethyl magnesium bromide with allyl pentafluorodithiopropanoate **1b** (761 mg, 3.22 mmol) in THF (10 mL) yielded ethyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **6** (714 mg, 90%) as an orange liquid.

¹⁹F NMR: δ -77.9 (d, 3F, $^3J_{\text{F-F}} = 6.5$ Hz, CF₃), -161.8 (ddq, $^3J_{\text{F-F}} = 6.5$ Hz, $^3J_{\text{F-H}} = 13.0$ Hz, $^3J_{\text{F-H}} = 36.0$ Hz, 1F, CF); ¹H NMR: δ 1.35 (t, $^3J_{\text{H-H}} = 7.0$ Hz, 3H, CH₃), 2.95 (ddd, $J_{\text{AB}} = 14.5$ Hz, $^3J_{\text{H-F}} = 13.0$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, 1H, CF-CHH), 3.21 (q, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, SCH₂), 3.28 (ddd, $^3J_{\text{H-F}} = 36.0$ Hz, $J_{\text{AB}} = 14.5$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, 1H, CF-CHH), 5.19 (m, 2H, CH=CH₂), 5.76 (ddt, $^3J_{\text{H-H}} = 17.0$ Hz, $^3J_{\text{H-H}} = 10.0$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, 1H, CH=CH₂); ¹³C NMR: δ 11.5 (CH₃), 30.3 (d, $^4J_{\text{C-F}} = 5.5$ Hz, SCH₂), 39.5 (d, $^2J_{\text{C-F}} = 21.0$ Hz, CF-CH₂), 103.0 (dq, $^1J_{\text{C-F}} = 201.0$ Hz, $^2J_{\text{C-F}} = 28.5$ Hz, CF), 121.3 (qd, $^1J_{\text{C-F}} = 287.2$ Hz, $^2J_{\text{C-F}} = 28.5$ Hz, CF₃), 121.7 (CH=CH₂), 127.5 (d, $^3J_{\text{C-F}} = 3.0$ Hz, CH=CH₂), 223.9 (d, $^2J_{\text{C-F}} = 19.5$ Hz, CS₂).

Benzyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate (7)

According to procedure A, reaction of benzyl magnesium bromide with allyl pentafluorodithiopropanoate **1b** (700 mg, 2.77 mmol) in THF (10 mL) yielded benzyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **7** (693mg, 85%) as an orange oil.

¹⁹F NMR: δ -77.7 (d, $^3J_{\text{F-F}} = 6.5$ Hz, 3F, CF₃), -161.7 (ddq, $^3J_{\text{F-F}} = 6.0$ Hz, $^3J_{\text{F-H}} = 13.0$ Hz, $^3J_{\text{F-H}} = 35.0$ Hz, 1F, CF); ¹H NMR: δ 2.90 (dddt, $^2J_{\text{AB}} = 14.5$ Hz, $^3J_{\text{H-F}} = 13.0$ Hz, $^3J_{\text{H-H}} = 6.5$ Hz, $^4J_{\text{H-F}} = 1.0$ Hz, 1H, CH_AH_B-CF), 3.21 (dddt, $^2J_{\text{AB}} = 14.5$ Hz, $^3J_{\text{H-F}} = 35.5$ Hz, $^3J_{\text{H-H}} = 6.5$ Hz, $^4J_{\text{H-F}} = 1.0$ Hz, 1H, CH_AH_B-CF), 4.31 (s, 2H, SCH₂), 5.12 (m, 2H, CH=CH₂), 5.60 (ddt, $^3J_{\text{H-H}}$

= 17.0 Hz, $^3J_{\text{H-H}} = 9.5$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, 1H, CH=CH₂), 7.23 (s, 5H, Ph); ^{13}C NMR: δ 39.8 (dq, $^2J_{\text{C-F}} = 20.5$ Hz, $^3J_{\text{C-F}} = 1.0$ Hz, CF-CH₂), 41.5 (d, $^4J_{\text{C-F}} = 6.0$ Hz, SCH₂), 103.6 (qd, $^1J_{\text{C-F}} = 200.0$ Hz, $^2J_{\text{C-F}} = 29.0$ Hz, CF), 121.3 (qd, $^1J_{\text{C-F}} = 286.5$ Hz, $^2J_{\text{C-F}} = 29.5$ Hz, CF₃), 122.2 (CH=CH₂), 127.7 (d, $^3J_{\text{C-F}} = 3.0$ Hz, CH=CH₂), 128.5 (C-aromatic para), 129.2 (2 C-aromatic), 129.7 (2 C-aromatic), 133.9 (C-aromatic ipso), 223.5 (d, $^2J_{\text{C-F}} = 20.0$ Hz, CS₂).

Reaction of Grignard reagents with tetrafluoro monounsaturated dithioesters 3-5 or 7 for the preparation of trifluoromethyl bis(unsaturated) dithioesters 9-14 (Procedure B):

To a solution of unsaturated dithioester **3-5** or **7** (1 equiv) in THF was added, at -78°C and under Ar, allylic Grignard reagent (1 equiv).¹ The reaction was then slowly warmed to room temperature. After 15 h of stirring, ^{19}F NMR analysis of the crude reaction mixture indicates the presence of both intermediate ketene dithioacetal of type **8** and the final bis(unsaturated) dithioester. The solution was then heated at reflux until completion of the rearrangement (checked by ^{19}F NMR). The reaction mixture was diluted with diethyl ether, washed with sat. aq. sol. NH₄Cl and brine. The organic layer was dried (MgSO₄) and concentrated under reduced pressure. Chromatography on silica gel (pentane) of the residue gave trifluoromethyl bis(unsaturated) dithioesters **9-14**.

“one-pot” preparation of trifluoromethyl bis(unsaturated) dithioesters 9,12 starting from pentafluoro dithioesters 1a-b (Procedure C). To a solution of pentafluorodithioester **1** (1 equiv) in THF was added, at -78°C and under Ar, the first Grignard reagent (1 equiv).¹ The reaction was then slowly warmed to room temperature and after 15 h of stirring the completion of the reaction was checked by ^{19}F NMR. The reaction was then cooled down to -78°C and the second Grignard reagent (1 equiv)¹ was added. The reaction was then slowly warmed to room temperature. After 15 h of stirring, ^{19}F NMR analysis of the crude reaction mixture indicates the presence of both intermediate ketene dithioacetal of type **8** and the final bis(unsaturated) dithioester **9** or **12**. The solution was heated at reflux until the completion of the rearrangement (checked by ^{19}F NMR). The reaction mixture was diluted with diethyl ether, washed with sat. aq. sol. NH₄Cl and brine. The organic layer was dried (MgSO₄) and concentrated under reduced pressure. Chromatography on silica gel (pentane) of the residue gave trifluoromethyl bis(unsaturated) dithioesters **9,12**.

Butyl 2,2-diallyl-3,3,3-trifluorodithiopropanoate (9)

According to procedure B, reaction of allyl magnesium bromide with butyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **3** (541 mg, 2.20 mmol) in THF (15 mL) led, after 15 min of reflux, to butyl 2,2-diallyl-3,3,3-trifluorodithiopropanoate **9** (456 mg, 70%) as an orange liquid.

According to “one-pot” procedure C, reaction of allyl magnesium bromide with butyl pentafluorodithiopropanoate **1a** (487 mg, 1.93 mmol) in THF (5 mL), followed by reaction with a second equivalent of allyl magnesium bromide (in solution in THF, 1 equiv) led, after 15 min of reflux, to butyl 2,2-diallyl-3,3,3-trifluorodithiopropanoate **9** (349 mg, 61%) as an orange liquid.

According to “one-pot” procedure C, reaction of *n*-butyl magnesium chloride with allyl pentafluorodithiopropanoate **1b** (232 mg, 0.98 mmol) in THF (5 mL), followed by reaction with allyl magnesium bromide (in solution in THF, 1 equiv) led, after 15 min of reflux, to butyl 2,2-diallyl-3,3,3-trifluorodithiopropanoate **9** (215 mg, 74%) as an orange liquid.

According to “one-pot” procedure C, reaction of allyl magnesium bromide with allyl pentafluorodithiopropanoate **1b** (212 mg, 0.90 mmol) in THF (5 mL), followed by reaction with *n*-butyl magnesium chloride (in solution in THF, 1 equiv) led, after 15 min of reflux, to butyl 2,2-diallyl-3,3,3-trifluorodithiopropanoate **9** (181 mg, 68%) as an orange liquid.

^{19}F NMR: δ -67.6 (s, 3F, CF_3); ^1H NMR: δ 0.94 (t, 3H, CH_3), 1.45 (sext, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, $\text{CH}_2\text{-CH}_3$), 1.67 (quint, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, $\text{CH}_2\text{-CH}_2\text{-CH}_3$), 2.99 (dd, $^2J_{\text{AB}} = 14.5$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, 4H, 2 $\text{CH}_2\text{-CH=CH}_2$), 3.23 (t, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, SCH_2), 5.12 (m, 2H, 2 CH=CHH), 5.15 (ddt, $^3J_{\text{H-H}} = 17.0$ Hz, $^2J_{\text{H-H}} = 2.0$ Hz, $^4J_{\text{H-H}} = 1.5$ Hz, 2H, 2 CH=CHH), 5.85 (ddtq, $^3J_{\text{H-H}} = 17.0$ Hz, $^3J_{\text{H-H}} = 10.0$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, $^5J_{\text{H-F}} = 1.0$ Hz, 2H, 2 CH=CH_2); ^{13}C NMR: δ 13.8 (CH_3), 22.5 (CH_2), 28.8 (CH_2), 37.5 (SCH_2), 39.3 (q, $^3J_{\text{C-F}} = 1.5$ Hz, 2 $\text{CH}_2\text{-CH=CH}_2$), 65.0 (q, $^2J_{\text{C-F}} = 21.5$ Hz, C-CF_3), 119.6 (2 CH=CH_2), 125.5 (q, $^1J_{\text{C-F}} = 286.0$ Hz, CF_3), 132.0 (2 CH=CH_2), 231.2 (CS_2).

Benzyl 2,2-diallyl-3,3,3-trifluorodithiopropanoate (10)

According to procedure B, reaction of allyl magnesium bromide with benzyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **7** (538 mg, 1.88 mmol) in THF (10 mL) led, after 15 min of reflux, to benzyl 2,2-diallyl-3,3,3-trifluorodithiopropanoate **10** (466 mg, 75%) as an orange oil.

^{19}F NMR: δ -67.5 (s, 3F, CF_3); ^1H NMR: δ 3.00 (m, 4H, 2 $\text{CH}_2\text{-CH=CH}_2$), 4.40 (s, 2H, SCH_2), 5.12 (m, 4H, 2 CH=CH_2), 5.77 (m, 2H, 2 CH=CH_2), 7.31 (s, 5H, Ph); ^{13}C NMR: δ 39.5 (q, $^3J_{\text{C-F}} = 1.5$ Hz, 2 $\text{CH}_2\text{-CH=CH}_2$), 43.4 (SCH_2), 65.3 (q, $^2J_{\text{C-F}} = 22.0$ Hz, 2 C-CF_3), 120.1 (q, $^5J_{\text{C-F}} = 0.5$ Hz, 2 CH=CH_2), 125.7 (q, $^1J_{\text{C-F}} = 287.0$ Hz, CF_3), 128.3 (C-aromatic para), 129.2 (2 C-aromatic), 129.8 (2 C-aromatic), 132.1 (q, $^4J_{\text{C-F}} = 1.0$ Hz, 2 CH=CH_2), 134.3 (C-aromatic ipso), 230.6 (q, $^3J_{\text{C-F}} = 1.0$ Hz, CS_2).

Butyl 2-allyl-2-(but-2'-enyl)-3,3,3-trifluorodithiopropanoate (11)

According to procedure B, reaction of 2-butenyl magnesium chloride with butyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **3** (107 mg, 0.43 mmol) in THF (5 mL) led, after 15 min of reflux, to butyl 2-allyl-2-(but-2'-enyl)-3,3,3-trifluorodithiopropanoate **11** (79 mg, 59%) as an orange liquid.

According to procedure B, reaction of allyl magnesium chloride with butyl 2-(but-2'-enyl)-2,3,3,3-tetrafluorodithiopropanoate **4** (259 mg, 0.90 mmol) in THF (10 mL) led, after 15 min of reflux, to butyl 2-allyl-2-(but-2'-enyl)-3,3,3-trifluorodithiopropanoate **11** (190 mg, 68%) as an orange liquid.

^{19}F NMR: δ -67.5 (s, 3F, CF_3); ^1H NMR: δ 0.95 (t, 3H, $^3J_{\text{H-H}} = 7.0$ Hz, $\text{CH}_2\text{-CH}_3$), 1.43 (sext, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, $\text{CH}_2\text{-CH}_3$), 1.65 (d, $^3J_{\text{H-H}} = 7.0$ Hz, 3H, CH=CH-CH_3), 1.68 (m, 2H, $\text{CH}_2\text{-CH}_2\text{-CH}_3$), 2.69 (m, 1H, $\text{CH}_\text{A}\text{H}_\text{B}\text{-CH=}$), 2.91 (m, 2H, CH_2), 3.04 (dd, $J_{\text{AB}} = 14.5$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, 1H, $\text{CH}_\text{A}\text{H}_\text{B}\text{-CH=}$), 3.23 (t, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, SCH_2), 5.12 (m, 2H, CH=CH_2), 5.37 (tdd, $^3J_{\text{H-H}} = 15.0$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, $^5J_{\text{H-F}} = 1.0$ Hz, 1H, CH=CH-CH_3), 5.60 (qdd, $^3J_{\text{H-H}} =$

15.0 Hz, $^3J_{\text{H-H}} = 6.5$ Hz, $^4J_{\text{H,H}} = 1.0$ Hz, 1H, CH=CH-CH₃), 5.76 (ddtq, $^3J_{\text{H-H}} = 17.5$ Hz, $^3J_{\text{H-H}} = 10.0$ Hz, $^3J_{\text{H-H}} = 7.5$ Hz, $^5J_{\text{H-F}} = 1.0$ Hz, 1H, CH=CH₂); ^{13}C NMR: δ 14.0 (CH₃), 18.4 (CH₃), 22.8 (CH₂), 29.1 (CH₂), 37.7 (SCH₂), 38.5 (CH₂), 39.5 (CH₂), 65.4 (q, $^2J_{\text{C-F}} = 21.5$ Hz, C-CF₃), 119.6 (q, $^4J_{\text{C-F}} = 1.0$ Hz, CH=CH₂), 124.4 (q, $^5J_{\text{C-F}} = 1.0$ Hz, CH=), 125.8 (q, $^1J_{\text{C-F}} = 287.0$ Hz, CF₃), 130.7 (CH=CH-CH₃), 132.5 (q, $^4J_{\text{C-F}} = 1.0$ Hz, CH=), 231.8 (q, $^4J_{\text{C-F}} = 1.0$ Hz, CS₂).

Butyl 2-allyl-2-(2'-methylallyl)-3,3,3-trifluorodithiopropanoate (12)

According to procedure B, reaction of 2-methylallyl magnesium chloride with butyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **3** (525 mg, 2.13 mmol) in THF (10 mL) led, after 45 min of reflux, to butyl 2-allyl-2-(2'-methylallyl)-3,3,3-trifluorodithiopropanoate **12** (516 mg, 78%) as an orange liquid.

According to procedure B, reaction of allyl magnesium chloride with butyl 2-(2'-methylallyl)-2,3,3,3-tetrafluorodithiopropanoate **5** (102 mg, 0.35 mmol) in THF (5 mL) led, after 1 h 30 min of reflux, to butyl 2-allyl-2-(2'-methylallyl)-3,3,3-trifluorodithiopropanoate **12** (71 mg, 65%) as an orange liquid.

According to “one-pot” procedure C, reaction of allyl magnesium bromide with butyl pentafluorodithiopropanoate **1a** (543 mg, 2.15 mmol) in THF (10 mL), followed by reaction with 2-methylallyl magnesium bromide led, after 45 min of reflux, to butyl 2-allyl-2-(2'-methylallyl)-3,3,3-trifluorodithiopropanoate **12** (494 mg, 74%) as an orange liquid.

^{19}F NMR: δ -67.5 (s, 3F, CF₃); ^1H NMR: δ 0.95 (t, 3H, $^3J_{\text{H-H}} = 7.0$ Hz, CH₂-CH₃), 1.43 (sext, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, CH₂-CH₃), 1.65 (d, $^3J_{\text{H-H}} = 7.0$ Hz, 3H, CH=CH-CH₃), 1.68 (m, 2H, CH₂-CH₂-CH₃), 2.69 (m, 1H, CH_AH_B-CH=), 2.91 (m, 2H, CH₂), 3.04 (dd, $J_{\text{AB}} = 14.5$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, 1H, CH_AH_B-CH=), 3.23 (t, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, SCH₂), 5.12 (m, 2H, CH=CH₂), 5.37 (tdd, $^3J_{\text{H-H}} = 15.0$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, $^5J_{\text{H-F}} = 1.0$ Hz, 1H, CH=CH-CH₃), 5.60 (qdd, $^3J_{\text{H-H}} = 15.0$ Hz, $^3J_{\text{H-H}} = 6.5$ Hz, $^4J_{\text{H,H}} = 1.0$ Hz, 1H, CH=CH-CH₃), 5.76 (ddtq, $^3J_{\text{H-H}} = 17.5$ Hz, $^3J_{\text{H-H}} = 10.0$ Hz, $^3J_{\text{H-H}} = 7.5$ Hz, $^5J_{\text{H-F}} = 1.0$ Hz, 1H, CH=CH₂); ^{13}C NMR: δ 14.0 (CH₃), 22.8 (CH₂), 24.7 (CH₃), 28.9 (CH₂), 29.1 (CH₂), 37.7 (SCH₂), 38.5 (CH₂), 39.5 (CH₂), 65.9 (q, $^2J_{\text{C-F}} = 21.5$ Hz, C-CF₃), 116.3 (=CH₂), 119.2 (=CH₂), 125.7 (q, $^1J_{\text{C-F}} = 287.0$ Hz, CF₃), 132.9 (q, $^4J_{\text{C-F}} = 1.5$ Hz, CH=CH₂), 139.8 (C=CH₂), 232.1 (CS₂).

Benzyl 2-allyl-2-(2'-methylallyl)-3,3,3-trifluorodithiopropanoate (13)

According to procedure B, reaction of 2-methylallyl magnesium chloride with benzyl 2-allyl-2,3,3,3-tetrafluorodithiopropanoate **7** (127 mg, 0.41 mmol) in THF (5 mL) led, after 1 h of reflux, to benzyl 2-allyl-2-(2'-methylallyl)-3,3,3-trifluorodithiopropanoate **13** (124 mg, 88%) as an orange oil.

^{19}F NMR: δ -67.2 (s, 3F, CF_3); ^1H NMR: δ 1.68 (s, 3H, CH_3), 2.86 (d, $J_{\text{AB}} = 15.0$ Hz, 1H, $\text{CH}_\text{A}\text{H}_\text{B}=\text{C}=\text{)$, 2.89 (m, 1H, $\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}=\text{)$, 3.11 (d, $J_{\text{AB}} = 15.0$ Hz, 1H, $\text{CH}_\text{A}\text{H}_\text{B}=\text{C}=\text{)$, 3.19 (dd, $J_{\text{AB}} = 14.5$ Hz, $^3J_{\text{H-H}} = 7.0$ Hz, 1H, $\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}=\text{)$, 4.41 (s, 2H, SCH_2), 4.74 (s, 1H, $\text{C}=\text{CHH}$), 4.88 (s, 1H, $\text{C}=\text{CHH}$), 5.09 (dq, $^3J_{\text{H-H}} = 17.0$ Hz, $^6J_{\text{H-F}} = 1.5$ Hz, 1H, $\text{CH}=\text{CHH}$), 5.13 (dq, $^3J_{\text{H-H}} = 10.0$ Hz, $^6J_{\text{H-F}} = 1.0$ Hz, 1H, $\text{CH}=\text{CHH}$), 5.86 (m, 1H, $\text{CH}=\text{CH}_2$), 7.31 (s, 5H, Ph); ^{13}C NMR: δ 24.8 (q, $^5J_{\text{C-F}} = 0.5$ Hz, CH_3), 40.1 (SCH_2), 43.3 ($\text{CH}_2-\text{C}-\text{CF}_3$), 43.8 (q, $^3J_{\text{C-F}} = 1.5$ Hz, $\text{CH}_2-\text{C}-\text{CF}_3$), 66.2 (q, $^2J_{\text{C-F}} = 21.5$ Hz, $\text{C}-\text{CF}_3$), 116.3 ($=\text{CH}_2$), 119.4 ($=\text{CH}_2$), 128.3 (C-aromatic para), 129.1 (2 C-aromatic), 129.8 (2 C-aromatic), 133.0 (q, $^4J_{\text{C-F}} = 1.5$ Hz, $\text{CH}=\text{CH}_2$), 134.4 (C-aromatic ipso), 140.0 ($\text{C}=\text{CH}_2$), 231.6 (CS_2).

Butyl 2,2-di(but-2'-enyl)-3,3,3-trifluorodithiopropanoate (14)

According to procedure B, reaction of 1-methyl-2-propenyl magnesium chloride with butyl 2-(but-2'-enyl)-2,3,3,3-tetrafluorodithiopropanoate **4** (233 mg, 0.81 mol) in THF (5 mL) led, after 1 h of reflux, to butyl 2,2-di(but-2'-enyl)-3,3,3-trifluorodithiopropanoate **14** (134 mg, 51%) as an orange liquid.

^{19}F NMR: δ -67.4 (s, 3F, CF_3); ^1H NMR: δ 0.95 (t, $^3J_{\text{H-H}} = 7.0$ Hz, 3H, CH_3), 1.45 (sext, $^3J_{\text{H-H}} = 7.0$ Hz, 2H, CH_2-CH_3), 1.64 (d, $^3J_{\text{H-H}} = 6.5$ Hz, 3H, $\text{CH}=\text{CH}-\text{CH}_3$), 1.65 (d, $^3J_{\text{H-H}} = 6.5$ Hz, 3H, $\text{CH}=\text{CH}-\text{CH}_3$), 1.66 (m, 2H, $\text{CH}_2-\text{CH}_2-\text{CH}_3$), 2.89 (m, 4H, 2 $\text{CH}_2-\text{CH}=\text{CH}$), 3.22 (t, $^3J_{\text{H-H}} = 7.5$ Hz, 2H, SCH_2), 5.40 (m, 2H, 2 $\text{CH}=\text{CH}$), 5.55 (m, 2H, 2 $\text{CH}=\text{CH}$); ^{13}C NMR: δ 14.0 (CH_3), 18.5 (CH_3), 22.8 (CH_2), 29.1 (CH_2), 37.6 (SCH_2), 38.4 (q, $^3J_{\text{C-F}} = 1.5$ Hz, CH_2),

65.6 (q, $^2J_{\text{C-F}} = 21.5$ Hz, C-CF₃), 124.7 (CH=CH), 125.9 (q, $^1J_{\text{C-F}} = 287.0$ Hz, CF₃), 130.3 (CH=CH), 232.2 (CS₂).

^9F NMR

S13

¹H NMR

S14

^{13}C NMR

120000

100000

80000

60000

40000

20000

0

200

180

160

140

120

100

80

60

40

20

0

(p p m)

S15

¹⁹F NMR

¹H NMR

S17

¹³C NMR

S18

¹⁹F NMR

S19

¹H NMR

S20

¹³C NMR

S21

¹⁹F NMR

S22

¹H NMR

S23

^{13}C NMR

1 0 0 0 0 0

8 0 0 0 0

6 0 0 0 0

4 0 0 0 0

2 0 0 0 0

0

ether

x

ether

x

(p p m)

S24

^{19}F NMR

S25

¹H NMR

S26

^{19}F NMR

¹H NMR

5 e + 0 0 5

4 e + 0 0 5

3 e + 0 0 5

2 e + 0 0 5

1 e + 0 0 5

0 e + 0 0 0

¹³C NMR

4 0 0 0 0

3 5 0 0 0

3 0 0 0 0

2 5 0 0 0

2 0 0 0 0

1 5 0 0 0

1 0 0 0 0

5 0 0 0

0

2 2 0

2 0 0

1 8 0

1 6 0

1 4 0

1 2 0

1 0 0

8 0

6 0

4 0

2 0

0

(p p m)

S30

^{19}F NMR

S31

¹³C NMR

. 0 e + 0 0 6

. 5 e + 0 0 6

. 0 e + 0 0 6

. 0 e + 0 0 5

. 0 e + 0 0 0

^{19}F NMR

1 2 0 0 0 0

1 0 0 0 0 0

8 0 0 0 0

6 0 0 0 0

4 0 0 0 0

2 0 0 0 0

0

- 6 1

- 6 2

- 6 3

- 6 4

- 6 5

- 6 6

- 6 7

- 6 8

- 6 9

- 7 0

- 7 1

- 7 2

- 7 3

- 7 4

- 7 5

- 7 6

(p p m)

9

S34

¹⁹F NMR

6 e + 0 0 5

5 e + 0 0 5

4 e + 0 0 5

3 e + 0 0 5

2 e + 0 0 5

1 e + 0 0 5

0 e + 0 0 0

- 6 4 . 0 - 6 5 . 0 - 6 6 . 0 - 6 7 . 0 - 6 8 . 0 - 6 9 . 0 - 7 0 . 0 - 7 1 . 0 - 7 2 . 0 - 7 3 . 0

(p p m)

¹H NMR

S38

¹³C NMR

. 4 e + 0 0 6

. 2 e + 0 0 6

. 0 e + 0 0 6

. 0 e + 0 0 5

. 0 e + 0 0 5

. 0 e + 0 0 5

. 0 e + 0 0 5

. 0 e + 0 0 0

^{19}F NMR

S40

¹H NMR

S41

¹³C NMR

^{19}F NMR

S43

¹H NMR

S44

¹³C NMR

. 6 e + 0 0 6

. 4 e + 0 0 6

. 2 e + 0 0 6

. 0 e + 0 0 6

. 0 e + 0 0 5

. 0 e + 0 0 5

. 0 e + 0 0 5

. 0 e + 0 0 5

. 0 e + 0 0 0

220

200

180

160

140

120

100

80

60

40

20

0

(p p m)

S45

^1H NMR

S47

¹⁹F NMR

S49

S50

¹³C NMR

5 e + 0 0 6

4 e + 0 0 6

3 e + 0 0 6

2 e + 0 0 6

1 e + 0 0 6

0 e + 0 0 0

2 4 0 2 2 0 2 0 0 1 8 0 1 6 0 1 4 0 1 2 0 1 0 0 8 0 6 0 4 0 2 0 0

(p p m)

S51