Supporting Information

Syntheses of New Liquid Crystals Derived From Thiophene Connected to Heterocyclic1,2,4-Oxadiazole

Edivandro Giroto,[†] Ivan H. Bechtold[‡] and Hugo Gallardo^{†,*}

[†]Departamento de Química and [‡]Departamento de Física, Universidade Federal de Santa Catarina, Florianópolis-SC, 88040-900,

*Email: hugo.gallardo@ufsc.br; fax: +55 48 37216850; tel: +55 48 37219544

Experimental

1.0. NMR spectra of compounds 8a-d and 13a-d.

Figure S1. ¹H NMR (a) spectrum of compound **8a** (CDCl₃).

Figure S2. ^1H NMR (a) and ^{13}C NMR (b) spectrum of compound **8b** (CDCl_3).

a)

b)

Figure S3. ^1H NMR (a) and ^{13}C NMR (b) spectrum of compound **8c** (CDCl_3).

a)

b)

Figure S4. ^1H NMR (a) and ^{13}C NMR (b) spectrum of compound **8d** (CDCl_3).

a)

Figure S5. ^1H NMR (a) and ^{13}C NMR (b) spectrum of compound **13a** (CDCl_3).

b)

Figure S6. ^1H NMR (a) and ^{13}C NMR (b) spectrum of compound **13b** (CDCl_3).

a)

b)

Figure S7. ^1H NMR (a) and ^{13}C NMR (b) spectrum of compound **13**(CDCl_3).

a)

b)

Figure S8. ^1H NMR (a) and ^{13}C NMR (b) spectrum of compound **13d** (CDCl_3).

b)

2.0 MS spectraDSC Thermograms