Supporting Information

Nickel(0)-Catalyzed Asymmetric Hydrocyanation of 1,3-Dienes

Biswajit Saha, T.V. RajanBabu

Department of Chemistry, 100 W. 18th Avenue, The Ohio State University,
Columbus, Ohio 43210

Supporting Information

Toxicity of Hydrogen Cyanide. Hydrogen cyanide is a highly toxic, volatile liquid (bp 27 °C) that is also susceptible to explosive polymerization in the presence of base catalysts. It should be handled only in well-ventilated fume hood or drybox. Sensible precautions include not working alone and having available proper first aid equipment. When handling large quantities of HCN, it should be handled by a team of at least two technically qualified individuals who have received appropriate medical training for treating HCN poisoning (for details see: *Prudent Practices for Handling Hazardous Chemicals in Laboratories*: National Academy Press: Washington, DC, 1981; pp 45-47). HCN prepared by the Ziegler procedure (Ziegler, K. *Org. Syn. Coll. Vol.* **1941**, *1*, 341) was diluted with cold (<5 °C) toluene to make a 2 M solution which was kept inside the freezer compartment of a drybox. The solution was further stabilized by adding ~ 20 milligrams of P₂O₅ to ~ 100 mL of the above solution. Excess HCN maybe disposed of by burning or in case of small amounts, by adding to aqueous sodium hypochlorite (which converts it to the cyanate). In the present case no more than 2 mmol (2 mL of toluene solution) was used at one time.

General methods

All catalyzed reactions were carried out under an inert atmosphere of nitrogen in a Vacuum Atmosphere drybox, or by Schlenk techniques. Methylene chloride, hexane, toluene were distilled from calcium hydride under nitrogen and stored over molecular sieves. Tetrahydrofuran (THF) was distilled from benzophenone ketyl under nitrogen and stored over molecular sieves. All chemicals were purchased from Aldrich Chemical Company unless otherwise noted. Chiral phosphinite ligands (**L1**, **L2**, **L3**, **L4**)¹, 1-(*E*)-phenylbutadiene², 1-vinyl-3,4-dihydronaphthalene³, 1-vinylindene⁴, methyl (2*E*)-2,4-pentadienoate,⁵ 1-methyl (*E*)-phenylbutadiene⁶ were prepared according to the literature procedures. Analytical TLC was done on E. Merck precoated (0.25

mm) silica gel 60F₂₅₄ plates. Column chromatography was conducted by using silica gel 40. ¹H NMR spectra were recorded on a Bruker DPX-500 (500 MHz) and DPX-400 (400 MHz). Gas chromatographic analyses were performed on a Hewlett-Packard 5890 equipped with an HP-ultra-1 cross-linked methyl silicon capillary column (25 m length x 0.2 mm i.d.) and an FID detector connected to an HP 3396 integrator. Chiral gas chromatographic separations were accomplished using cyclodex-B column (25m x 0.25 mm, 0.12 μm film thickness), purchased from Chrompack (1130 Route 202 South, Raritan, New Jersey 08869). As carrier gas helium was used. HPLC analysis of chiral compounds was performed using Daicel Chiralcel OD, OJ or Chiralpak AS-H columns, and, eluent hexane/isopropanol was used as solvents where base-line separation was obtained. Optical rotations were recorded on a Perkin-Elmer Model 241 polarimeter at the sodium D line in chloroform.

Preparation of Ligands

To a 100 mL flask was added chlorodiphenylphosphine (2.03 g, 9.2 mmol), 21 mL of anhydrous toluene and 4-dimethylaminopyridine (DMAP) (1.23 g, 10 mmol). To this solution was then added *trans*-1,2-dihydroxy cyclohexane (0.5 g, 4.4 mmol) in portions over 5 minutes. The mixture was stirred for additional 16 h, and the precipitated DMAP hydrochloride is filtered inside the drybox, washed with toluene (2 x 2 mL), and the filtrate was concentrated to dryness in vacuo to provide the crude ligand (1.84 g) which was crystallized from benzene to obtain 1.4 g white solid (65% yield).

The ligands **L1-L4** were synthesized by similar procedure.¹

General Procedure for Hydrocyanation of 1,3-dienes. To a mixture of the phosphinite ligand and Ni(COD)₂ was added 0.2 mL toluene and the mixture was stirred for 5 minutes at room temperature, 0.2 mL of toluene solution of respective diene (0.5 mmol) (see Table 1) was added dropwise, and the mixture was stirred for 10 minutes at room temperature and after that 0.2 mL

of 2M HCN in toluene was added slowly, dropwise, and the reaction was stirred for 20 – 48 h at room temperature, or heated at 50 °C for 24 h in a sealed vial. The reaction flask was taken out from the dry box and N₂ gas passed into the reaction mixture under well ventilated hood to remove excess HCN. The crude mixture was passed through a small pad of silica to remove the catalyst and the solvent was removed under vacuo and purified by short silicagel column chromatography to afford pure product using 100% hexane to hexane/Et₂O (19:1) as eluent. The products were subjected to gas chromatography and spectroscopic analysis to ascertain purity and isomeric composition. Isolated yield of the desired products are given in the Table 1 of paper.

(*E*)-2-Methyl-4-phenylbut-3-enitrile (1):

Catalyst : 10 mol%, 20 h at room temperature. Colorless oil. ¹H NMR (500 MHz, CDCl₃) δ 7.42-7.3 (m, 5 H), 6.75 (d, *J* = 16 Hz, 1 H), 6.07 (dd, *J* = 6.5 Hz, 15.5 Hz, 1 H), 3.52 (m, 1 H), 1.53 (d, *J* = 7.5 Hz, 3 H); ¹³C NMR (125.7 MHz, CDCl₃) δ 135.74, 132.55, 128.74, 128.29, 126.57, 124.38, 120.9, 28.38, 19.06.

(*E*)-2-Methyl-4-phenylpent-3-enitrile (2):

Catalyst : 10 mol%. 24h at 50°C. Colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.41-7.29 (m, 5 H), 5.66 (d, *J* = 8.4 Hz, 1H), 3.64 (dq, *J* = 7.2, 8.4 Hz, 1 H), 2.14 (d, *J* = 1.2 Hz, 3 H), 1.51 (d, *J* = 7.2 Hz, 3 H); ¹³C NMR (125.7 MHz, CDCl₃) δ 142.05, 139.66, 128.43, 127.81, 125.91, 122.80, 121.61, 24.57, 19.06, 16.42.

Cyclohex-2-enecarbonitrile (3):

Catalyst: 5 mol%. 20h at room temperature. Colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 5.95-5.91 (m, 1 H), 5.64-5.61 (m, 1 H), 3.25-3.23 (m, 1 H), 2.09-2.04 (m, 2 H), 1.97-

1.92 (m, 2H), 1.85-1.82 (m, 1H), 1.69-1.65 (m, 1H). ^{13}C NMR (125.7 MHz, CDCl_3) 131.84, 121.45, 120.75, 26.49, 26.26, 24.17, 19.91.

2-(3,4-Dihydronaphthalene)-propionitrile (4):

Catalyst: 10 mol%, 24h at room temperature. Colorless oil. ^1H NMR (500 MHz, CDCl_3) δ 7.28-7.2 (m, 4 H), 6.36 (t, J = 4.5 Hz, 1 H), 3.92 (q, J = 7 Hz, 1 H), 2.79 (t, J = 8 Hz, 2 H), 2.39-2.35 (m, 2 H), 1.6 (d, J = 7.5 Hz, 3 H). ^{13}C NMR (125.7 MHz, CDCl_3) δ 136.86, 132.74, 131.83, 128.19, 127.61, 127.23, 126.63, 121.84, 121.66, 27.98, 27.88, 22.9, 18.88.

2-(Indene)-propionitrile (5):

Catalyst : 5 mol%, 48h at room temperature. Colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.53-7.27 (m, 4 H), 6.60 (br, m, 1 H), 3.96-3.93 (m, 1 H), 3.44 (br, m, 2 H), 1.72 (d, J = 7.2 Hz, 3 H). ^{13}C NMR (100.6 MHz, CDCl_3) 144.43, 141.95, 139.54, 130.63, 126.39, 125.49, 124.26, 120.90, 118.94, 37.77, 24.86, 18.14.

2,6-Dimethyl-2-propenyl-hept-5-enenitrile (6):

Catalyst : 10 mol%. 24 h at 50 $^{\circ}\text{C}$. Colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 5.88-5.83 (m, 1 H), 5.20 (d, J = 15.6 Hz, 1 H), 5.06 (t, J = 1.6 Hz, 1 H), 2.09-2.06 (m, 2 H), 1.71 (dd, J = 1.6 Hz, 6.4 Hz, 3 H), 1.68 (br, s, 3 H), 1.60 (br, s, 3 H), 1.53-1.49 (m, 2 H), 1.38 (s, 3 H); ^{13}C NMR (125.7 MHz, CDCl_3) δ 132.79, 131.13, 126.65, 122.95, 122.63, 40.40, 39.43, 26.32, 25.66, 24.04, 17.69, 17.55.

2-Propenyl-hexanenitrile (A) and 2-Methyl-oct-3-enenitrile (B) [7 and 8]:

(A)

(B)

Catalyst : 10 mol%, 24 h at 50°C. Colorless oil. For compound

7: ^1H NMR (400 MHz, CDCl_3) δ 5.81-5.75 (m, 1 H), 5.35-5.29 (m, 1 H), 3.15-3.14 (m, 1 H), 2.03 (q, J = 6.8 Hz, 2 H), 1.70 (dd, J = 1.2, 5.2 Hz, 3 H), 1.64-1.6 (m, 2 H), 1.45-1.26 (m, 4 H), 0.92-0.86 (m, 3 H). For compound **8**: ^1H NMR (400 MHz, CDCl_3) δ 5.81-5.75 (m, 1 H), 5.35-5.29 (m, 1 H), 3.25 (dq, J = 6.4, 7.2 Hz, 1 H), 2.03 (q, J = 6.8 Hz, 2 H), 1.64-1.6 (m, 2 H), 1.38 (d, J = 6 Hz, 3 H), 1.45-1.26 (m, 4 H), 0.92-0.86 (m, 3 H). For compounds **7** and **8**: ^{13}C NMR (125.7 MHz) 134.02, 129.34, 125.00, 124.86, 121.44, 120.74, 34.03, 32.97, 31.72, 31.02, 28.84, 28.02, 22.12, 19.10, 17.58, 13.85, 13.79.

General Procedure for Asymmetric Hydrocyanation of 1,3-dienes: To a mixture of the phosphinite ligand (**L1** – **L4**) (2.1 mol%) and $\text{Ni}(\text{COD})_2$ (2 mol%) was added 0.2 mL of toluene and the mixture was stirred for 5 minutes at room temperature, 0.2 mL toluene solution of respective dienes (see Table 2, 0.5 mmol) was added dropwise. The solution turned brown and the mixture was stirred for 10 minutes at room temperature and after that 0.14 mL of 2M HCN in toluene was added slowly, dropwise and the reaction was stirred for 12 h and additional 1 mol% of $\text{Ni}(\text{COD})_2$, 0.1 mL of 2M HCN in toluene were added respectively and the reaction mixture was stirred for additional 12 to 48 h. The reaction flask was taken out from the dry box and N_2 gas passed into the reaction mixture under well-ventilated hood to remove any excess HCN. The crude mixture was passed through a small pad of silica to remove the catalyst and the solvent was removed under vacuo and purified by short silica gel column chromatography to afford pure product using 100% hexane to hexane/ Et_2O (19:1) as eluent. The enantiomeric ratio of the final product was determined using chiral HPLC or GC (attached).

(E)-2-Methyl-4-phenylbut-3-enenitrile (1):

Enantiomer Separation: Chiralcel OD Column 95:5 hexanes/2-propanol, 0.75 mL/min, 254 nm, $R_{t1} = 11.34$ min, $R_{t2} = 12.23$ min. 78% ee, $([\alpha]_D^{20} = +19, c 0.7$ in CHCl_3).

2-(3,4-dihydronaphthalene)-propionitrile (4):

Enantiomer Separation: Chiralcel OJ Column 95:5 hexanes/2-propanol, 0.5 mL/min, 254 nm, $R_{t1} = 20.53$ min, $R_{t2} = 23.62$ min. 75% ee, $([\alpha]_D^{20} = -61, c 0.18$ in CHCl_3).

2-(indene)-propionitrile (5):

Enantiomer Separation: Chiralcel OD Column 95:5 hexanes/2-propanol, 0.75 mL/min, 254 nm, $R_{t1} = 14.76$ min, $R_{t2} = 17.50$ min. 68% ee, $([\alpha]_D^{20} = -32, c 0.11$ in CHCl_3).

(E)-2-Methyl-4-phenylpent-3-enitrile (2):

Enantiomer Separation: Chiralpak AS-H Column 95:5 hexanes/2-propanol, 0.75 mL/min, 254 nm, $R_{t1} = 12.81$ min, $R_{t2} = 13.69$ min.

2,6-Dimethyl-2-propenyl-hept-5-enitrile (6):

Enantiomer Separation: Cyclodex-B Column, 5 min at 60°C, 0.7°C/min, 120 min at 160°C.

Determination of the Absolute Configuration of (S)-(-)-(E)-2-Methyl-4-phenylbut-3-enitrile (1): To a toluene solution of (E)-2-methyl-4-phenylbut-3-enitrile (78% ee, from ligand **L2**) (30 mg, 0.19 mmol) was added 1.0 M DIBAL-H in hexane (0.2 mL, 0.2 mmol) dropwise during about 5 minutes at -78°C, and the reaction mixture was stirred at that

temperature for 1.5 h and quenched with 1M NH₄Cl (0.25 mL), the cold bath was removed and diethyl ether (8 mL) was added and warm to room temperature. The organic layer was separated and washed with 1M citric acid (2 X 2.5 mL), water (1 X 3 mL) and dried over Na₂SO₄ and solvents were removed and the yellow residue was dried under vacuum.

The crude yellow mixture was dissolved into 0.75 mL of MeOH and NaBH₄ (0.76 mmol) was added portion-wise at room temperature. The reaction mixture was stirred for 30 minutes at the same temperature and quenched with water (0.2 mL), MeOH was removed and diethyl ether (4 mL) was added and decanted; dried over Na₂SO₄ and filtered; solvent was evaporated and the crude residue was purified by short silica gel column chromatography using hexanes/Et₂O (9 : 1 to 4 : 1) as eluant to obtain (*E*)-2-methyl-4-phenylbut-3-en-1-ol in 57% yield. As can be seen from the data below some racemization ensued during the aldehyde formation and subsequent reduction.

Enantiomer separation: Chiralcel OD Column 95:5 hexanes/2-propanol, 1 mL/min, 254 nm, R_{t1} = 9.46 min, R_{t2} = 11.32 min. 61% ee, ([α]_D²⁰ = – 32, *c* 0.16 in CHCl₃), [lit.⁷: [α]_D²⁰ = – 47, *c* 0.8 in CHCl₃, 100% ee].

Determination of the Absolute Configuration of (*S*)-2-(3,4-Dihydronaphthalene)-propionitrile (4): To a solution of 2-(3,4-dihydronaphthalene)-propionitrile (75% ee from **L2**, 15 mg, 0.08 mmol) in benzene (1.5 mL) was added DDQ (55 mg, 0.24 mmol) at room temperature under nitrogen. After the mixture was stirred at room temperature for 27 h, the solvent was removed on rotary evaporator and the resulting residue was taken into 1 mL hexane/Et₂O mixture (1:1) and was filtered through short pad of celite, solvents were removed under vacuo and the crude mixture was purified using short silica gel column chromatography with 19:1 hexane/Et₂O mixture to afford 2-(1-naphthalenyl)-propionitrile, the spectral data is in agreement with the known sample.¹ The absolute configuration *S* was assigned by comparing

with the elution order of HPLC analysis of the reported compound.¹ Daicel Chiralcel OJ Column 95:5 hexanes/2-propanol, 0.5 mL/min, 254 nm, *R* isomer: $R_{t1} = 29.8$ min, *S*-isomer: $R_{t2} = 32.15$ min. (74 % ee).

References

1. Casalnuovo, A. L.; RajanBabu, T. V.; Ayeres, T. A.; Warren, T. H. *J. Am. Chem. Soc.* **1994**, *116*, 9869.
2. Stille, J. K.; Groh, B. L. *J. Am. Chem. Soc.* **1987**, *109*, 813.
3. Pal, K. *Synthesis* **1995**, 1485.
4. He, Z.; Yi, C. S.; Donaldson, W. A. *Org. Lett.* **2003**, *5*, 1567.
5. Rodriguez, J.; Waegell, B. *Synthesis*, **1988**, 535.
6. Ken-Tsung W.; Ying-Yueh, H. *Tetrahedron Lett.* **2003**, *44*, 8033.
7. Ohfuné, Y.; Tomita, M. *J. Am. Chem. Soc.* **1982**, *104*, 3511.

Table 1. Diarylphosphinites for Hydrocyanation of 1-Phenylbuta-1,3-diene

Ligands L	conv. (%)	%ee	cat. (mol%)	time(h)
	29	28	2	72
	0	--	--	--
<p>Ar = C_6F_5</p>	21	27	2	72
<p>Ar = </p>	12	7.5	2	72
	56	71	2	72
<p>Tr = Ph_3C</p>	62	27	2	72

1.541
1.526

3.538
3.523
3.511
3.509

7.416
7.402
7.383
7.378
7.369
7.353
7.324
7.310
7.296
6.766
6.734
6.119
6.106
6.087
6.075

Compound # 1

Current Data Parameters
NAME: 14-05-273
EXPNO: 1
PROCNO: 1
F2 - Acquisition Parameters
Date_: 20060615
Time: 18.09
INSTRUM: spect
PROBHD: 5 mm Multinucl
PULPROG: zg30
TD: 65536
SOLVENT: CDCl3
NS: 16
DS: 2
SWH: 10310.572 Hz
FIDRES: 0.157632 Hz
AQ: 3.171923 sec
RG: 327.5
DM: 48.400 usec
DE: 6.00 usec
TE: 300.2 K
DWT: 1.000000 sec
MCREST: 0.0000000 sec
MCWID: 0.01500000 sec
===== CHANNEL f1 =====
NUC1: 1H
P1: 14.80 usec
PL1: 0.00 dB
SFO1: 500.023978 MHz
F2 - Processing parameters
SI: 32768
SF: 500.020000 MHz
WDW: EM
SSB: 0
LB: 0.30 Hz
GB: 0
PC: 1.00

76.86
77.12
79.37

130.90
126.98
126.97
126.29
126.24
132.95
135.74

Compound # 1

Current Data Parameters
NAME IV-B5-293-C13
EXPNO 1
PROCNO 1
F2 - Acquisition Parameters
Date_ 20080814
Time 18.14
INSTRUM spect
PROBHD 5 mm MUX-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
DS 4
SS 4
SMR 30010.029 Hz
FIDRES 0.01324 Hz
AQ 1.01324 sec
RG 1635.5
EW 16.650 usec
TE 300.2 K
D1 2.0000000 sec
d11 0.0000000 sec
d12 0.0000000 sec
d13 0.0000000 sec
DELTA 0.0000000 sec
MCHEST 0.0000000 sec
MCWPR 0.0150000 sec
===== CHANNEL f1 =====
NUC1 13C
P1 13.00 usec
PL1 0.00 dB
SFO1 125.762700 MHz
===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 0.00 dB
PL12 18.80 dB
PL13 120.00 dB
SFO2 500.0220001 MHz
F2 - Processing parameters
SI 32768
SF 500.136460 MHz
WDW EM
SSB 0
GB 1.00 Hz
PC 1.40

2.138
2.135
1.558
1.519
1.501

3.663
3.645
3.642
3.624

7.408
7.403
7.398
7.387
7.382
7.380
7.363
7.358
7.348
7.343
7.333
7.328
7.323
7.319
7.311
7.295
7.281
5.672
5.665
5.651
5.647

Compound # 2

Current Data Parameters
NAME IV-BS-159
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters:
Date_ 20060301
Time_ 12.40
INSTRUM spect
PROBHD 5 mm Multinucl
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 4
DW 60.400 use
DE 6.00 use
TE 300.2 K
D1 1.00000000 sec
MCREST 0.00000000 sec
MCWRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.50 use
PL1 -6.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 32768
SF 400.1300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Current Data Parameters
NAME IV-85-297-C13
EXPNO 1
PROCNO 1
F2 - Acquisition Parameters
Date_ 20060623
Time_ 10.37
INSTRUM spect
PROBHD 5 mm Multinucl
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 274
DS 4
SWH 30030.075 Hz
FIDRES 0.458222 Hz
AQ 1.0912244 sec
RG 32768
RW 16.650 usec
TE 300.2 K
D1 2.00000000 sec
d11 0.03000000 sec
DELTA 1.89999998 sec
ACQRES 0.00000000 sec
ACQRES 0.01300000 sec
===== CHANNEL f1 =====
NUC1 13C
P1 13.00 usec
PL1 0.00 dB
SFO1 125.7427020 MHz
===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.40 usec
PL2 -1.00 dB
PL12 18.80 dB
PL13 120.00 dB
SFO2 500.0220001 MHz
F2 - Processing Parameters
SI 32768
SF 125.7301290 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Compound # 3

Current Data Parameters
 NAME IV-BS-302
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20060620
 Time_ 12.08
 INSTRUM spect
 PROBHD 5 mm Multinucl
 PULPROG zg30
 TD 65536
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 8278.146 Hz
 FIDRES 0.126314 Hz
 AQ 3.9584243 sec
 RG 181
 DW 60.400 usec
 DE 6.00 usec
 TE 297.2 K
 D1 1.00000000 sec
 MCREST 0.00000000 sec
 MCWRK 0.01500000 sec
 ===== CHANNEL f1 =====
 NUC1 1H
 P1 7.50 usec
 PL1 -6.00 dB
 SFO1 400.1324710 MHz
 F2 - Processing parameters
 SI 32768
 SF 400.1300097 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

26.49
26.26
24.17
19.91

77.30
77.05
76.79

131.84
128.33
121.45
120.75

```

Current Data Parameters
NAME      V-BE-7
EXPNO     2
PROCNO    1

F2 - Acquisition Parameters
Date_     20060629
Time      19.44
INSTRUM   spect
PROBHD    5 mm Multinuc
PULPROG   zgpg30
TD         65536
SOLVENT   CDCl3
NS         3128
DS         4
SWH        30030.049 Hz
FIDRES     0.458222 Hz
AQ         1.0912244 sec
RG          2288
DM          16680 usec
DE          6.00 usec
TE         300.2 K
D1         2.00000000 sec
DELTA      0.03000000 sec
DELTAT     0.00000000 sec
MCREST     0.00000000 sec
MCWRRK     0.01500000 sec

===== CHANNEL f1 =====
NUC1       13C
P1         13.00 usec
PL1        3.00 dB
SFO1       125.7627020 MHz

===== CHANNEL f2 =====
CPDPRG2    waltz16
NUC2       1H
P2         100.00 usec
PL2        19.00 dB
SFO2       500.0220001 MHz
PL12       18.80 dB
PL13       120.00 dB
SFO2       500.0220001 MHz

F2 - Processing parameters
SI         32768
SF         125.7301290 MHz
WDW        EM
SSB        0
GB         1.0 Hz
PC         1.40
  
```

3.933
3.931
3.919
3.917
2.809
2.793
2.777
2.389
2.379
2.377
2.373
2.363
2.361
2.358
2.356
2.353
2.346
1.607
1.592

7.283
7.267
7.263
7.260
7.253
7.247
7.237
7.223
7.219
7.213
7.204
6.348
6.356
6.365

Compound # 4

Current Data Parameters
NAME IV-BS-295
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060615
Time 18.26
INSTRUM spect
PROBHD 5 mm Multinucl
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 10330.578 Hz
FIDRES 0.157632 Hz
AQ 3.1719923 sec
RG 4
RW 4.000000 sec
DE 6.000000 sec
TE 300.2 K
D1 1.00000000 sec
NCREST 0.00000000 sec
NCRMK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 14.80 usec
PL1 -1.00 dB
SFO1 500.0230878 MHz

F2 - Processing parameters
SI 32768
SF 500.0200000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

27.99
27.98
23.96
18.88

77.32
77.07
76.91

136.86
132.74
131.83
129.19
127.61
127.23
126.63
121.96
121.66

Compound # 4

Current Data Parameters
NAME IV-BS-295-C13
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050615
Time 18.30
INSTRUM spect
PROBHD 5 mm Multinucl
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 173
DS 4
SWH 30030.025 Hz
FIDRES 1.01322 Hz
AQ 1.092244 sec
RG 32768
DM 16.650 usec
DE 6.00 usec
TE 300.2 K
D1 2.00000000 sec
d11 0.03000000 sec
DELTA 1.89999998 sec
MCREST 0.00000000 sec
MCWRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 13.00 usec
PL1 3.00 dB
SFO1 125.7627020 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 -1.00 dB
PL12 18.00 dB
PL13 12.00 dB
SFO2 500.0220001 MHz

F2 - Processing parameters
SI 32768
SF 125.7601250 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1.733
1.726
1.715
1.707

3.956
3.942
3.938
3.935
3.440

7.535
7.517
7.461
7.442
7.387
7.368
7.350
7.311
7.292
7.282
7.276
7.274
6.603

Current Data Parameters
NAME IV-BS-60
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20051108
Time 21.20
INSTRUM spect
PROBHD 5 mm Multinucl
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 16
DS 2
SWH 8278.146 Hz
FIDRES 0.252629 Hz
AQ 1.9792372 sec
RG 114
DM 60.400 usec
DE 6.00 usec
TE 300.2 K
D1 1.00000000 sec
MCREST 0.00000000 sec
MCWRK 0.01500000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.50 usec
PL1 -6.00 dB
SFO1 400.1324710 MHz

F2 - Processing parameters
SI 16384
SF 400.1300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

Compound # 5

18.14
24.86
37.77

77.33
77.21
77.01
76.69

144.43
141.95
139.54
139.63
126.39
125.49
124.26
120.90
118.94

Current Data Parameters
NAME BS-Cyano
EXPNO 2
PROCNO 1
F2 - Acquisition Parameters
Date_ 20060629
Time_ 20.11
INSTRUM spect
PROBHD 5 mm Multinucl
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1024
DS 4
SWH 23980.814 Hz
FIDRES 0.365518 Hz
AQ 1.3664756 sec
RG 32768
DM 20.850 usec
DE 6.00 usec
TE 300.2 K
D1 2.00000000 sec
d11 0.03000000 sec
DELTA 1.89999998 sec
MCREST 0.00000000 sec
MCWRR 0.01500000 sec
===== CHANNEL f1 =====
NUC1 13C
P1 5.90 usec
PL1 -6.00 dB
SFO1 100.628298 MHz
===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 -2.00 dB
PL12 15.80 dB
PL13 120.00 dB
SFO2 400.1316005 MHz
F2 - Processing parameters
SF 100.628298 MHz
WDW EM
SSB 0
LB 0
GB 0
PC 1.40

Compound # 5

210 200 190 180 170 160 150 140 130 120 110 100 90 80 70 60 50 40 30 20 10 0 ppm

Current Data Parameters
 NAME IV-B8-234-Pdt
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20060428
 Time_ 11.40
 INSTRUM spect
 PROBRD 5 mm Multinuc
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 8978.14 Hz
 FIDRES 0.136314 Hz
 AQ 3.9584243 sec
 RG 256
 DW 60.400 usec
 DE 6.00 usec
 TE 300.2 K
 D1 1.00000000 sec
 D2 0.00000000 sec
 MCOREST 0.00000000 sec
 MCNFK 0.01500000 sec
 ===== CHANNEL f1 =====
 NUC1 13C
 P1 14.00 usec
 PL1 -6.00 dB
 SF01 400.1324710 MHz
 F2 - Processing parameters
 SI 32768
 SF 400.1300161 MHz
 MDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

Compound # 6

17.55
17.59
14.04
13.92

40.40
39.43

77.30
77.04
76.79

132.79
131.13
126.65
122.63

Compound # 6

Current Data Parameters
NAME IN-198
EXPNO 1
PROCNO 1
F2 - Acquisition Parameters
Date_ 20060623
Time 11:12
INSTRUM spect
PROBHD 5 mm Multinuc1
PULPROG zgpg30
SOLVENT CDCl3
NS 87
DS 4
SWH 30030.098 Hz
FIDRES 0.458222 Hz
AQ 1.0912244 sec
RG 655
AQ 16.680 usec
DE 6.00 usec
TE 296.2 K
D1 2.000000 sec
d11 0.0300000 sec
DELTA 1.8999998 sec
PCYCL 0.0000000 sec
NOVER 0.0150000 sec
===== CHANNEL f1 =====
NUC1 13C
P1 13.00 usec
PL1 3.00 dB
SFO1 125.747040 MHz
===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
P2 100.00 usec
PL2 -1.00 dB
PL3 18.00 dB
PL13 12.00 dB
SFO2 500.0220001 MHz
F2 - Processing parameters
SI 32768
SF 125.7301280 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

Current Data Parameters
 NAME IV-SB-294
 EXPNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20060622
 Time 17.16
 INSTRUM spect
 PROBHD 5 mm Multinucl
 PULPROG zgpg30
 SOLVENT CDCl3
 NS 99
 DS 4
 SWH 23980.814 Hz
 FIDRES 0.0000000 Hz
 AQ 1.3667156 sec
 RG 1448.2
 DW 20.850 usec
 DE 6.00 usec
 TE 300.2 K
 D1 2.0000000 sec
 d11 0.03000000 sec
 DELTA 1.89999998 sec
 MCREST 0.00000000 sec
 MCWRR 0.01500000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 5.90 usec
 PL1 -6.00 dB
 SFO1 100.628298 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 FCPD2 80.00 usec
 PL2 18.00 dB
 PL12 18.00 dB
 PL13 120.00 dB
 SFO2 400.1316005 MHz

F2 - Processing parameters
 SI 32768
 SF 100.612760 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 FC 1.40

Compound # 7 and 8

* ATT 2^3.0
 *RN
 RUN # 2791 MAY 30, 2006 11:43:38
 START
 TF
 4F 239
 7 624
 5 473

Compound # 3

FNN OF STANAI

Cyclodex B, 35 min at 35°C

Closing signal file M:STANAI RNR

RUN# 2791 MAY 30, 2006 11:43:38

STANAI FTIF: M:STANAI RNR

RT	ARFA	TYPF	WINTH	ARFA2
4 239	1571	AR	020	49295
5 473	31333	PA	060	98 50547
6 220	1950	AR	035	61187
7 624	1242	AR	095	38971

TOTAL ARFA= 318696
 MIN FACTOR=1 0000F+00

V-33

c:\users\IV-33, Channel A

* RUN # 1018 FEB 19, 1992 07:36:33
START

Dehydrogenation product of compound # 4

Closing signal file M:SIGNAL.BNC

RUN# 1018 FEB 19, 1992 07:36:33

SIGNAL FILE: M:SIGNAL.BNC

AREA%	RT	AREA	TYPE	WIDTH	AREA%
5.795	1716181	BB	.356	5.78378	
24.480	4946720	VV	1.053	16.67118	
26.770	21294688	VV	1.298	71.76624	
33.978	1714705	VV	1.736	5.77881	

TOTAL AREA=2.9672E+07
MUL FACTOR=1.0000E+00