Supporting Information

A novel alternative for the *N*–*S* bond formation and its application to the synthesis of benzisothiazol-3-ones

Arkaitz Correa, Imanol Tellitu, Esther Domínguez,* and Raul SanMartin*

Departamento de Química Orgánica II, Facultad de Ciencia y Tecnología,

Universidad del País Vasco – Euskal Herriko Unibertsitatea

P. O. Box 644, 48080 Bilbao, Spain

imanol.tellitu@ehu.es

Supporting Information Available. Experimental details for compounds **2a–j** and **1a–j** and ¹H NMR and ¹³C NMR spectra of all new compounds are included.

General remarks	S1
Typical reaction procedure for amides 2a–j	S2
Spectroscopic data of amides 2a–j	S3
Typical reaction procedure for benzisothiazolones 1a–j	S7
Spectroscopic data of benzisothiazolones 1a–j	S8
¹H NMR and ¹³C NMR spectra	S12

General Methods and Materials. All reagents were purchased and used as received. Melting points were measured using open glass capillaries and are uncorrected. Infrared spectra were recorded as KBr plates or as thin films and peaks are reported in cm^{–1}. Only representative absorptions are given. Flash chromatography was carried out on

SiO₂ (silica gel 60, 230–400 mesh ASTM). NMR spectra were recorded on a 300 instrument (300 MHz for ¹H and 75.4 MHz for ¹³C) at 20 °C. Chemical shifts (τ^m) were measured in ppm relative to chloroform (τ^m=7.26 for ¹H or 77.00 for ¹³C) as internal standard. Coupling constants, *J*, are reported in Hertz. DEPT experiments were used to assist with the assignment of the signals. HRMS spectra were recorded on a Waters GCT Mass Spectrometer.

Typical reaction procedure for the synthesis of 2-mercaptoamides 2a–j

2-Mercapto-*N*-(*p*-methoxyphenyl)benzamide (2a**).¹ Typical procedure:**

A solution of $AlMe_3$ (11.52 mmol, 2.0 M in toluene) was added dropwise to a cooled (0 °C) suspension of *p*-anisidine (1.42 g, 11.52 mmol) in CH_2Cl_2 (30 mL). When the addition was complete, the reaction mixture was allowed to warm to room temperature and stirring was continued for 30 minutes until the gas evolution ceased. Then, a solution of commercially available methyl thiosalicylate (**3**) (0.70 mL, 5.76 mmol) was added and the mixture was heated under reflux overnight. The reaction mixture was cooled to room temperature and carefully quenched with 5% aq HCl (20 mL). The organic layer was separated and the aqueous layer was extracted with CH_2Cl_2 (3x15 mL). The combined organic extracts were washed with a saturated aqueous solution of $NaHCO_3$ (15 mL) and brine (15 mL). Then, the organic layer was dried over sodium sulfate, filtered and the solvent was evaporated *in vacuo*. The resulting residue was purified by crystallization from Et_2O to afford amide **2a** as a white solid (92% yield). The ¹H and ¹³C NMR spectra of **2a** were in total agreement with the reported data: mp 140–141 °C (Et_2O). Lit.¹ 149–151 °C.

¹ Waisser, K.; Pesina, M.; Holý, P.; Pour, M.; Bures, O.; Kunes, J.; Klimesova, V.; Buchta, V.; Kubanová, P.; Kaustová, J. *Arch. Pharm. Pharm. Med. Chem.* **2003**, 336, 322.

2-Mercaptobenzanilide (**2b**).¹

According to the general procedure using aniline (0.53 mL, 5.76 mmol), benzanilide **2b** was obtained as a white solid in 95% yield after purification by crystallization from Et₂O. The ¹H and ¹³C NMR spectra of **2b** were in total agreement with the reported data: mp 117–118 °C (Et₂O). Lit.¹ 109–110 °C.

N-(*p*-Ethylphenyl)-2-mercaptobenzamide (**2c**).

According to the general procedure using *p*-ethylaniline (0.71 mL, 5.76 mmol), benzamide **2c** was obtained as a white solid in 95% yield after purification by crystallization from Et₂O: mp 217–218 °C (Et₂O); ¹H NMR (CDCl₃) _ 1.26 (t, *J*=7.5, 3H, CH₃), 2.66 (q, *J*=7.5, 2H, CH₂), 4.62 (br s, 1H, SH), 7.18–7.39 (m, 5H, H_{arom}), 7.53–7.60 (m, 3H, H_{arom}), 7.81 (br s, 1H, NH); ¹³C NMR (DMSO-*d*₆) _ 15.7 (CH₃), 27.7 (CH₂), 120.2 (CH), 124.6 (CH), 126.2 (CH), 127.8 (CH), 130.3 (CH), 131.2 (CH), 133.3 (C), 134.1 (C), 136.8 (C), 139.4 (C), 165.8 (C); IR (KBr) 3296, 1637; MS (EI) *m/z* (%) 257 (M⁺, 12), 137 (42), 121 (95), 106 (100), 91 (21); HRMS calc. for C₁₅H₁₅NOS 257.0874, found 257.0874.

2-Mercapto-*N*-(1-naphthyl)benzamide (**2d**).

According to the general procedure 1-naphthylamine (825 mg, 5.76 mmol), benzamide **2d** was obtained as a white solid in 95% yield after purification by crystallization from Et₂O: mp 196–197 °C (Et₂O); ¹H NMR (DMSO-*d*₆) _ 5.31 (br s, 1H, SH), 7.28–7.42 (m, 7H, H_{arom}), 7.54–8.12 (m, 4H, H_{arom}), 10.52 (br s, 1H, NH); ¹³C NMR (DMSO-*d*₆) _ 123.4 (CH), 123.5 (CH), 124.8 (CH), 125.6 (CH), 126.0 (CH), 126.2 (CH), 126.3 (CH),

128.2 (CH), 128.7 (CH), 128.9 (CH), 130.7 (CH), 133.6 (C), 133.7 (C), 133.9 (C), 167.5 (C); IR (KBr) 3249, 1631; MS (EI) m/z (%) 279 (M^+ , 11), 143 (100), 137 (30), 115 (57), 109 (34); HRMS calc. for $C_{17}H_{13}NOS$ 279.0718, found 279.0721.

***N*-(2-Iodophenyl)-2-mercaptobenzamide (2e).**

According to the general procedure using 2-iodoaniline (1.29 g, 5.76 mmol), benzamide **2e** was obtained as a white solid in 95% yield after purification by crystallization from Et_2O : mp 147–148 °C (Et_2O); 1H NMR ($CDCl_3$) δ 4.67 (br s, 1H, SH), 6.91–6.96 (m, 1H, H_{arom}), 7.26–7.75 (m, 4H, H_{arom}), 7.83–7.87 (m, 2H, H_{arom}), 8.10 (br s, 1H, NH), 8.43 (d, $J=7.5$, 1H, H_{arom}); ^{13}C NMR ($DMSO-d_6$) δ 98.6 (C), 124.7 (CH), 128.5 (CH), 128.6 (CH), 128.8 (CH), 128.9 (CH), 130.8 (CH), 130.9 (CH), 132.4 (C), 134.4 (C), 138.9 (CH), 139.5 (C), 166.5 (C); IR (KBr) 3249, 1631; MS (EI) m/z (%) 354 (M^+ , 1), 228 (48), 137 (27), 109 (37), 84 (66), 82 (100); HRMS calc. for $C_{13}H_{10}INOS$ 354.9528, found 354.9510.

***N*-(2-Bromo-4-methylphenyl)-2-mercaptobenzamide (2f).**

According to the general procedure using 2-bromo-4-methylaniline (0.73 mL, 5.76 mmol), benzamide **2f** was obtained as a white solid in 91% yield after purification by crystallization from Et_2O : mp 135–136 °C (Et_2O); 1H NMR ($CDCl_3$) δ 2.33 (s, 3H, CH_3), 4.64 (br s, 1H, SH), 7.16–7.40 (m, 5H, H_{arom}), 7.65 (d, $J=7.8$, 1H, H_{arom}), 8.17 (br s, 1H, NH), 8.33 (d, $J=8.3$, 1H, H_{arom}); ^{13}C NMR ($DMSO-d_6$) δ 20.2 (CH_3), 120.3 (C), 124.4 (CH), 128.1 (CH), 128.4 (CH), 128.5 (CH), 130.4 (CH), 130.5 (CH), 132.4 (C), 132.6 (CH), 133.6 (C), 134.2 (C), 137.9 (C), 166.6 (C); IR (KBr) 3402, 1660; MS (EI) m/z (%) 322 (M^++2 , 4), 320 (M^+ , 3), 242 (100), 184 (65), 109 (59); HRMS calc. for $C_{14}H_{12}BrNOS$ 320.9823, found 320.9825.

2-Mercapto-*N*-(6-methoxy-3-pyridyl)benzamide (2g).

According to the general procedure using 3-amino-6-methoxypyridine (578 mg, 4.66 mmol), benzamide **2g** was obtained as a white solid in 61% yield after purification by crystallization from Et₂O: mp 135–136 °C (Et₂O); ¹H NMR (DMSO-*d*₆) δ 3.02 (br s, 1H, SH), 3.85 (s, 3H, OCH₃), 6.86 (d, *J*=8.9, 1H, H_{arom}), 7.39 (t, *J*=7.4, 1H, H_{arom}), 7.52 (t, *J*=7.5, 1H, H_{arom}), 7.72–7.83 (m, 2H, H_{arom}), 8.05 (dd, *J*=8.7, 1.6, 1H, H_{arom}), 8.50 (s, 1H, H_{arom}), 10.6 (br s, 1H, NH); ¹³C NMR (DMSO-*d*₆) δ 53.3 (CH₃), 110.2 (CH), 126.3 (CH), 126.4 (CH), 128.6 (C), 129.8 (CH), 131.7 (CH), 132.4 (CH), 134.1 (C), 136.7 (C), 138.7 (CH), 160.1 (C), 165.7 (C); IR (KBr) 3272, 1643; MS (EI) *m/z* (%) 260 (M⁺, 35), 258 (100), 229 (81), 137 (91), 109 (84), 95 (37), 80 (77); HRMS calc. for C₁₃H₁₂N₂O₂S 260.0619, found 260.0618.

***N*-Benzyl-2-mercaptobenzamide (2h).²**

According to the general procedure using benzylamine (0.40 mL, 5.76 mmol), benzamide **2h** was obtained as a white solid in 60% yield after purification by crystallization from Et₂O. The ¹H and ¹³C NMR spectra of **2h** were in total agreement with the reported data: mp 108–109 °C (Et₂O). Lit.² 108 °C.

***N*-Allyl-2-mercaptobenzamide (2i).**

According to the general procedure using allylamine (0.43 mL, 5.76 mmol), benzamide **2i** was obtained as a colourless oil in 63% yield after purification by column

² El-Barbary, A. A.; Clausen, K.; Lawesson, S. –O. *Tetrahedron* **1980**, 36, 3309.

chromatography (CH₂Cl₂): mp 135–136 °C (Et₂O); ¹H NMR (CDCl₃) _ 3.91–3.96 (m, 2H, CH₂CH), 4.66 (br s, 1H, SH), 5.07–5.22 (m, 2H, CH=CH₂), 5.77–5.90 (m, 1H, CH=CH₂), 6.64 (br s, 1H, NH), 7.00–7.25 (m, 3H, H_{arom}), 7.39 (d, *J*=7.6, 1H, H_{arom}); ¹³C NMR (CDCl₃) _ 42.1 (CH₂), 116.4 (CH₂), 124.1 (CH), 127.8 (CH), 130.4 (CH), 130.7 (CH), 133.6 (CH), 132.7 (C), 132.8 (C), 168.4 (C); IR (KBr) 3359, 1641; MS (EI) *m/z* (%) 193 (M⁺, 50), 164 (42), 151 (34), 135 (100), 108 (81); HRMS calc. for C₁₀H₁₁NOS 193.0561, found 193.0560.

2-Mercapto-*N*-methylbenzamide (2j).

According to the general procedure using NH₂Me·HCl (401 mg, 5.94 mmol), benzamide **2j** was obtained as a white solid in 76% yield after purification by crystallization from Et₂O: mp 90–91 °C (Et₂O). Lit.³ 90–92 °C; ¹H NMR (CDCl₃) _ 2.89 (s, 3H, CH₃), 4.68 (br s, 1H, SH), 6.41 (br s, 1H, NH), 6.51–7.07 (m, 1H, H_{arom}), 7.19–7.26 m, 2H, H_{arom}), 7.35–7.38 (m, 1H, H_{arom}); ¹³C NMR (CDCl₃) _ 26.2 (CH₃), 124.5 (CH), 127.7 (CH), 129.9 (CH), 130.2 (CH), 132.4 (C), 132.5 (C), 169.1 (C); IR (KBr) 3368, 1653; MS (EI) *m/z* (%) 167 (M⁺, 14), 166 (100), 137 (51), 109 (33); HRMS calc. for C₈H₉NOS 167.0405, found 167.0406.

³ Nagy, P.; Csámpai, A.; Szabó, D.; Varga, J.; Ramat, V.; Ruff, F.; Kucsman, A. *J. Chem. Soc., Perkin Trans. 2* **2001**, 339.

Typical reaction procedure for the synthesis of benzisothiazol-3-ones **1a–j.**

2-(*p*-Methoxyphenyl)-1,2-benzisothiazol-3(2*H*)-one (1a**).⁴ Typical procedure:**

A solution of PIFA (249 mg, 0.58 mmol) in 9 mL of CH₂Cl₂ was added at 0 °C to a solution of benzamide **2a** (100 mg, 0.39 mmol) and TFA (0.09 mL, 1.16 mmol) in 6 mL of the same solvent, and the new solution was stirred for 1 hour. Then, the solvent was evaporated at reduced pressure and the resulting residue was purified by column chromatography (CH₂Cl₂) followed by crystallization from EtOH to afford benzisothiazolone **1a** as a white solid (78%). The ¹H and ¹³C NMR spectra of **1a** were in total agreement with the reported data: mp 145–146 °C (EtOH). Lit.⁴ 147–148 (EtOH).

2-Phenyl-1,2-benzisothiazol-3(2*H*)-one (1b**).⁵**

According to the general procedure, benzisothiazolone **1b** was obtained from benzanilide **2b** (100 mg, 0.44 mmol) as a white solid in 71% yield after purification by column chromatography (CH₂Cl₂) followed by crystallization from EtOH. The ¹H and ¹³C NMR spectra of **1b** were in total agreement with the reported data: mp 139–140 °C (EtOH). Lit.⁵ 138–140 (EtOH).

⁴ Sano, T.; Takagi, T.; Gama, Y.; Shibuya, I.; Shimizu, M. *Synthesis* **2004**, 1585.

⁵ Shimizu, M.; Sugano, Y.; Konakahara, T.; Gama, Y.; Shibuya, I. *Tetrahedron* **2002**, 58, 3779.

2-(*p*-Ethylphenyl)-1,2-benzisothiazol-3(2*H*)-one (1c).

According to the general procedure benzisothiazolone **1c** was obtained from benzamide **2c** (100 mg, 0.39 mmol) as a white solid in 66% yield after purification by column chromatography (hexanes/EtOAc, 1:1) followed by crystallization from EtOH: mp 122–123 °C (EtOH); ¹H NMR (CDCl₃) _ 1.24 (t, *J*=7.6, 3H, CH₃), 2.67 (q, *J*=7.6, 2H, CH₂), 7.27–7.44 (m, 3H, H_{arom}), 7.54–7.63 (m, 4H, H_{arom}), 8.07–8.10 (m, 1H, H_{arom}); ¹³C NMR (CDCl₃) _ 15.4 (CH₃), 28.4 (CH₂), 119.9 (CH), 124.6 (CH), 125.5 (CH), 126.9 (CH), 128.6 (CH), 132.0 (CH), 134.6 (C), 139.8 (C), 139.9 (C), 143.3 (C), 164.0 (C); IR (KBr) 1649; MS (EI) *m/z* (%) 255 (M⁺, 45), 240 (100), 212 (20); HRMS calc. for C₁₅H₁₃NOS 255.0718, found 255.0719.

2-(1-Naphthyl)-1,2-benzisothiazol-3(2*H*)-one (1d).⁶

According to the general procedure benzisothiazolone **1d** was obtained from benzamide **2d** (100 mg, 0.36 mmol) as a white solid in 62% yield after purification by column chromatography (hexanes/EtOAc, 1:1) followed by crystallization from EtOH: mp 140–150 °C (EtOH). Lit.⁶ 154–155 °C (CCl₄/CH₂Cl₂); ¹H NMR (CDCl₃) _ 7.23 (s, 1H, H_{arom}), 7.48–7.73 (m, 7H, H_{arom}), 7.97 (d, *J*=8.4, 2H, H_{arom}), 8.18 (d, *J*=7.9, 1H, H_{arom}); ¹³C NMR (DMSO-*d*₆) _ 122.1 (CH), 122.8 (CH), 123.4 (C), 125.8 (CH), 126.1 (CH), 126.4 (CH), 126.9 (CH), 127.5 (CH), 127.9 (CH), 128.6 (CH), 129.8 (CH), 131.6 (C), 132.6 (CH), 134.4 (C), 136.1 (C), 141.7 (C), 165.7 (C); IR (KBr) 1655; MS (EI) *m/z* (%) 277 (M⁺, 100), 260 (42), 140 (40), 115 (39); HRMS calc. for C₁₇H₁₁NOS 277.0561, found 277.0564.

⁶ Uchida, Y.; Kozuka, S. *Bull. Chem. Soc. Jpn.* **1982**, *55*, 1183.

2-(2-Iodophenyl)-1,2-benzisothiazol-3(2H)-one (1e).

According to the general procedure, benzisothiazolone **1e** was obtained from benzamide **2e** (100 mg, 0.28 mmol) as a white solid in 64% yield after purification by column chromatography (hexanes/EtOAc, 1:1) followed by crystallization from EtOH: mp 125–127 °C (EtOH); ^1H NMR (CDCl_3) δ 7.13–7.20 (m, 1H, H_{arom}), 7.43–7.48 (m, 3H, H_{arom}), 7.58–7.70 (m, 2H, H_{arom}), 7.97 (d, $J=7.9$, 1H, H_{arom}), 8.13 (d, $J=7.9$, 1H, H_{arom}); ^{13}C NMR (CDCl_3) δ 99.4 (C), 120.4 (CH), 123.5 (C), 125.8 (CH), 127.4 (CH), 129.5 (CH), 130.7 (CH), 131.1 (CH), 132.5 (CH), 138.7 (C), 140.1 (CH), 141.0 (C), 164.0 (C); IR (KBr) 1661; MS (EI) m/z (%) 353 (M^+ , 24), 227 (100), 198 (69), 171 (34), 154 (49), 127 (40); HRMS calc. for $\text{C}_{13}\text{H}_8\text{INOS}$ 352.9371, found 352.9372.

2-(2-Bromo-4-methylphenyl)-1,2-benzisothiazol-3(2H)-one (1f).⁷

According to the general procedure, benzisothiazolone **1f** was obtained from benzamide **2f** (100 mg, 0.31 mmol) as a white solid in 67% yield after purification by column chromatography (CH_2Cl_2) followed by crystallization from hexanes: mp 125–127 °C (hexanes). Lit.⁷ 139 °C (petroleum ether/ CH_2Cl_2); ^1H NMR (CDCl_3) δ 2.36 (s, 3H, CH_3), 7.18–7.20 (m, 1H, H_{arom}), 7.32–7.44 (m, 2H, H_{arom}), 7.52–7.66 (m, 3H, H_{arom}), 8.10 (d, $J=7.9$, 1H, H_{arom}); ^{13}C NMR (CDCl_3) δ 20.8 (CH_3), 120.1 (CH), 123.2 (C), 123.5 (C), 125.4 (CH), 127.0 (CH), 129.0 (CH), 130.5 (CH), 132.1 (CH), 132.3 (C), 133.9 (CH), 141.1 (C), 141.5 (C), 164.0 (C); IR (KBr) 1670; MS (EI) m/z (%) 320 ($\text{M}^+ + 2$, 95), 318 (M^+ , 96), 240 (100), 212 (80), 136 (54), 108 (85); HRMS calc. for $\text{C}_{14}\text{H}_{10}\text{BrNOS}$ 318.9666, found 318.9665.

⁷ Hellwinkel, D.; Lenz, R.; Lämmerzahl, F. *Tetrahedron* **1983**, 39, 2073.

2-(6-Methoxy-3-pyridyl)-1,2-benzisothiazol-3(2H)-one (1g).

According to the general procedure, benzisothiazolone **1g** was obtained from benzamide **2g** (100 mg, 0.38 mmol) as a white solid in 70% yield after purification by column chromatography (hexanes/EtOAc, 1:1) followed by crystallization from EtOH: mp 125–126 °C (EtOH); ^1H NMR (CDCl_3) δ 3.96 (s, 3H, OCH_3), 6.83 (d, $J=8.8$, 1H, H_{arom}), 7.44 (t, $J=7.9$, 1H, H_{arom}), 7.56–7.69 (m, 2H, H_{arom}), 7.89 (dd, $J=8.8$, 2.7, 1H, H_{arom}), 8.08 (d, $J=7.9$, 1H, H_{arom}), 8.36 (d, $J=2.7$, 1H, H_{arom}); ^{13}C NMR (CDCl_3) δ 53.9 (CH_3), 111.2 (CH), 120.1 (CH), 123.9 (C), 125.8 (CH), 127.1 (CH), 127.6 (C), 132.4 (CH), 136.1 (CH), 139.9 (C), 143.5 (CH), 162.7 (C_{ar}), 165.0 (C); IR (KBr) 1655; MS (EI) m/z (%) 258 (M^+ , 64), 229 (39), 136 (41), 108 (58), 80 (100); HRMS calc. for $\text{C}_{13}\text{H}_{10}\text{N}_2\text{O}_2\text{S}$ 258.0463, found 258.0463.

2-Benzyl-1,2-benzisothiazol-3(2H)-one (1h).⁸

According to the general procedure, benzisothiazolone **1h** was obtained from benzamide **2h** (100 mg, 0.41 mmol) as a white solid in 60% yield after purification by column chromatography (hexanes/EtOAc, 1:1) followed by crystallization from EtOH: mp 79–80 °C (EtOH). Lit.⁸ 86–89 °C ($i\text{PrOH}$); ^1H NMR (CDCl_3) δ 5.02 (s, 2H, CH_2), 7.31–7.77 (m, 8H, H_{arom}), 8.06 (d, $J=7.9$, 1H, H_{arom}); ^{13}C NMR (CDCl_3) δ 47.4 (CH_3), 120.3 (CH), 124.3 (C), 125.4 (CH), 126.7 (CH), 128.2 (CH), 128.3 (CH), 128.7 (CH), 131.7 (CH), 136.0 (C), 140.3 (C), 165.2 (C); IR (KBr) 1670; MS (EI) m/z (%) 241 (M^+ , 97), 152 (99), 108 (70), 91 (100); HRMS calc. for $\text{C}_{14}\text{H}_{11}\text{NOS}$ 241.0561, found 241.0565.

⁸ Grivas, J. C. *J. Org. Chem.* **1975**, *40*, 2029.

2-Methyl-1,2-benzisothiazol-3(2H)-one (1j).

According to the general procedure, benzisothiazolone **1j** was obtained from benzamide **2j** (100 mg, 0.60 mmol) as a white solid in 60% yield after purification by column chromatography (hexanes/EtOAc, 1:1) followed by crystallization from EtOH: mp 50–51°C (EtOH). Lit.⁹ 51–52 °C; ¹H NMR (CDCl₃) _ 3.61 (s, 3H, CH₃), 7.13–7.17 (m, 1H, H_{arom}), 7.37–7.60 (m, 2H, H_{arom}), 7.85–7.89 (m, 1H, H_{arom}); ¹³C NMR (CDCl₃) _ 30.2 (CH₃), 120.0 (CH), 123.2 (C), 125.2 (CH), 126.2 (CH), 131.4 (CH), 139.7 (C), 165.3 (C); IR (KBr) 1680; MS (EI) *m/z* (%) 165 (M⁺, 76), 136 (100), 109 (37); HRMS calc. for C₈H₇NOS 165.0248, found 165.0250.

⁹ Lawson, A J. *J. Chem. Soc., Chem. Commun.* **1981**, 23, 1238.

2e

2