Novel Ozone-Mediated Cleavage of the Benzhydryl Protecting Group from Aziridinyl Esters

Aniruddha P. Patwardhan, Zhenjie Lu, V. Reddy Pulgam and William D. Wulff*

Department of Chemistry

Michigan State University

East Lansing, MI 48824

General Information

All chemicals were purchased from Aldrich. All solvents were reagent grade and were distilled under argon before use: dichloromethane (CH_2Cl_2), toluene and benzene from calcium hydride (CaH_2), tetrahydrofuran and diethyl ether from benzophenone ketyl. Ozonolysis was performed on a Welsbach T23 ozonoator at a flow rate of 0.0965 scfm (standard cubic feet per min) of air. NMR spectra were recorded on a Varian 300 MHz spectrometers using ^1H and ^{13}C solvent peaks as an internal reference (e.g., 7.240 ppm for CDCl_3 and the CDCl_3 triplet at 77.00 ppm). FT-IR were recorded on a Nicolet IR/42 spectrometer. Both low and high resolution mass spectra were obtained from the Michigan State University Mass Spectrometry Facility. Elemental analyses were done by the Michigan State University Elemental Analysis Facility.

General experimental procedure for the deprotection via hydrogenolysis illustrated for 3b.

***cis*-(2R)-Carboxyethyl-(3R)-cyclohexylaziridine (6b):** To a 250 mL flame-dried round bottom flask filled with argon was added 363 mg *cis*-N-benzylhydryl-2-carboxyethyl-3-cyclohexylaziridine (1 mmol), 170 mg Pd(OH)_2 (0.1mmol) (Pd(OH)_2 on carbon 20%, moisture ca 60%) and 80 mL MeOH. The flask was equipped with a vacuum transfer adapter connected to

vacuum and a balloon filled with hydrogen. The valve to vacuum was opened for a few seconds, then switched to the hydrogen balloon, this process was repeated for 4 times. The suspension was stirred at r.t under hydrogen for 4h and filtered through a pad of Celite, washed with ether (20 mL) and the filtrate was concentrated by rotary evaporation, followed by flash chromatography (15 mm $\times$ 300 mm silica gel, 1:1 ether/hexane) to afford **6b** as a white solid in 93% yield (183 mg). White solid: mp: 58-59 °C; R_f =0.19 (1:1 Et₂O/hexane). Spectral data for **6b**: ¹H-NMR(CDCl₃, 300 MHz) δ = 0.90 (br s, 1H), 1.05-1.22 (m, 6H), 1.24 (t, J = 7.1Hz, 3H), 1.40-1.45 (m, 1H), 1.52-1.70 (m, 3H), 1.80-2.00 (m, 2H), 2.60 (d, J = 6.1 Hz, 1H), 4.18 (q, J = 7.1 Hz, 2H); ¹³C (CDCl₃, 75 MHz) δ = 14.14, 25.39, 25.40, 26.01, 30.84, 31.61, 34.24, 36.91, 43.98, 61.06, 170.95; IR (thin film, cm⁻¹) 3165(w), 2924(vs), 2849(s), 1730(vs), 1194(s), 1180(s); mass spectrum, m/z (% rel intensity) 197 M⁺ (4), 168 (32), 124 (100), 95 (84). Anal calcd for C₁₁H₁₉NO₂: C, 66.97; H, 9.71; N, 7.10. Found: C, 66.98; H, 9.30; N, 7.08.

cis-(2R)-Carboxyethyl-(3R)-n-propylaziridine (6a): This reaction was carried out with the general procedure with a slight change in the purification. The product evaporates if rotary evaporation is used to remove the solvent after chromatography, so the eluent was adjusted to 1:1 ether/ pentane and distillation was used to remove the solvent. The aziridine **6a** was afforded as a colorless liquid in 83 % yield (130 mg). R_f =0.13 (1:1 Et₂O/hexane). Spectral data for **6a**: ¹H-NMR (CDCl₃, 300 MHz) δ = 0.79 (t, J = 7.1 Hz, 3H), 1.16 (t, J = 7.1 Hz, 3H), 1.22-1.40 (m, 3H), 1.40-1.50 (m, 2H), 2.04-2.10 (m, 1H), 2.49 (d, J = 6.0 Hz, 1H), 4.08 (q, J = 7.1 Hz, 2H); ¹³C (CDCl₃, 75 MHz) δ = 13.46, 13.99, 20.73, 29.60, 34.27, 38.31, 60.97, 170.73; IR (thin film, cm⁻¹) 3269(br), 2963(m), 1728(s), 1196(s); mass spectrum, m/z (% rel intensity) 157 M⁺ (<1), 128 (31), 68 (29), 54 (100), 41 (38). Anal calcd for C₈H₁₅NO₂: C, 61.12; H, 9.62; N, 8.91. Found: C, 61.17; H, 10.09; N, 8.87.

***cis*-(2R)-Carboxyethyl-(3R)-1,1-dimethylethylaziridine (6c):** This reaction was performed with the general procedure and the change in workup noted for **6a**. Colorless liquid (will evaporate under high vacuum). $R_f = 0.10$ (1:1 Et₂O/hexane). Spectral data for **6c**: ¹H-NMR (CDCl₃, 300 MHz) δ = 0.89 (s, 9H), 1.23 (t, J = 6.9 Hz, 3H), 1.48 (br, s, 1H), 2.09 (d, J = 6.1 Hz, 1H), 2.61 (d, J = 6.6 Hz, 1H), 4.04-4.23 (m, 2H); ¹³C (CDCl₃, 75 MHz) δ = 14.00, 27.42, 31.46, 35.25, 47.36, 60.98, 170.20; IR (thin film, cm⁻¹) 3276(w), 2957(w), 1738(s), 1202(s), 1032(s); mass spectrum, m/z (% rel intensity) 171 M⁺ (<1), 156 (43), 114 (30), 97 (63), 82 (100), 70(88), 55 (63). Anal. calcd for C₉H₁₇NO₂: C, 63.13; H, 10.01; N, 8.18. Found: C, 63.10; H, 9.94; N, 8.32.

General experimental procedure for the deprotection via ozonolysis illustrated for the aziridine 3e.

***cis*-(2R)-Carboxyethyl-(3R)-benzylaziridine (6e):** A 250 mL flame-dried 3-necked round bottom flask was equipped with a CaCl₂ drying tube and a glass tube for ozone inlet and then filled with argon. To the flask was added 357 mg *cis*-N-benzylhydryl-2-carboxyethyl-3-phenylaziridine **3e** (1 mmol) and 100 mL CH₂Cl₂. The flask was then immersed in a dry ice-acetone cooling bath at -78° C while nitrogen was bubbled through the solution. The nitrogen purge was stopped and then ozone was bubbled through the stirring solution until the reaction was complete (~3 h). Upon completion, the addition of ozone was discontinued and then the solution was purged with nitrogen for 5 min to remove unreacted ozone (the color of the solution changed from blue to colorless). The reaction was quenched by the addition of 10 equiv. of methanolic sodium borohydride and then allowed to gradually warm up to r.t, followed by addition of 50 mL deionized water. The reaction

mixture was then extracted with CH_2Cl_2 , dried with MgSO_4 , filtered and the filtrate was concentrated on a rotary evaporator and the resulting oil was purified by flash chromatography (15 mm $\times$ 300 mm silica gel, 1:1 ether/hexane) to afford **6e** as white solid in 60 % yield (114 mg). White solid, mp: 67-68 °C; $R_f = 0.13$ (1:1 Et_2O /hexane). Spectral data for **6e**: ^1H -NMR (CDCl_3 , 300 MHz) $\delta = 0.99$ (t, $J = 7.1$ Hz, 3H), 1.87 (br s, 1H), 3.00 (d, $J = 6.1$ Hz, 1H), 3.47 (d, $J = 6.1$ Hz, 1H), 3.90-4.00 (m, 2H), 7.24-7.33 (m, 5H); ^{13}C (CDCl_3 , 75 MHz) $\delta = 13.90, 29.65, 37.14, 61.07, 127.47, 127.62, 128.01, 134.80, 169.02$; IR (thin film, cm^{-1}) 3218(w), 1738(s), 1186(s), 698(s); mass spectrum, m/z (% rel intensity) 191 M^+ (<1), 162 (58), 118 (100), 104 (12), 91 (100), 88 (21), 77 (8). Anal calcd for $\text{C}_{11}\text{H}_{13}\text{NO}_2$: C, 69.09; H, 6.85; N, 7.32. Found: C, 68.74; H, 6.91; N, 7.25.

All of the reactions were carried out on racemic aziridines **3** except for compound **3e**. A sample of **3e** that was 98 % ee was deprotected to give **6e** which was found to be 97 % ee. The optical purity of **6e** was determined by HPLC analysis on a CHIRALCEL OD-H column with a 90:10 mixture of hexane/2-propanol at 228 nm with a flow rate of 0.7 mL/min. Retention times: $R_t = 6.16$ min (minor) and $R_t = 9.47$ min (major). $[\alpha]_{23\text{D}} = -12.4$ ($c = 1$ in EtOAc) on material that was 99 % ee.

1-Aza-spiro[2.5]octane-2-carboxylic acid ethyl ester (6d): $R_f = 0.06$, yield = 40 %. Spectral data for **6d**: ^1H (300 MHz, CDCl_3) δ 1.2 (t, $J = 7.0$ Hz, 3H), 1.2-1.8 (m, 10), 2.5 (s, 1H), 2.95 (bs, 1H), 4.23 (q, $J = 7.0$ Hz, 2H); ^{13}C (75 MHz, CDCl_3): 14.28, 25.10, 25.46, 25.59, 29.54, 29.66, 36.60, 41.87, 61.20, 170.45; IR: 3362, 2928, 2855, 1740, 1635, 1194; HRMS (FAB) calcd for $\text{C}_{10}\text{H}_{18}\text{NO}_2^+$ m/z 184.1332 ($\text{M}+1$), measd 184.1337.

cis-1H-2-Carboxyethyl-2-methyl-3-phenyl aziridine (6f): $R_f = 0.20$ (1:1 hexanes: Et_2O). Yield 50 %. Spectral data for **6f**: ^1H (300 MHz, CDCl_3) δ 0.94 (t, $J = 7.1$ Hz, 3H), 1.65 (s, 3H), 1.94 (br s, 1H), 3.23 (s, 1H), 3.91 (q, $J = 7.1$ Hz, 2H), 7.2-7.4 (bs, 5H); ^{13}C (75 MHz, CDCl_3) δ 13.82, 20.35, 42.99, 47.56, 61.23, 127.28, 127.47, 127.85, 127.92, 128.01, 135.49, 170.78; IR: 3303, 2980, 2936, 1729, 1197, 698; HRMS (FAB) calcd for m/z 205.1102 (M^+), measd 205.1097.

cis-1H-2-Carboxyethyl-3-(4-tolyl)-aziridine (6g): White solid, mp = 54-55° C. $R_f = 0.14$ (1:1 hexanes: Et_2O), yield = 51 %. Spectral data for **6g**: ^1H (300 MHz, CDCl_3) δ 1.05 (t, $J = 7.0$ Hz, 3H), 1.75 (bs, 1H), 2.30 (s, 3H), 2.96 (d, $J = 6.36$ Hz, 1H), 3.43 (d, $J = 6.4$ Hz, 1H), 3.98 (m, 2H), 7.10 (d, $J = 8.1$ Hz, 2H), 7.21 (d, $J = 8.05$ Hz, 2H); ^{13}C (75 MHz, CDCl_3) δ 13.90, 21.05, 37.03, 39.85, 61.008, 127.28, 128.67, 131.69, 137.22, 169.11; IR: 3310, 2984, 1734, 1383, 1200, 1028, 833; HRMS (FAB) calcd for $\text{C}_{12}\text{H}_{16}\text{NO}_2^+$ m/z 206.1175 ($\text{M}+1$), measd 206.1177.

***cis*-1H-2-Carboxyethyl-3-biphenyl-aziridine (6h):** White solid, mp = 61-62° C, R_f = 0.07 (1:1 hexane:Et₂O), yield = 60 %. Spectral data for **6h**: ¹H (300 MHz, CDCl₃) δ 1.0 (bs, 3H), 1.78 (bs, 1H), 3.0 (bs, 1H), 3.45 (bs, 1H), 4.0 (bs, 2H), 7.2-7.62 (m, 9H). ¹³C (75 MHz, CDCl₃) δ 13.95, 37.34, 40.68, 61.07, 126.70, 126.97, 127.27, 127.29, 127.92, 128.69, 133.94, 140.54, 169.02; IR: 3310, 2982, 1734, 1487, 1204, 1030, 760, 698; HRMS (FAB) calcd for C₁₇H₁₈NO₂⁺ m/z 268.1332 (M+1), measd 268.1341.

***cis*-(2R)-Carboxyethyl-(3R)-*p*-bromobenzylaziridine (6i):** White solid, mp = 78.5-79 °C; R_f = 0.12 (1:1 Et₂O/hexane). Spectral data for **6i**: ¹H-NMR (CDCl₃, 300 MHz) δ = 1.02 (t, J = 7.1Hz, 3H), 1.67 (br s, 1H), 3.01 (d, J = 6.6 Hz, 1H), 3.40 (d, J = 6.4 Hz, 1H), 3.93-3.98 (m, 2H), 7.20 (d, J = 8.5 Hz, 2H), 7.41 (d, J = 8.5 Hz, 2H); ¹³C (CDCl₃, 75 MHz) δ = 13.88, 37.05, 38.98, 61.06, 121.45, 129.18, 130.98, 133.85, 168.62; IR (thin film, cm⁻¹) 3222(br s), 1734(vs), 1194(vs), 1170(s), 843(w); mass spectrum, m/z (% rel intensity) 271 M⁺ (⁸¹Br, 2), 269 M⁺ (⁷⁹Br, 2), 226 (⁸¹Br, 8), 224 (⁷⁹Br, 11), 197 (⁸¹Br, 100), 195 (⁷⁹Br, 100), 116 (96), 89 (97), 88 (21), 77 (13), 63 (26). Anal calcd for C₁₁H₁₂BrNO₂: C, 48.91; H, 4.48; N, 5.19. Found: C, 48.88; H, 4.50; N, 5.13.