A Novel Synthesis of Desymmetrized Resorcinol Derivatives: Aryl Fluoride Displacement on Deactivated Substrates

*Aujin Kim[†], Jeremiah D. Powers, Jennifer F. Toczko**

Chemical Development, GlaxoSmithKline, Research Triangle Park, NC, 27709

jennifer.f.toczko@gsk.com

Table of Contents:

Synthesis and Characterization of Compounds 2a-12c	S2-S12
References	S12

General. All starting materials, reagents, and solvents were obtained from commercial suppliers and used without further purification. Organic extracts were dried over anhydrous MgSO_4 , filtered, and concentrated under reduced pressure. Silica gel chromatography was carried out with EMD silica gel, 230-400 mesh according to Still's procedure.¹ HPLC analyses were carried out using a Phenomenex® Luna 3 μm C18(2) column (50 x 2.0 mm). ^1H NMR (300 MHz) were measured in CDCl_3 . ^{13}C NMR (100 MHz) were carried in DMSO-d_6 unless otherwise indicated. IR spectra were collected using DATR (Diamond Attenuated Total Reflectance).

3-Fluorophenyl phenylmethylether (2a). To a solution of KOtBu (24.6 g, 219 mmol) and toluene (50 mL), benzyl alcohol (29 mL, 280 mmol) was slowly added, followed by DMPU (20 mL). The solution was heated at 80 °C for 30 min. 1,3-Difluorobenzene (**1**) (9 mL, 92 mmol) was then added and the solution heated for an additional 3.5 hrs at 80 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine(x 2), dried and concentrated. The resulting oil was purified by flash chromatography (1% EtOAc in hexanes) to produce 16.7 g (89%) of **2a** as a colorless oil.²

^1H NMR δ 5.05 (s, 2H), 6.64-6.78 (m, 3H), 7.23 (q, 1H, J = 8.5, 15.2 Hz), 7.31-7.45 (m, 5H). ^{13}C NMR (125 MHz, CDCl_3) δ 70.2, 102.6 (d, J = 24 Hz), 107.7 (d, J = 21 Hz), 110.6, 127.5, 128.1, 128.6, 130.2, 136.5, 160.1 (d, J = 11 Hz), 163.6 (d, J = 245 Hz). IR 1025, 1164, 1262, 1277, 1488, 1592, 1610, 2873, 2917, 2931, 3034, 3067, 3090 cm^{-1} . Analysis calculated for $\text{C}_{13}\text{H}_{11}\text{OF}$: C, 77.21; H, 5.48; O, 7.91; F, 9.1. Found: C 77.34; H, 5.57; O, 8.06; F, 9.43.

3-Fluoroanisole (2b). To a solution of KOtBu (7.4 g, 66 mmol) and DME (15 mL), methanol (3.2 mL, 79 mmol) was slowly added, followed by DMPU (6 mL). The solution was heated at 80 °C for 30 min.

1,3-Difluorobenzene (**1**, 2.6 mL, 26 mmol) was then added and the solution heated for an additional 3 hrs at 80 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (1% EtOAc in hexanes) to produce 2.7 g (80%) of **2b** as a colorless oil.

¹H NMR δ 3.79 (s, 3H), 6.58-6.70 (m, 3H), 7.18-7.25 (m, 1H). Analysis calculated for C₇H₁₇OF: C, 66.66; H, 5.59. Found: C, 66.80; H, 5.69. The spectral data are in good agreement when compared with an authentic sample.

1-(Butyloxy)-3-[(phenylmethyl)oxy]benzene (3a). To a solution of KOtBu (6.66 g, 59.4 mmol) and toluene (11 mL), butanol (7 mL, 77 mmol) was slowly added, followed by DMPU (6 mL). The solution was heated at 100 °C for 30 min. 3-Fluorophenyl phenylmethyl ether (**2a**, 3.19 g, 15.8 mmol) in toluene (1 mL) was then added and the solution heated for an additional 21 hrs at 100 °C (After 4 h, additional KOtBu (3.3 g, 29 mmol) was added to the reaction mixture). The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), 6N HCl, dried and concentrated. The resulting oil was purified by flash chromatography (0.5% EtOAc in hexanes) to produce 3.6 g (90%) of **3a** as a colorless oil.

¹H NMR δ 0.96 (t, 3H, *J* = 7.3 Hz), 1.41-1.53 (m, 2H), 1.70-1.79 (m, 2H), 3.93 (t, 2H, *J* = 6.45 Hz), 5.03 (s, 2H), 6.49-6.57 (m, 3H), 7.13-7.19 (m, 1H), 7.31-7.44 (m, 5H). ¹³C NMR δ 13.6, 18.7, 30.7, 67.0, 69.1, 101.4, 106.8, 106.9, 127.6, 127.7, 128.4, 129.9, 137.1, 159.5, 159.9. IR 1027, 1145, 1177, 1261, 1285, 1454, 1491, 1589, 2871, 2932, 2957, 3032, 3065, 3089 cm⁻¹. Analysis calculated for C₁₇H₂₀O₂: C, 79.58; H, 7.80; O, 12.48. Found: C 79.75; H, 7.84; O, 12.60.

1-(Butyloxy)-3-(methyloxy)benzene (3b). To a solution of KOtBu (3.5 g, 30.3 mmol) and toluene (5 mL), 1-butanol (3.8 mL, 42 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 100 °C for 30 min. 3-Fluoroanisole (**2b**, 0.90 mL, 7.9 mmol) was then added and the solution heated for an additional 21 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2) and 6N HCl. The combined aqueous layers were extracted with EtOAc and the combined organic layers were dried and concentrated. The resulting oil was purified by flash chromatography (2% EtOAc in hexanes) to produce 1.0 g (71%) of **3b** as a colorless oil.³

¹H NMR δ 0.96 (t, 3H, J = 7.4 Hz), 1.41-1.54 (m, 2H), 1.70-1.80 (m, 2H), 3.78 (s, 3H), 3.93 (t, 2H, J = 6.5 Hz) 6.45-6.51 (m, 3H), 7.16 (t, 1H, J = 8 Hz). ¹³C NMR δ 13.6, 18.7, 30.7, 55.0, 67.0, 100.6, 106.1, 106.5, 129.8, 159.9, 160.5. IR 1042, 1148, 1199, 1264, 1286, 1453, 1492, 1590, 2835, 2872, 2935, 2985, 3030, 3067, 3090 cm⁻¹. Analysis calculated for C₁₁H₁₆O₂: C, 73.30; H, 8.95. Found: C 73.23; H, 9.04.

1-[(Cyclopropylmethyl)oxy]-3-[(phenylmethyl)oxy]benzene (4a). To a solution of KOtBu (3.3 g, 30 mmol) and toluene (7.5 mL), cyclopropylmethanol (3 mL, 37 mmol) was slowly added, followed by DMPU (3 mL). The solution was heated at 100 °C for 30 min. 3-Fluorophenyl phenylmethyl ether (**2a**, 1.50 g, 7.42 mmol) was then added and the solution heated for an additional 21 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (6% EtOAc in hexanes) to produce 1.66 g (88%) of **4a** as a yellow oil.

¹H NMR δ 0.30-0.35 (m, 2H), 0.60-0.66 (m, 2H), 1.19-1.33 (m, 1H), 3.77 (d, 2H, J = 7.1 Hz), 5.03 (s, 2H), 6.50-6.57 (m, 3H), 7.16 (t, 1H, J = 8.6 Hz), 7.31-7.43 (m, 5H). ¹³C NMR δ 3.0, 10.0, 69.1, 71.9, 101.4, 106.8, 106.9, 127.5, 127.7, 128.3, 129.8, 137.1, 159.5, 159.8. IR 1026, 1144, 1179, 1261,

1284, 1405, 1453, 1490, 1590, 2871, 2910, 3007, 3030, 3066, 3082 cm^{-1} . Analysis calculated for $\text{C}_{17}\text{H}_{18}\text{O}_2$: C, 80.29; H, 7.13; O, 12.58. Found: C 80.19; H, 7.05; O, 12.50.

1-[(Cyclopropylmethyl)oxy]-3-(methoxy)benzene (4b). To a solution of KOtBu (3.6 g, 32 mmol) and toluene (5 mL), cyclopropylmethanol (3.2 mL, 42 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 100 °C for 30 min. 3-Fluoroanisole (**2b**, 0.91 mL, 7.9 mmol) was then added and the solution heated for an additional 22 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (6% EtOAc in hexanes) to produce 1.5 g (75%) of **4b** as a colorless oil.

^1H NMR δ 0.31-0.36 (m, 2H), 0.60-0.67 (m, 2H), 1.20-1.33 (m, 1H), 3.77 (d, 2H, $J = 6.8$ Hz), 3.78 (s, 3H), 6.46-6.51 (m, 3H), 7.16 (t, 1H, $J = 8.3$ Hz). ^{13}C NMR δ 3.0, 10.1, 55.0, 71.9, 100.6, 106.1, 106.5, 129.8, 159.9, 160.5. IR 1015, 1041, 1145, 1198, 1264, 1285, 1405, 2835, 2874, 2911, 2941, 3005, 3082 cm^{-1} . Analysis calculated for $\text{C}_{11}\text{H}_{14}\text{O}_2$: C, 74.13; H, 7.92. Found: C 74.28; H, 7.98.

1-{[2-(Methoxy)ethyl]oxy}-3-[(phenylmethyl)oxy]benzene (5a). To a solution of KOtBu (3.32 g, 29.7 mmol) and toluene (6 mL), 2-methoxyethanol (3.6 mL, 38 mmol) was slowly added, followed by DMPU (3 mL). The solution was heated at 100 °C for 30 min. 3-Fluorophenyl phenylmethyl ether (**2a**, 1.55 g, 7.67 mmol) in toluene (1 mL) was then added and the solution heated for an additional 24 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine, and 6N HCl. The combined aqueous layers were extracted with EtOAc and the combined organic layers were dried and

concentrated. The resulting oil was purified by flash chromatography (1% EtOAc in hexanes) to produce 1.25 g (65%) of **5a** as a colorless oil.

^1H NMR δ 3.44 (s, 3H), 3.71-3.75 (m, 2H), 4.07-4.10 (m, 2H), 5.03 (s, 2H), 6.52-6.60 (m, 3H), 7.14-7.19 (m, 1H), 7.29-7.44 (m, 5H). ^{13}C NMR δ 58.1, 66.8, 69.2, 70.3, 101.5, 106.8, 107.2, 127.6, 127.8, 128.4, 129.9, 137.1, 159.5, 159.7. IR 1026, 1124, 1151, 1178, 1262, 1286, 1451, 1490, 1589, 2819, 2877, 2926, 3032 cm^{-1} . Analysis calculated for $\text{C}_{16}\text{H}_{18}\text{O}_3$: C, 74.38; H, 7.04; O, 18.58. Found: C 74.37; H, 7.15; O, 18.63.

1-(Methoxy)-3-[[2-(methoxy)ethyl]oxy]benzene (5b). To a solution of KOtBu (3.6 g, 32 mmol) and toluene (5 mL), 2-methoxyethanol (4 mL, 50 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 100 $^{\circ}\text{C}$ for 30 min. 3-Fluoroanisole (**2b**, 0.91 mL, 7.9 mmol) was then added and the solution heated for an additional 46 hrs at 100 $^{\circ}\text{C}$. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (25% EtOAc in hexanes) to produce 0.96 g (67%) of **5b** as a colorless oil.

^1H NMR δ 3.44 (s, 3H), 3.72-3.75 (m, 2H), 3.77 (s, 3H), 4.07-4.12 (m, 2H), 6.49-6.53 (m, 3H), 7.13-7.19 (m, 1H). ^{13}C NMR δ 55.0, 58.1, 66.8, 70.4, 100.6, 106.3, 106.5, 129.9, 159.7, 160.5. IR 1035, 1123, 1152, 1199, 1265, 1286, 1492, 1589, 2820, 2836, 2881, 2928, 3070, 3089 cm^{-1} . Analysis calculated for $\text{C}_{10}\text{H}_{14}\text{O}_2$: C, 65.92; H, 7.74. Found: C 65.92; H, 7.81.

1-[(2,2-Dimethylpropyl)oxy]-3-[(phenylmethyl)oxy]benzene (6a). To a solution of KOtBu (3.37 g, 30.0 mmol) and toluene (7.5 mL), neopentyl alcohol (3.4 g, 39 mmol) was slowly added, followed by

DMPU (3 mL). The solution was heated at 100 °C for 30 min. 3-Fluorophenyl phenylmethyl ether (**2a**, 1.48 g, 7.32 mmol) was then added and the solution heated for an additional 44 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried, and concentrated. The resulting oil was purified by flash chromatography (3% EtOAc in hexanes) to produce 1.55 g (77%) of **6a** as a colorless oil.

^1H NMR δ 1.02 (s, 9H), 3.56 (s, 2H), 5.04 (s, 2H), 6.50-6.607 (m, 3H), 7.14-7.19 (m, 1H), 7.32-7.45 (m, 5H). ^{13}C NMR δ 26.3, 31.5, 69.1, 77.1, 101.4, 106.9, 127.5, 127.7, 128.3, 129.8, 137.1, 159.5, 160.3. IR 1026, 1049, 1147, 1177, 1261, 1288, 1453, 1491, 1591, 2868, 2904, 2955, 3033, 3065, 3090 cm^{-1} . Analysis calculated for $\text{C}_{18}\text{H}_{22}\text{O}_2$: C, 79.96; H, 8.22; O, 11.84. Found: C 80.13; H, 8.27; O, 11.70

1-[(2,2-Dimethylpropyl)oxy]-3-(methoxy)benzene (6b). To a solution of K₂CO₃ (3.6 g, 32 mmol) and toluene (5 mL), neopentyl alcohol (3.55 g, 40.3 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 100 °C for 30 min. 3-Fluoroanisole (**2b**, 0.91 mL, 7.9 mmol) was then added and the solution heated for an additional 41 hrs at 100 °C (At 19 h, additional K₂CO₃ (1.8 g, 16 mmol) was added). The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated under reduced pressure. The resulting oil was purified by flash chromatography (2% EtOAc in hexanes) to produce 1.25 g (81%) of **6b** as a colorless oil.

^1H NMR δ 1.04 (s, 9H), 3.58 (s, 2H), 3.80 (s, 3H), 6.48-6.53 (m, 3H), 7.17 (t, 1H, $J = 7.8$ Hz). ^{13}C NMR δ 26.3, 31.5, 55.0, 77.1, 100.5, 106.2, 106.6, 129.8, 160.3, 160.5. IR 1021, 1051, 1148, 1198, 1264, 1289, 1364, 1400, 2835, 2869, 2904, 2955, 3030, 3070, 3089 cm^{-1} . Analysis calculated for $\text{C}_{12}\text{H}_{18}\text{O}_2$: C, 74.18; H, 9.36; O, 16.47. Found: C 74.41; H, 9.55; O, 16.20.

1-[(1-Ethylpropyl)oxy]-3-[(phenylmethyl)oxy]benzene (7a). To a solution of KOtBu (3.4 g, 30 mmol) and toluene (7.5 mL), 3-pentanol (4 mL, 37 mmol) was slowly added, followed by DMPU (3 mL). The solution was heated at 100 °C for 30 min. 3-Fluorophenyl phenylmethyl ether (**2a**, 1.5 g, 7.4 mmol) was then added and the solution heated for an additional 46 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (2% EtOAc in hexanes) to produce 1.8 g (91%) of **7a** as a colorless oil.

^1H NMR δ 0.94 (t, 6H, $J = 7.5$ Hz), 1.61-1.70 (m, 4H), 4.03-4.11 (m, 1H), 5.03 (s, 2H), 6.49-6.56 (m, 3H), 7.13-7.18 (m, 1H), 7.32-7.45 (m, 5H). ^{13}C NMR δ 9.3, 25.5, 69.1, 78.9, 102.6, 106.8, 108.0, 127.6, 127.7, 128.4, 129.9, 137.1, 159.5, 159.6. IR 1027, 1143, 1178, 1260, 1283, 1488, 1586, 2877, 2935, 2965, 3032, 3065, 3089 cm^{-1} . Analysis calculated for $\text{C}_{18}\text{H}_{22}\text{O}_3$: C, 79.96; H, 8.20. Found: C 79.88; H, 8.31.

1-[(1-Ethylpropyl)oxy]-3-(methoxy)benzene (7b). To a solution of KOtBu (3.6 g, 32 mmol) and toluene (5 mL), 3-pentanol (4.4 mL, 41 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 100 °C for 30 min. 3-Fluoroanisole (**2b**, 0.91 mL, 7.9 mmol) was then added and the solution heated for an additional 44 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (3% EtOAc in hexanes) to produce 1.3 g (84%) of **7b** as a colorless oil.

^1H NMR δ 0.94 (t, 6H, J = 7.5 Hz), 1.62-1.71 (m, 4H), 3.78 (s, 3H), 4.05-4.12 (m, 1H), 6.46-6.51 (m, 3H), 7.15 (t, 1H, J = 8.8 Hz). ^{13}C NMR δ 9.3, 25.5, 54.9, 78.9, 101.8, 105.9, 107.6, 129.9, 159.5, 160.5. IR 1041, 1145, 1198, 1263, 1283, 2835, 2878, 2937, 2965, 3029, 3070, 3089 cm^{-1} . Analysis calculated for $\text{C}_{12}\text{H}_{18}\text{O}_2$: C, 74.19; H, 9.34. Found: C 74.14; H, 9.45.

2-Fluorophenyl phenylmethyl ether (10). To a solution of K₂CO₃ (2.46 g, 21.9 mmol) and toluene (5 mL), benzyl alcohol (2.8 mL, 27 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 80 °C for 30 min. 1,2-Difluorobenzene (**8**, 0.86 mL, 8.8 mmol) was then added and the solution heated for an additional 3 hrs at 80 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (3% EtOAc in hexanes) to produce 1.36 g (77%) of **10** as white crystals mp, 35 °C.⁴

^1H NMR δ 5.14 (s, 2H), 6.86-7.12 (m, 4H), 7.29-7.46 (m, 5H). ^{13}C NMR δ 70.1, 115.5, 116.0 (d, J = 19 Hz), 121.2 (d, J = 7 Hz), 124.7 (d, J = 10 Hz), 127.8, 128.0, 128.5, 136.6, 146.2 (d, J = 10.7 Hz), 151.9 (d, J = 243 Hz). IR 1015, 1106, 1202, 1254, 1280, 1502, 1589, 1613, 2867, 2915, 2924, 3032, 3036, 3078, 3088 cm^{-1} . Analysis calculated for $\text{C}_{13}\text{H}_{11}\text{OF}$: C, 77.21; H, 5.48. Found: C, 77.18; H, 5.47.

4-Fluorophenyl phenylmethyl ether (11). To a solution of K₂CO₃ (1.0 g, 8.8 mmol) and toluene (5 mL), benzyl alcohol (1.1 mL, 11 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 80 °C for 30 min. 1,4-Difluorobenzene (**9**, 0.85 mL, 8.8 mmol) was then added and the

solution heated for an additional 5 hrs at 80 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (2% EtOAc in hexanes) to produce 0.45 g (25%) of **11** as white crystals.⁵

¹H NMR δ 5.05 (s, 2H), 6.90-7.02 (m, 4H), 7.32-7.46 (m, 5H) ¹³C NMR δ 69.7, 115.8 (d, J = 19 Hz), 116.0 (d, J = 6 Hz), 127.7, 127.8, 128.4, 136.9, 155.6 (d, J = 97 Hz), 156.1 (d, J = 286 Hz). IR 694, 731, 759, 831, 1026, 1040, 1197, 1248, 1451, 1502, 1598, 2856, 2888, 3036, 3040, 3052, 3064 cm⁻¹. Analysis calculated for C₁₃H₁₁FO: C, 77.21; H, 5.48. Found: C, 77.54; H, 5.43.

1-(Butyloxy)-2-[(phenylmethyl)oxy]benzene (12a). To a solution of KOtBu (0.73 g, 6.5 mmol) and toluene (4 mL), butanol (0.71 mL, 7.8 mmol) was slowly added, followed by DMPU (1 mL). The solution was heated at 100 °C for 30 min. 2-Fluorophenyl phenylmethyl ether (**10**, 0.52 g, 2.6 mmol) in toluene (1 mL) was then added and the solution heated for 18 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (4% EtOAc in hexanes) to produce 0.51 g (77%) of **12a** as a colorless oil.

¹H NMR δ 0.97 (t, 3H, J = 7.3 Hz), 1.45-1.58 (m, 2H), 1.77-1.86 (m, 2H), 4.03 (t, 2H, J = 6.5 Hz), 5.13 (s, 2H), 6.81-6.92 (m, 4H), 7.29-7.46 (m, 5H). ¹³C NMR δ 13.7, 18.8, 30.9, 68.0, 70.0, 113.9, 114.7, 120.8, 121.3, 127.4, 127.7, 128.3, 137.5, 148.1, 148.8. IR 695, 734, 1123, 1211, 1251, 1451, 1500, 1592, 2871, 2932, 2957, 3033, 3064 cm⁻¹. Analysis calculated for C₁₇H₂₀O₂: C, 79.65; H, 7.86. Found: C, 79.61; H, 8.02.

1-[(2,2-Dimethylpropyl)oxy]-2-[(phenylmethyl)oxy]benzene (12b). To a solution of KOtBu (2.16 g, 19.2 mmol) and toluene (5 mL), neopentyl alcohol (2.18 mL, 24.7 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 80 °C for 30 min. 2-Fluorophenyl phenylmethyl ether (**10**, 1.0 g, 5.0 mmol) in toluene (1 mL) was then added and the solution heated for 29 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting oil was purified by flash chromatography (2% EtOAc in hexanes) to produce 1.1 g (84%) of **12b** as a colorless oil.

^1H NMR δ 1.07 (s, 9H), 3.66 (s, 2H), 5.11 (s, 2H), 6.83-6.97 (m, 4H), 7.27-7.49 (m, 5H). ^{13}C NMR δ 28.4, 31.8, 70.1, 78.1, 113.8, 115.0, 120.7, 121.6, 127.0, 127.5, 128.3, 137.6, 148.1, 149.5. IR 694, 737, 1016, 1121, 1219, 1251, 1364, 1451, 1501, 1593, 2867, 2902, 2954, 3033, 3064 cm^{-1} . Analysis calculated for $\text{C}_{18}\text{H}_{22}\text{O}_2$: C, 79.96; H, 8.20. Found: C, 79.92; H, 8.28.

1-[(1-Ethylpropyl)oxy]-2-[(phenylmethyl)oxy]benzene (12c). To a solution of KOtBu (2.22 g, 19.8 mmol) and toluene (4 mL), 3-pentanol (2.7 mL, 25 mmol) was slowly added, followed by DMPU (2 mL). The solution was heated at 100 °C for 30 min. 2-Fluorophenyl phenylmethyl ether (**10**, 1.0 g, 5.0 mmol) in toluene (1 mL) was then added and the solution heated for 44 hrs at 100 °C. The reaction mixture was cooled to rt, and washed with water, 10% brine (x 2), dried and concentrated. The resulting

oil was purified by flash chromatography (3% EtOAc in hexanes) to produce 1.0 g (75%) of **12c** as a colorless oil.

¹H NMR δ 0.97 (t, 6H, J = 7.6 Hz), 1.63-1.76 (m, 4H), 4.09-4.16 (m, 1H), 5.12 (s, 2H), 6.82-6.95 (m, 4H), 7.26-7.46 (m, 5H). ¹³C NMR δ 9.2, 25.5, 70.0, 80.4, 114.9, 116.6, 121.1, 121.3, 127.4, 127.6, 128.3, 137.5, 148.0, 149.1. IR 695, 737, 1122, 1208, 1252, 1451, 1495, 1591, 2877, 2935, 2964, 3033, 3064 cm⁻¹. Analysis calculated for C₁₈H₂₂O₂: C, 79.96; H, 8.20. Found: C, 79.89; H, 8.25.

[†] GlaxoSmithKline Summer Intern, 2005

¹ Still, W.C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923.

² Durrani, A. A.; Tyman, J.H.P. *J. Chem. Soc., Perkin Trans. 1* **1979**, 2079.

³ Wolter, M.; Nordmann, G.; Job, G.E.; Buchwald, S.L. *Org. Lett.* **2002**, *4*, 973.

⁴ Marsh, J.P.; Goodman, L. *J. Org. Chem.* **1965**, *30*, 2491.

⁵ Sinhababu, A.K.; Kawase, M.; Borchardt, R.T. *Tetrahedron Lett.* **1987**, *28*, 4139.