Supporting Information

Novel Tunable CuX₂-mediated Cyclization Reaction of Cyclopropylideneacetic Acids and Esters for the Facile Synthesis of 4-Halomethyl-2(5*H*)-furanones and 4-Halo-5,6-dihydro-2*H*-pyran-2-ones.

Xian Huang^{*,a,b} and Hongwei Zhou^a

^a*Department of Chemistry, Zhejiang University(Campus Xixi), Hangzhou 310028, P.R. China*

^b*State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, P.R. China*

huangx@mail.Hz.zj.cn Fax: +86-571-88807077

Experimental Section

Typical experimental procedure

General. All ¹H NMR and ¹³C NMR spectra were measured in CDCl₃ and recorded on Bruker Avance – 400 (100M Hz) spectrometer spectra with TMS as the internal standard. Chemical shifts are expressed in ppm and *J* values are given in Hz. IR spectra were run on a Bruker vector 22 spectrometer. EIMS were determined with a

HP5989B mass spectrometer. Melting points are uncorrected. All the reactions in this paper were performed directly in air. Solvents except for diethyl ether used in the reaction described here were distilled before use.

Typical Procedure. Condition A. A solution of **2a** (63 mg, 0.5 mmol) with CuBr₂ (450 mg, 2.0 mmol) in CH₃CN/H₂O (4:1, 10 mL) was stirred at 85°C (reflux) for 10 h. The mixture was then diluted with 20mL of saturated NH₄Cl and extracted three times with diethyl ether. The ether phases were combined and dried over MgSO₄. After evaporation, the residues were purified via chromatography on silica gel with *n*-hexane/diether ether (2:1) as the eluent to afford 70 mg (79%) of **3a**.

Condition B. A solution of **2a** (63 mg, 0.5 mmol) with I₂ (510 mg, 2.0 mmol) and CuI (10 mg, 0.05 mmol) in CH₃CN/H₂O (4:1, 10 mL) was stirred at 85°C (reflux) for 10 h. The mixture was then diluted with 20mL of saturated Na₂S₂O₃ and extracted three times with diethyl ether. The ether phases were combined and dried over MgSO₄. After evaporation, the residues were purified via chromatography on silica gel with *n*-hexane/diether ether (2:1) as the eluent to afford 55 mg (49%) of **3b**.

4-bromomethyl-2(5H)-furanone (3a): oil, ¹H NMR (400M Hz, CDCl₃) δ 6.14 (d, 1H, *J* = 1.15 Hz), 4.94 (t, 2H, *J* = 1.10 Hz), 4.23 (s, 2H) ¹³C NMR (100M Hz, CDCl₃) δ 172.5, 163.2, 118.9, 71.9, 22.5. EIMS *m/z* (relative intensity, %) 178 [M⁺ (⁸¹Br), 6.53], 176 [M⁺ (⁷⁹Br), 6.27], 149 [M⁺ (⁸¹Br)–29, 100], 147 [M⁺ (⁷⁹Br) – 29, 97.18], 96 [M⁺ –Br, 58.56], 81 [⁸¹Br⁺, 13.87], 79 [⁷⁹Br⁺, 14.24] IR (neat) 1785.4, 1751.6 cm⁻¹; Anal. Calcd. For C₅H₅BrO₂: C, 33.93; H, 2.85. Found: C, 33.81; H, 2.81.

4-iodomethyl-2(5H)-furanone (3b): oil, ^1H NMR (400M Hz, CDCl_3) δ 6.14 (m, 1H), 5.02 (t, 2H, $J = 0.76$ Hz), 4.15 (d, 2H, $J = 0.80$ Hz) EIMS m/z (relative intensity, %) 224 [M^+ , 46.75], 127 [I^+ , 33.79], 97 [$\text{M}^+ - \text{I}$, 87.63], 67 [100]. IR (neat), 1780, 1740 cm^{-1} ; Anal. Calcd. For $\text{C}_5\text{H}_5\text{IO}_2$: C, 26.81; H, 2.25. Found: C, 26.70; H, 2.22.

4-iodo-5,6-dihydro-2H-pyran-2-one (4a), oil, ^1H NMR (400M Hz, CDCl_3) δ 6.77(s, 1H), 4.37-4.40 (t, 3H, $J = 6.12$ Hz), 2.91-2.95 (m, 2H) EIMS m/z (relative intensity, %) 224 [M^+ , 4.69], 127 [I^+ , 5.64], 97 [$\text{M} - \text{I}^+$, 7.87], 43 [100.] IR (neat), 1716 cm^{-1} ; Anal. Calcd. For $\text{C}_5\text{H}_5\text{BrO}_2$: C, 26.81; H, 2.25. Found: C, 26.90; H, 2.28.

3-methyl-4-bromo-5,6-dihydro-2H-pyran-2-one (4b): oil, ^1H NMR (400M Hz, CDCl_3) δ 4.36-4.39 (t, 2H, $J = 6.25$ Hz), 2.91-2.95 (m, 2H), 2.06-2.07 (t, 3H, $J = 1.89$ Hz) ^{13}C NMR (100M Hz, CDCl_3) δ 162.7, 138.4, 115.8, 65.4, 35.3, 16.9. EIMS m/z (relative intensity, %) 192 [M^+ (^{81}Br), 22.61], 190 [M^+ (^{79}Br), 23.25], 111 [$\text{M}^+ - \text{Br}$, 42.83], 53 [100] IR (neat) 1716.9 cm^{-1} ; Anal. Calcd. For $\text{C}_6\text{H}_7\text{BrO}_2$: C, 37.73; H, 3.69. Found: C, 37.59; H, 3.65.

3-methyl-4-iodo-5,6-dihydro-2H-pyran-2-one (4c): oil, ^1H NMR (400M Hz, CDCl_3) δ 4.31-4.34 (t, 2H, $J = 6.14$ Hz), 2.99-3.03 (m, 2H), 2.14-2.15 (t, 3H, $J = 1.87$ Hz) ^{13}C NMR (100M Hz, CDCl_3) δ 159.9, 133.8, 66.0, 39.3, 21.9. EIMS m/z (relative intensity, %) 238 [M^+ , 2.56], 127 [I^+ , 6.59], 111 [$\text{M} - \text{I}$, 7.51], 43 [100] IR (neat) 1714.0 cm^{-1} ; Anal. Calcd. For $\text{C}_6\text{H}_7\text{IO}_2$: C, 30.28; H, 2.96. Found: C, 30.15; H, 2.93.

3-ethyl-4-bromo-5,6-dihydro-2H-pyran-2-one (4d): oil, ^1H NMR (400M Hz, CDCl_3) δ 4.34-4.37 (t, 2H, $J = 6.18$ Hz), 2.90-2.94 (t, 2H, $J = 6.25$ Hz), 2.51-2.56 (q, 2H, $J = 7.48$ Hz), 1.07-1.10 (t, 2H, $J = 7.46$ Hz) ^{13}C NMR (100M Hz, CDCl_3) δ 161.7, 137.3,

133.5, 64.8, 34.9, 24.1, 11.7. EIMS m/z (relative intensity, %) 206 [M^+ (^{81}Br), 37.15], 204 [M^+ (^{79}Br), 36.36], 125 [$M^+ - \text{Br}$, 68.42], 79 [Br^+ , 100] IR (neat) 1717.8cm^{-1} ; Anal. Calcd. For $\text{C}_7\text{H}_9\text{BrO}_2$: C, 41.00; H, 4.42. Found: C, 41.15; H, 4.39.

3-ethyl-4-iodo-5,6-dihydro-2H-pyran-2-one (4e): oil, ^1H NMR (400M Hz, CDCl_3) δ 4.27-4.30 (t, 2H, $J = 6.09$ Hz), 2.98-3.01 (t, 2H, $J = 6.20$ Hz), 2.51-2.56 (q, 2H, $J = 7.48$ Hz) 1.07-1.09 (t, 3H, $J = 7.46$ Hz) ^{13}C NMR (100M Hz, CDCl_3) δ 159.4, 138.8, 115.1, 66.0, 39.4, 29.3, 11.8. EIMS m/z (relative intensity, %) 253 [$M^+ + 1$, 34.93], 252 [M^+ , 38.43], 127 [I^+ , 11.37], 125 [$M^+ - \text{I}$, 36.67], 41 [100] IR (neat) 1714.6 cm^{-1} ; Anal. Calcd. For $\text{C}_7\text{H}_9\text{IO}_2$: C, 33.36; H, 3.60. Found: C, 33.24; H, 3.56.

3-*n*-propyl-4-bromo-5,6-dihydro-2H-pyran-2-one (4f): oil, ^1H NMR (400M Hz, CDCl_3) δ 4.34-4.37 (t, 2H, $J = 6.25$ Hz), 2.91-2.95 (t, 2H, $J = 6.27$ Hz), 2.48-2.52 (t, 2H, $J = 7.84$ Hz), 1.50-1.56 (m, 2H), 0.94-0.98 (t, 3H, $J = 7.38$ Hz) ^{13}C NMR (100M Hz, CDCl_3) δ 162.3, 138.2, 132.9, 65.3, 35.5, 32.9, 21.2, 13.8. EIMS m/z (relative intensity, %) 220 [M^+ (^{81}Br), 29.93], 218 [M^+ (^{79}Br), 30.23], 139 [$M^+ - \text{Br}$, 40.18], 81 [Br^+ , 75.48], 79 [Br^+ , 76.97], 43 [100] IR (neat) 1714.4 cm^{-1} ; Anal. Calcd. For $\text{C}_8\text{H}_{11}\text{BrO}_2$: C, 43.86; H, 5.06. Found: C, 43.62; H, 5.11.

3-*n*-propyl-4-iodo-5,6-dihydro-2H-pyran-2-one (4g): oil, ^1H NMR (400M Hz, CDCl_3) δ 4.28-4.31 (t, 2H, $J = 6.19$ Hz), 3.00-3.03 (t, 2H, $J = 6.09$ Hz), 2.50-2.54 (t, 2H, $J = 7.96$ Hz), 1.50-1.56 (m, 2H), 0.96-0.99 (t, 3H, $J = 7.35$ Hz) ^{13}C NMR (100M Hz, CDCl_3) δ 160.0, 138.4, 116.1, 66.5, 40.2, 38.0, 21.4, 13.9. EIMS m/z (relative intensity, %) 266 [M^+ , 25.41], 139 [$M^+ - \text{I}$, 100], 127 [I^+ , 12.33] IR (neat) 1714.0 cm^{-1} ; Anal. Calcd. For $\text{C}_8\text{H}_{11}\text{IO}_2$: C, 36.11; H, 4.17. Found: C, 36.32; H, 4.10

3-benzyl-4-bromo-5,6-dihydro-2H-pyran-2-one (4h): mp 46-47°C, ¹H NMR (400M Hz, CDCl₃) δ 7.19-7.33 (m, 5H), 4.33-4.36 (t, 2H, *J* = 6.28 Hz), 3.90 (s, 2H), 2.96-2.99 (t, 2H, *J* = 6.31 Hz), ¹³C NMR (100M Hz, CDCl₃) δ 161.7, 138.8, 137.3, 131.4, 128.2, 127.9, 126.0, 64.7, 35.8, 25.0. EIMS *m/z* (relative intensity, %) 268 [M⁺ (⁸¹Br), 13.02], 266 [M⁺ (⁷⁹Br), 12.99], 187 [M⁺ – Br, 40.52], 128 [100], 91 [Bn⁺, 27.58] IR (neat) 1718.3cm⁻¹; Anal. Calcd. For C₁₂H₁₁BrO₂: C, 53.96; H, 4.15. Found: C, 53.77; H, 4.22.

3-benzyl-4-iodo-5,6-dihydro-2H-pyran-2-one (4i): mp 49-51°C, ¹H NMR (400M Hz, CDCl₃) δ 7.1-7.35 (m, 5H), 4.28-4.31(t, 2H, *J* = 6.09 Hz), 3.95(s, 2H), 3.06-3.09(t, 2H, *J* = 6.15 Hz), ¹³C NMR (100M Hz, CDCl₃) δ 159.4, 137.2, 137.1, 128.1, 127.9, 126.1, 116.9, 65.8, 40.7, 39.7. EIMS *m/z* (relative intensity, %) 314[M⁺, 15.09], 187[M⁺ – I, 19.45], 128[94.56], 91[Bn⁺, 100] IR (neat) 1717.3cm⁻¹; Anal. Calcd. For C₁₂H₁₁IO₂: C, 45.88; H, 3.53. Found: C, 46.02; H, 3.49.

6-bromo-4,5,6,7-tetrahydro-3H-cyclopenta[c]-pyran-1-one (5a): oil, ¹H NMR (400M Hz, CDCl₃) δ 4.77-4.78 (m, 1H), 3.76-3.82(m, 2H), 3.54-3.61(m, 4H), 2.87-3.21(m, 2H) ¹³C NMR (100M Hz, CDCl₃) δ 165.1, 141.4, 128.2, 73.4, 38.4, 37.2, 34.4, 29.4. EIMS *m/z* (relative intensity, %) 218 [M⁺ (⁸¹Br), 6.63], 216 [M⁺ (⁷⁹Br), 6.84], 137[M⁺ – Br, 4.64], 65[100], IR (neat) 1754.3, 1652.7 cm⁻¹; Anal. Calcd. For C₈H₉BrO₂: C, 44.27; H, 4.18. Found: C, 44.09; H, 4.23.

6-iodo-4,5,6,7-tetrahydro-3H-cyclopenta[c]-pyran-1-one (5b): oil, ¹H NMR (400M Hz, CDCl₃) δ 4.56-4.58 (m, 1H), 3.84-3.88(m, 2H), 3.30-3.44(m, 4H), 2.68-3.07(m, 2H) ¹³C NMR (100M Hz, CDCl₃) δ 162.8, 133.6, 125.0, 73.4, 44.1, 42.2, 8.29, 2.85.

ELMS m/z (relative intensity, %) 264 [M^+ , 14.79], 137 [$M^+ - I$, 88.32], 127 [I^+ , 49.49],

91 [100] IR (neat) 1749.1, 1632.8 cm^{-1} ; Anal. Calcd. For $C_8H_9BrO_2$: C, 36.39; H, 3.44.

Found: C, 36.18; H, 3.48.