Novel and Direct Oxidative Cyanation Reactions of Heteroaromatic Compounds Mediated by A Hypervalent Iodine(III) Reagent

Toshifumi Dohi, Koji Morimoto, Yorito Kiyono, Hirofumi Tohma,
and Yasuyuki Kita*

*Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamada-oka, Suita,
Osaka 565-0871, Japan*

General Information ^1H NMR and ^{13}C NMR spectra were recorded on a JEOL JMN-300 spectrometer in CDCl_3 with tetramethylsilane as an internal standard. Data are reported as follows: chemical shift in ppm (δ), multiplicity (s = singlet, d = doublet, t = triplet, br = broad singlet, m = multiplet, c = complex), coupling constant (Hz), integration, and interpretation. Infrared spectra (IR) were obtained on a Hitachi 270-50 spectrometer; absorptions are reported in reciprocal centimeters with the following relative intensities: s (strong), m (medium), or w (weak). Mass spectra were obtained on a Shimadzu GCMS-QP 5000 instrument with ionization voltages of 70 eV. Elemental analyses and high resolution mass spectra were performed by the Elemental Analysis Section of Osaka University. Column chromatography was performed with SiO_2 (Merck Silica Gel 60 (230-400 mesh)). Preparative high performance liquid chromatography (HPLC) was performed on a Japan Analytical Industry LC-908 instrument, equipped with an RI absorbance detector.

Materials PIFA was prepared from commercially available phenyliodine bisacetate (PIDA) and trifluoroacetic acid.¹ Absolute dichloromethane, $\text{BF}_3 \cdot \text{Et}_2\text{O}$ and heteroaromatic compounds (**1a-d**, **3a-b**, **3d-e**) are commercially available and used without further purification. Other N-tosylated pyrroles (**1e-i**) and indoles (**5a-c**) were prepared from corresponding 1H-pyrroles and p-toluenesulfonylchloride. Pyrrole (**1f**, **g**)², thiophene (**3c**)³, indole (**6c**)⁴ were prepared according to the literature procedure.

3-Heptyl-1-(toluene-4-sulfonyl)-pyrrole (1e) Brown oil ; $R_f = 0.57$ (hexane/EtOAc = 4/1) ; ^1H NMR (CDCl_3): δ 0.84 (t, $J = 6.9$ Hz, 3H, CH_2CH_3), 1.07-1.33 (c, 8H, CH_2), 1.46 (m, 1H, ArCH_2CH_2), 2.32 (t, $J = 7.8$ Hz, 2H, ArCH_2), 2.35 (s, 3H, Ts), 6.10 (m, 1H, pyrrole 4-H), 6.85 (m, 1H, pyrrole 2-H), 7.02 (m, 1H, pyrrole 5-H), 7.23 (d, $J = 8.4$ Hz, 2H, Ts), 7.68 (d, $J = 8.4$ Hz, 2H, Ts). ^{13}C NMR (CDCl_3): δ 14.03, 21.51, 22.57, 26.69, 29.00, 29.10, 29.88, 31.74, 114.71, 117.14, 120.76, 126.65, 129.80, 130.01, 136.29, 144.58. IR (KBr): 3142 m, 2923 s, 2855 s, 1732 m, 1597 m, 1454 s, 1360 s, 1252 s, 1171 s, 1101 s, 1058 s, 956 m, 912 m, 798 m, 667 s cm^{-1} . HRFABMS Calcd for $\text{C}_{18}\text{H}_{26}\text{O}_2\text{NS}$ ($\text{M}+\text{H}$) $^+$; 320.1686. Found 320.1685.

3-(4-Bromophenyl)-1-(toluene-4-sulfonyl)-pyrrole (1f) White powder; $R_f = 0.55$ (hexane/EtOAc = 4/1) ; ^1H NMR (CDCl_3): δ 2.39 (s, 3H, Ts), 6.53 (m, 1H, pyrrole 4-H), 7.12-7.38 (c, 6H, Ar), 7.43 (d, $J = 8.7$ Hz, 2H, Ar) 7.76 (d, $J = 8.4$ Hz, 2H, Ts). ^{13}C NMR (CDCl_3): δ 21.62, 111.80, 116.37, 120.71, 121.79, 126.94, 127.11, 128.35, 130.08, 131.83, 132.45, 135.82, 145.24. MS: m/z (relative intensity, %): 377 (M^+ , 26), 375 (22), 222 (53), 220 (57), 194 (13), 192 (13), 155 (10), 113 (14), 91 (100), 64 (13). Anal. Calcd for $\text{C}_{17}\text{H}_{14}\text{BrNO}_2\text{S}$: C, 54.26; H, 3.75; N, 3.72. Found: C, 54.45; H, 3.87; N, 3.72.

3-(2-Bromophenyl)-1-(toluene-4-sulfonyl)-pyrrole (1g) Brown powder ; $R_f = 0.51$ (hexane/EtOAc = 4/1) ; ^1H NMR (CDCl_3): δ 2.40 (s, 3H, Ts), 6.55 (dd, $J = 3.4, 1.8$ Hz, 1H, pyrrole 4-H), 7.09 (td, $J = 7.8, 1.8$ Hz, H, Ar), 7.14 (m, 1H, Ar), 7.20-7.34 (c, 4H, Ar), 7.46 (m, 1H, Ar), 7.61 (dd, $J = 8.7$ Hz, 1.8 Hz, 1H, Ar), 7.77 (d, $J = 8.4$ Hz, 2H, Ts) ^{13}C NMR (CDCl_3): δ 21.67, 114.81, 119.39, 120.20, 122.14, 126.78, 127.30, 127.92, 128.31, 129.90, 130.63, 133.43, 134.48, 135.80, 144.98. IR (KBr): 3143 m, 3064 m, 2923 m, 1919 w, 1596 m, 1558 m, 1488 s, 1459 m, 1435 m, 1371 s, 1308 s, 1255 m, 1174 s, 1134 s, 1058 s, 958 m, 793 s, 757 s, 704 s, 674 s cm^{-1} . MS: m/z (relative intensity, %): 377 (M^+ , 38), 375 (38), 313 (3), 311 (3), 222 (22), 220 (22), 194 (10), 192 (10), 155 (38), 113 (23), 91 (100), 64 (22). HRFABMS Calcd for $\text{C}_{17}\text{H}_{14}\text{O}_2\text{NSBr}$ (M

+ Na)⁺; 397.9826. Found 397.9837.

General Procedure for PIFA-mediated Cyanation Reaction of Heteroaromatic Compounds. To a stirred solution of PIFA (2 mmol) and BF₃ • Et₂O (4 mmol) in CH₂Cl₂ (1 mL) added trimethylsilyl cyanide (3 mmol) at room temperature. After stirring 30 min., N-tosyl pyrrole **1d** (1 mmol) was added in one portion and stirred for additional 3 hours under the same condition while checking the reaction progress by GC or TLC. After the reaction completed, saturated aqueous sodium thiosulfate (ca. 5 mL) was added to the mixture. The organic layer was separated and aqueous phase was extracted with CH₂Cl₂. The combined extract was dried with MgSO₄ and evaporated to dryness. The residue was purified by column chromatography (SiO₂/n-hexane-Et₂O) to give pure **2d** (83%).

All reactions have finished within 3 hours. Compounds **2h-i**, **6a-b** were identified by comparing the reported data after deprotection of tolyloxy group.^{7, 8, 11}

1-Methyl-pyrrole-2-carbonitrile (2c).⁵ ¹H NMR (CDCl₃): δ 3.72 (s, 3H, CH₃), 6.10 (dd, *J* = 4.0, 2.0 Hz, 1H, pyrrole), 6.71 (m, 1H, pyrrole), 6.78 (dd, *J* = 4.0, 2.0 Hz, 1H, pyrrole). ¹³C NMR (CDCl₃): δ 35.2, 104.3, 109.4, 113.8, 119.8, 127.6. IR (KBr): 2235 (CN) cm⁻¹.

1-(toluene-4-sulfonyl)-pyrrole-2-carbonitrile (2d)⁶ White powder, mp 114-115 °C, *R_f* = 0.51 (hexane/EtOAc = 4/1); ¹H NMR (CDCl₃): δ 2.44 (s, 3H, Ts), 6.32 (t, *J* = 3.4 Hz, 1H, pyrrole), 6.95 (dd, *J* = 3.4, 1.6 Hz, 1H, pyrrole), 7.37 (d, *J* = 8.7 Hz, 2H, Ts), 7.47 (dd, *J* = 3.4, 1.6 Hz, 1H, pyrrole), 7.93 (d, *J* = 8.7 Hz, 2H, Ts). ¹³C NMR (CDCl₃): δ 21.69, 103.74, 111.63, 112.30, 126.56, 126.57, 127.85, 130.36, 134.12, 146.52. IR (KBr): 2225 s (CN) cm⁻¹.

3-Heptyl-1-(toluene-4-sulfonyl)-pyrrole-2-carbonitrile (2e) Brown oil; *R_f* = 0.41 (hexane/EtOAc = 4/1); ¹H NMR (CDCl₃): δ 0.75 (t, *J* = 7.5 Hz, 3H, CH₃), 1.02-1.30 (c, 8H, CH₂), 1.43 (m, 2H, ArCH₂CH₂), 2.34 (s, 3H, Ts) 2.41 (t, *J* = 7.3 Hz, 2H,

ArCH₂), 6.10 (d, *J* = 3.4 Hz, 1H, pyrrole 4-H), 7.20-7.30 (c, 3H, Ar), 7.81 (d, *J* = 8.4 Hz, 2H, Ts). ¹³C NMR (CDCl₃): δ 14.00, 21.68, 22.51, 26.56, 28.82, 28.93, 29.52, 31.61, 101.52, 111.71, 113.07, 126.33, 127.72, 130.26, 134.35, 144.14, 146.16. IR (KBr): 3144 w, 2928 m, 2856 m, 2220 s (C≡N), 1595 m, 1556 m, 1493 m, 1454 m, 1385 s, 1267 m, 1163 s, 1088 m, 1049 m, 970 m, 912 s, 814 m, 743 s, 675 m cm⁻¹. HRFABMS Calcd for C₁₉H₂₅N₂O₂S (M+H)⁺; 345.1637. Found 345.1636.

3-(4-Bromophenyl)-1-(toluene-4-sulfonyl)-pyrrole-2-carbonitrile (2f) White powder; *R_f* = 0.41 (hexane/EtOAc = 4/1); ¹H NMR (CDCl₃): δ 2.37 (s, 3H, Ts), 6.46 (d, *J* = 3.1 Hz, 1H, pyrrole 4-H), 7.31 (d, *J* = 8.7 Hz, 2H, Ts), 7.37-7.55 (c, 5H, Ar), 7.89 (d, *J* = 8.7 Hz, 2H, Ts). ¹³C NMR (CDCl₃): δ 21.78, 99.46, 111.57, 112.37, 123.34, 126.86, 128.09, 128.69, 129.75, 130.46, 132.20, 133.98, 139.89, 146.69. IR (KBr): 3145 m, 2926 w, 2220 s (C≡N), 1593 m, 1493 s, 1385 s, 1290 m, 1169 s, 1088 m, 1074 m, 1007 s, 912 m, 829 m, 814 m, 762 m, 739 s, 702 m, 648 m, 624 s cm⁻¹. MS: *m/z* (relative intensity, %): 401 (M⁺, 11), 399 (8), 247 (3), 245 (3), 195 (6), 194 (6), 155 (29), 113 (4), 91 (100), 194 (13), 64 (13). HRFABMS Calcd for C₁₈H₁₄O₂N₂BrS (M+H)⁺; 400.9959. Found 400.9944

3-(2-Bromophenyl)-1-(toluene-4-sulfonyl)-pyrrole-2-carbonitrile (2g) Colorless oil; *R_f* = 0.41 (hexane/EtOAc = 4/1); ¹H NMR (CDCl₃): δ 2.44 (s, 3H, Ts), 6.56 (d, *J* = 3.4 Hz, 1H, pyrrole 4-H), 7.21 (m, 1H, Ar), 7.27-7.45 (c, 4H, Ar), 7.50 (d, *J* = 3.4 Hz, 1H, pyrrole 5-H), 7.62 (d, *J* = 8.7 Hz, 1H, Ar), 7.96 (d, *J* = 8.4 Hz, 2H, Ts). ¹³C NMR (CDCl₃): δ 21.78, 102.05, 111.50, 114.47, 122.78, 125.63, 127.65, 128.02, 130.42, 131.26, 131.92, 133.49, 134.12, 140.68, 146.60. IR (KBr): 3143 m, 3060 m, 2923 m, 2223 s (CN), 1923 w, 1595 s, 1564 m, 1479 s, 1442 s, 1385 s, 1300 s, 1255 s, 1169 s, 1088 s, 1011 s, 930 m, 814 s, 758 s, 675 s cm⁻¹. HRFABMS Calcd for C₁₈H₁₄O₂N₂BrS (M+H)⁺; 400.9959. Found 400.9946

3,5-Dimethyl-1-(toluene-4-sulfonyl)-pyrrole-2-carbonitrile (2h)⁷ White powder; *R_f* = 0.38 (hexane/EtOAc = 4/1); ¹H NMR (CDCl₃): δ 2.08 (s, 3H, CH₃), 2.37 (s, 3H,

CH₃), 2.38 (s, 3H, CH₃), 5.80 (s, 1H, pyrrole 4-H), 7.28(d, $J = 8.4$ Hz, 2H, Ts), 7.80 (d, $J = 8.4$ Hz, 2H, Ts). ¹³C NMR (CDCl₃): δ 11.91, 14.95, 21.71, 112.55, 114.95, 126.99, 129.95, 130.79 134.95, 137.21, 137.54, 145.55. IR (KBr): 2926 m, 2855 w, 2214 s (CN), 1585w, 1504 m, 1445 m, 1377 s, 1254 s, 1180 s, 1153 m, 1090 m, 1051 m, 912 s, 814 m, 743 s, 673 m cm⁻¹. HRFABMS Calcd for C₁₄H₁₅O₂N₂S (M+H)⁺; 275.0884. Found 275.0869

3-Methyl-thiophene-2-carbonitrile (4a)⁹ Yellow oil; $R_f = 0.43$ (hexane/EtOAc = 4/1) ; ¹H NMR (CDCl₃): δ 2.45 (s, 3H, CH₃), 6.94 (d, $J = 5.1$ Hz, 1H, thiophene 4-H), 7.46 (d, $J = 5.1$ Hz, 1H, thiophene 5-H). IR (KBr): 2214 cm⁻¹(CN).

3-Hexyl-thiophene-2-carbonitrile (4b) Yellow oil; $R_f = 0.42$ (hexane/EtOAc = 4/1) ; ¹H NMR (CDCl₃): δ 0.89 (t, $J = 6.9$ Hz, 3H, CH₃), 1.20-1.48 (c, 6H, CH₂), 1.51-1.73 (m, 2H, ArCH₂CH₂), 2.79 (t, $J = 7.5$ Hz, 2H, ArCH₂), 6.97 (d, $J = 5.1$ Hz, 1H, thiophene 4-H), 7.26 (d, $J = 5.1$ Hz, 1H, thiophene 5-H). ¹³C NMR (CDCl₃): δ 14.01, 22.50, 28.76, 29.83, 30.13, 31.45, 105.42, 114.33, 128.49, 131.63, 154.81. IR (KBr): 3153 w, 3111 w, 2930 s, 2858 m, 2214 s (C N), 1794 w, 1529 w, 1468 m, 1414 m, 1379 m, 1215 m, 1163 w, 1092 m, 912 s, 743 s, 650 m cm⁻¹. HRFABMS Calcd for C₁₁H₁₆NS (M+H)⁺; 194.1047. Found 194.1025

3-Cyclohexyl-thiophene-2-carbonitrile (4c) Yellow oil; $R_f = 0.41$ (hexane/EtOAc = 4/1) ; ¹H NMR (CDCl₃): δ 1.05-1.98 (c, 10H, CH₂), 2.85 (m, 1H, CH), 6.93 (d, $J = 5.1$ Hz, 1H, thiophene 4-H), 7.40 (d, $J = 5.1$ Hz, 1H, thiophene 5-H). ¹³C NMR (CDCl₃): δ 25.75, 26.31, 33.62, 39.77, 104.23, 114.42, 126.52, 131.80, 159.91. IR (KBr): 3153 w, 3109 w, 2932 m, 2855 m, 2214 s (C N), 1526w, 1448 m, 1416 m, 1159 w, 1128 w, 1091 w, 912 s, 743 s, 650 m. HRFABMS Calcd for C₁₁H₁₄NS (M+H)⁺; 192.0847. Found 192.0824.

3-Methoxy-thiophene-2-carbonitrile (4d)¹⁰ colorless oil; $R_f = 0.39$ (hexane/EtOAc = 4/1) ; ¹H NMR (CDCl₃): δ 4.02 (s, 3H, CH₃), 6.72 (d, $J = 5.1$ Hz, 1H, thiophene 4-H), 7.36 (d, $J = 5.1$ Hz, 1H, thiophene 5-H). IR (KBr): 2214 s (CN).

3-Phenyl-thiophene-2-carbonitrile (4e) Yellow solid; $R_f = 0.42$ (hexane/EtOAc = 4/1) ; ^1H NMR (CDCl_3): δ 7.23 (d, $J = 5.1$ Hz, 1H, thiophene 4-H), 7.23-7.50 (c, 3H, Ar), 7.54 (d, $J = 5.1$ Hz, 1H, thiophene 5-H), 7.65 (dd, $J = 8.4, 1.8$ Hz, 1H, Ar). ^{13}C NMR (CDCl_3): δ 104.32, 114.73, 127.72, 128.23, 128.96, 129.08, 131.83, 132.99, 151.35. IR (KBr): 3034 w, 2214 s (C $\equiv$ N), 1493 m, 1448 m, 1402 m, 1379 m, 1157 m, 1101 w, 1074 w, 912 s, 883 m, 835 w, 743 s, 694 m, 654 m cm^{-1} . HRFABMS Calcd for $\text{C}_{11}\text{H}_8\text{NS}$ ($\text{M}+\text{H}$) $^+$; 186.0377. Found 186.0387.

2-Methyl-1-(toluene-4-sulfonyl)-indole-3-carbonitrile (6a)¹¹ White powder; $R_f = 0.31$ (hexane/EtOAc = 10/1) ; ^1H NMR (CDCl_3): 2.39 (s, 3H, Ts), 2.81 (s, 3H, CH_3), 7.22-7.46 (c, 4H, Ar), 7.59 (dd, $J = 7.1, 1.4$ Hz, 1H, Ar), 7.74 (d, $J = 8.4$ Hz, 2H, Ts), 8.20 (dd, $J = 7.1, 1.4$ Hz, 1H, Ar). ^{13}C NMR (CDCl_3): 15.00, 21.74, 113.98, 114.58, 119.10, 124.65, 125.73, 126.55, 126.84, 127.75, 129.69, 130.22, 135.12, 135.39, 145.99. IR (KBr): 3077 w, 2928 s, 2226 s (C $\equiv$ N), 1597 m, 1566 w, 1454 m, 1385 s, 1308 m, 1254 m, 1220m, 1180 s, 1120 m, 1089 m, 1022 m, 972 m, 912 s, 812 m, 745 s, 684 m, 665 s cm^{-1} .

3-Methyl-1-(toluene-4-sulfonyl)-indole-2-carbonitrile (6b)¹¹ Yellow powder; $R_f = 0.31$ (hexane/EtOAc = 4/1) ; ^1H NMR (CDCl_3): 2.39 (s, 3H, Ts), 2.81 (s, 3H, CH_3), 7.21-7.45 (c, 4H, Ar), 7.58 (dd, $J = 7.1, 1.4$ Hz, 1H, Ar), 7.74 (d, $J = 8.4$ Hz, 2H, Ts), 8.20 (dd, $J = 7.1, 1.4$ Hz, 1H, Ar). ^{13}C NMR (CDCl_3): 10.20, 21.73, 107.04, 112.04, 114.74, 120.57, 124.28, 127.02, 128.58, 128.65, 130.02, 134.25, 134.40, 136.58, 145.67. IR (KBr): 2932 s, 2226 s (CN), 1598 s, 1452 s, 1387 s, 1301 s, 1254 s, 1181 w, 1090 s, , 912 s, 743 s, 665 s cm^{-1} . HRFABMS Calcd for $\text{C}_{17}\text{H}_{15}\text{N}_2\text{O}_2\text{S}$ ($\text{M}+\text{H}$) $^+$; 311.0854. Found 311.0850.

Transformation of 2f and 2g into 2-oxazole-3-aryl pyrroles.

ZnCl_2 (0.2 mmol) was added to the mixture of **2f** or **2g** (0.2 mmol) and 2-amino-1-methyl-propanol (1 mmol) in chlorobenzene (1mL), and the mixture was

heated at reflux for 20 h. Then, solvent was removed in vacuo and the residue was chromatographed on silica gel (hexane/EtOAc) to give pure **7a** and **7b**.

2-[3-(4-Bromophenyl)-1H-pyrrol-2-yl]-4,4-dimethyl-4,5-dihydro-oxazole (7a)

White powder; $R_f = 0.24$ (hexane/EtOAc = 4/1) ; ^1H NMR (CDCl_3): 1.33 (s, 6H, CH_3), 4.00 (s, 2H, CH_2), 6.28 (d, $J = 2.7$ Hz, 1H, Pyrrole 4-H), 6.89 (d, $J = 2.7$ Hz, 1H, Pyrrole 5-H), 7.39 (d, $J = 8.7$ Hz, 2H, Ar), 7.46 (d, $J = 8.7$ Hz, 2H, Ar), 10.52 (bs, 1H, NH). ^{13}C NMR (CDCl_3): 28.33, 66.28, 79.35, 100.49, 111.51, 120.55, 121.37, 127.80, 130.72, 130.92, 134.36, 157.52. IR (KBr): 3450 w, 3176 w, 2970 m, 1643 m ($\text{C}=\text{N}$), 1593 w, 1546 w, 1502 m, 1427 m, 1359 w, 1248 w, 1175 w, 1117 m, 1011 w, 984 w, 912 s, 743 s, 650 w cm^{-1} . HRFABMS Calcd for $\text{C}_{15}\text{H}_{15}\text{ON}_2\text{Br}$ ($\text{M}+\text{H}$) $^+$; 319.0446. Found 319.0453.

2-[3-(2-Bromophenyl)-1H-pyrrol-2-yl]-4,4-dimethyl-4,5-dihydro-oxazole (7b)

White powder; $R_f = 0.31$ (hexane/EtOAc = 10/1) ; ^1H NMR (CDCl_3): 1.25 (s, 6H, CH_3), 3.85 (s, 2H, CH_2), 6.17 (d, $J = 2.7$ Hz, 1H, Pyrrole 4-H), 6.85 (d, $J = 2.7$ Hz, 1H, Pyrrole 5-H), 7.08 (td, $J = 7.8, 1.8$ Hz, 1H, Ar), 7.13-7.38 (c, 4H, Ar), 7.54 (d, $J = 7.8$ Hz, 1H, Ar), 10.50 (bs, 1H, NH). ^{13}C NMR (CDCl_3): 28.32, 66.01, 79.52, 104.32, 112.23, 116.99, 120.88, 124.49, 126.51, 128.19, 131.77, 132.25, 138.86, 154.96. IR (KBr): 3176 w, 2970 m, 1651 m ($\text{C}=\text{N}$), 1593 w, 1553 w, 1504 w, 1454 m, 1433 m, 1366 w, 1302 w, 1250 w, 1175 w, 1121 m, 1047 w, 1026 w, 983 m, 912 s, 743 s, 650 m cm^{-1} . HRFABMS Calcd for $\text{C}_{15}\text{H}_{15}\text{ON}_2\text{Br}$ ($\text{M}+\text{H}$) $^+$; 319.0446. Found 319.0459.

References

- 1) White, J. D.; Caravatti, G.; Kline, T. B.; Edstrom, E.; Rice, K. C.; Brossi, A.; *Tetrahedron* **1983**, 39, 2393.
- 2) Balasubramanian, T.; Strachan, J-P.; Boyle, P. D.; Lindsey, J. S. *J. Org. Chem.* **2000**, 65, 7919.
- 3) Bongini, A.; Barbarella, G.; Favaretto, L.; Sotgin, G.; Zambianchi, M.; Casarini, D.

- Tetrahedron* **2002**, 58, 10151.
- 4) Ketcha, D. M.; Lieurance, B. A.; Homan, D. F. J.; Gribble, G. W. *J. Org. Chem.* **1989**, 54, 4350.
- 5) Brueckner, C.; Xie, L. Y.; Dolphin, D. *Tetrahedron* **1998**, 54, 2021
- 6) Highes, T. V.; Cava, M. P. *J. Org. Chem.* **1999**, 64, 313.
- 7) Cheng, L.; Lightner, D. A. *Synthesis* **1999**, 46.
- 8) Elsom, J. *J. Chem. Soc. B.* **1970**, 79.
- 9) Terpstra, J. W.; Leusen, A. M. *J. Org. Chem.* **1986**, 51, 230.
- 10) Rossy, P. A.; Hoffmann, W.; Mueller, N. *J. Org. Chem.* **1980**, 45, 617.
- 11) Tamura, Y.; Adachi, M.; Kawasaki, T.; Yasuda, H.; Kita, Y. *J Chem. Soc., Perkin Trans. I* **1980**, 1132.