Olefination of Ketones Using A Gold(III)-Catalyzed Meyer-Schuster Rearrangement

Douglas A. Engel and Gregory B. Dudley*

Department of Chemistry and Biochemistry, Florida State University,

Tallahassee, Florida 32306

Supporting Information

CONTENTS:

General experimental procedures	S1-S4
Characterization (analytical) data	S5-S12
Copies of ^1H NMR and ^{13}C NMR spectra	S13-S38

GENERAL EXPERIMENTAL PROCEDURES

^1H -NMR and ^{13}C -NMR spectra were recorded on 300 MHz spectrometer using CDCl_3 as the deuterated solvent. The chemical shifts (δ) are reported in parts per million (ppm) relative to the residual CHCl_3 peak (7.26 ppm for ^1H NMR, 77.0 ppm for ^{13}C NMR). The coupling constants (J) were reported in Hertz (Hz). IR spectra were recorded on an FTIR spectrometer on NaCl discs. Mass spectra were recorded using chemical ionization (CI) or electron ionization (EI) techniques. Yields refer to isolated material judged to be $\geq 95\%$ pure by ^1H NMR spectroscopy following silica gel chromatography. All chemicals were used as received unless otherwise stated. Tetrahydrofuran (THF) was purified by passing through a column of activated alumina. The $n\text{-BuLi}$ solutions were titrated with menthol dissolved in tetrahydrofuran using 1,10-phenanthroline as the

indicator. The purifications were performed by flash chromatography using silica gel F-254 (230-499 mesh particle size).

Typical procedure for the preparation of phenyl propargylic alcohols 2a, 2b, 2c

(Table 1: entries 1-3).

To a THF solution (10 mL) of phenylacetylene (1.55 mL, 14.1 mmol) was added *n*-BuLi (4.6 mL, 11 mmol, 2.3 M) dropwise over 5 min at $-78\text{ }^{\circ}\text{C}$ under argon atmosphere. The solution was allowed to warm to $0\text{ }^{\circ}\text{C}$ over 1 h and held at $0\text{ }^{\circ}\text{C}$ for an additional 30 min. The solution was then re-cooled to $-78\text{ }^{\circ}\text{C}$ and 3-pentanone (0.75 mL, 7.1 mmol) was added in one portion. The solution was allowed to warm to room temperature over 1 h and held at room temperature for an additional 3 h. Saturated aqueous NH_4Cl solution was added to quench the reaction and the mixture was extracted with ethyl acetate. The organic layer was washed with water, saturated aqueous sodium bicarbonate, and brine. The organic layer was dried over MgSO_4 , filtered, and concentrated. The residue was purified using silica gel column chromatography (hexanes/ethyl acetate = 20/1 – 10/1) to give 3-ethyl-1-phenyl-pent-1-yn-3-ol (**2b**) in 100% yield (1.33 g).

Typical procedure for the preparation of ethoxy propargylic alcohols 2d, 2e, 2f, 2g

(Table 1: entries 4-8, Table 2: entries 1-8).

To a THF solution (7 mL) of ethyl ethynyl ether (0.7 g, ca. 40% by weight in hexanes, ca. 9 mmol) was added *n*-BuLi (1.5 mL, 3.4 mmol, 2.3 M) dropwise over 5 min at $-78\text{ }^{\circ}\text{C}$ under argon atmosphere. The solution was allowed to warm to $0\text{ }^{\circ}\text{C}$ over 1 h and held at $0\text{ }^{\circ}\text{C}$ for an additional 30 min. The solution was then re-cooled to $-78\text{ }^{\circ}\text{C}$ and pinacolone

(0.30 mL, 2.4 mmol) was added in one portion. The solution was allowed to warm to room temperature over 1 h and held at room temperature for an additional 3 h. Saturated aqueous NH₄Cl solution was added to quench the reaction and the mixture was extracted with ethyl acetate. The organic layer was washed with water, saturated aqueous sodium bicarbonate, and brine. The organic layer was dried over MgSO₄, filtered, and concentrated. The residue was purified using silica gel column chromatography (hexanes/ethyl acetate = 20/1 – 7/1) to give 1-ethoxy-3-methyl-3-*tert*-butyl-1-propyn-3-ol (**2g**) in 83% yield (0.34g).

Typical procedure for preparation of α,β -unsaturated esters 3a-3g

(Table 1: entries 1-8, Table 2: entries 1-7)

To a CH₂Cl₂ solution (8 mL) of 1-ethoxy-3-methyl-3-*tert*-butyl-1-propyn-3-ol (**2g**) in an open flask was added 95% ethanol (0.16 mL, 2.9 mmol) followed by AuCl₃ (35 mg, 0.12 mmol, fine powder). After 5 min, the reaction mixture was filtered through a plug of SiO₂ with the aid of an ethyl acetate/hexane mixture (1/7). The filtrate was concentrated and purified using silica gel column chromatography (hexanes/ethyl acetate = 50/1) to give 3,4,4-trimethyl-pent-2-enoic acid ethyl ester (**3g**) in 86% yield (85 mg).

Typical two step procedure for preparation of α,β -unsaturated esters 3d, 3f-3j

(Table 3: entries 1-6)

To a THF solution (2.6 mL) of ethyl ethynyl ether (0.13g, ca. 40% by weight in hexanes, ca. 2 mmol) was added *n*-BuLi (0.40 mL, 0.75 mmol, 2.0 M) dropwise over 5 min at –78 °C under argon atmosphere. The solution was allowed to warm to 0 °C over 1 h and

held at 0 °C for an additional 30 min. The solution was then re-cooled to –78°C and 2-adamantanone (75 mg, 0.5 mmol) was added in one portion. The solution was allowed to warm to room temperature over 1 h and held at room temperature for an additional 3 h. Saturated aqueous NH₄Cl solution was added to quench the reaction and the mixture was extracted with ethyl acetate. The organic layer was washed with water, saturated aqueous sodium bicarbonate, and brine. The organic layer was dried over MgSO₄, filtered, and concentrated. To the concentrated mixture in an open flask was added 6 mL of CH₂Cl₂, followed by 95% ethanol (0.14 mL, 2.5 mmol) and AuCl₃ (7.6 mg, .025 mmol, fine powder). [NOTE: 15.2 mg, 0.05 mmol of AuCl₃ was used for the preparation of **3j** and **3k**.] After 30 min, the reaction mixture was filtered through a plug of SiO₂ with the aid of an ethyl acetate/hexane mixture (1/7). The filtrate was concentrated and purified using silica gel column chromatography (hexanes/ethyl acetate = 50/1) to give adamantan-2-ylidene-acetic acid ethyl ester (**3f**) in 99% yield (109 mg) over two steps.

Analytical Data

3-ethyl-1-phenylpent-1-yn-3-ol (2b): light yellow oil; ^1H NMR (300 MHz, CDCl_3) δ 1.11 (t, $J = 7.4$ Hz, 6H), 1.77 (q d, $J = 7.4, 4.5$ Hz, 4H), 1.94 (br s, 1H), 7.28-7.33 (m, 3H), 7.40-7.44 (m, 2H). For full characterization refer to Galli, C. et al. *J. Org. Chem.* **1994**, 59, 6786-6795.

1,3-diphenylprop-2-yn-1-ol (2c): colorless oil; NMR (300 MHz, CDCl_3) δ 2.27 (d, $J = 8.1$ Hz, 1H), 5.70 (s, 1H), 7.31-7.50 (m, 8H), 7.62-7.65 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 64.6, 86.3, 88.9, 122.3, 126.6, 128.1, 128.3, 128.4, 131.5, 140.5; IR (neat) 3350, 3062, 2874, 2229, 1953, 1884, 1807, 1598, 1490, 1384, 1280, 1190, 1030, 917, 756, 691 cm^{-1} ; HRMS (CI) Calcd for $\text{C}_{15}\text{H}_{12}\text{O}$ ($\text{M}-\text{H}^+$) 207.0810. Found 207.0808.

3-ethoxy-1,1-diphenyl-prop-2-yn-1-ol (2d): yellow oil; ^1H NMR (300 MHz, CDCl_3) δ 1.41 (t, $J = 7.1$ Hz, 3H), 2.64 (s, 1H), 4.18 (q, $J = 7.1$ Hz, 2H), 7.21-7.34 (m, 6H), 7.60-7.63 (m, 4H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.4, 42.2, 74.4, 74.8, 95.9, 125.9, 127.2, 128.0, 146.1; IR (neat) 3543, 3444, 3060, 2982, 2263, 1450, 1170, 1004, 640 cm^{-1} ; HRMS (EI) Calcd for $\text{C}_{17}\text{H}_{16}\text{O}_2$ (M^+) 252.1150. Found 252.1153.

4-tert-butyl-1-ethoxyethynyl-cyclohexanol (2e): white solid; ^1H NMR (300 MHz, CDCl_3) δ 0.87 (s, 9H), 0.98 (t t, $J = 11.9, 3.5$ Hz, 1H), 1.38 (t, $J = 7.1$ Hz, 3H), 1.26-1.55 (m, 4H), 1.71 (br d, $J = 12.8$ Hz, 2H), 1.89 (s, 1H), 1.94 (br d, $J = 10.6$ Hz, 2H), 4.09 (q, $J = 7.1$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.3, 25.0, 27.6, 32.2, 41.1, 41.3, 47.1, 69.6, 74.4, 94.0; IR (neat) 3390, 2944, 2865, 2257, 1479, 1444, 1393, 1365, 1178, 1059, 900, 834 cm^{-1} ; HRMS (CI) Calcd for $\text{C}_{14}\text{H}_{24}\text{O}_2$ ($\text{M}+\text{H}^+$) 225.1855. Found 225.1856.

2-ethoxyethynyl-adamantan-2-ol (2f): clear, colorless oil; ^1H NMR (300 MHz, CDCl_3) δ 1.37 (t, $J = 7.1$ Hz, 3H), 1.53 (br s, 1H), 1.57 (br s, 1H), 1.68-1.80 (m, 7H), 1.88 (br s, 2H), 2.14 (br t, $J = 12.6$ Hz, 4H), 4.09 (q, $J = 7.1$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.6, 27.1, 27.3, 32.0, 35.9, 38.0, 39.8, 43.2, 72.7, 74.6, 93.9; IR (neat) 3426, 2902, 2854, 2666, 2259, 1712, 1450, 1247, 1010, 916, 866 cm^{-1} ; HRMS (CI) Calcd for $\text{C}_{14}\text{H}_{20}\text{O}_2$ ($\text{M}+\text{H}^+$) 221.1542. Found 221.1538.

1-ethoxy-3-methyl-3-tert-butyl-1-propyn-3-ol (2g): clear, colorless oil; ^1H NMR (300 MHz, CDCl_3) δ 1.03 (s, 9H), 1.37 (t, $J = 7.1$ Hz, 3H), 1.41 (s, 3H), 1.71 (s, 1H), 4.08 (q, $J = 7.1$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.3, 25.2, 25.6, 38.4, 41.9, 73.9, 74.2, 92.8; IR (neat) 3479, 2971, 2873, 2261, 1481, 1392, 1369, 1219, 1094, 1007, 908, 878 cm^{-1} ; HRMS (CI) Calcd for $\text{C}_{10}\text{H}_{19}\text{O}_2$ ($[\text{M}+\text{H}]^+$) 171.1385. Found 171.1390.

1,3,3-triphenylpropenone (3a): yellow oil; ^1H NMR (300 MHz, CDCl_3) δ 7.12 (s, 1H), 7.16-7.21 (m, 2H), 7.25-7.28 (m, 3H), 7.35-7.4 (m, 7H), 7.46-7.51 (m, 1H), 7.91 (d, J = 7.1 Hz, 2H); For full characterization refer to Gurdeep et al. *J. Chem. Soc. Perkin Trans. I*. **1985**, 1289-1294.

ethyl 3,3-diphenylpropenoate (3d): yellow oil; ^1H NMR (300 MHz, CDCl_3) δ 2.01 (t, J = 7.1 Hz, 3H), 4.95 (q, J = 7.1 Hz, 2H), 8.09-8.12 (m, 2H), 8.18-8.30 (m, 8H); For full characterization refer to Le Tadic-Biadatti et al. *J. Org. Chem.* **1997**, 62, 559-563 and Rathke, M. et al. *Synth. Commun.* **1990**, 20, 869-875.

4-tert-butylcyclohexylidene-acetic acid ethyl ester (3e): clear, colorless oil; ^1H NMR (300 MHz, CDCl_3) δ 0.84 (s, 9H), 1.04-1.28 (m, 3H), 1.26 (t, $J = 7.1$ Hz, 3H), 1.76-1.95 (m, 3H), 2.14 (t d, $J = 13.5, 3.9$ Hz, 1H), 2.27-2.32 (m, 1H), 3.86 (d q, $J = 13.7, 2.7$, 1H), 4.12 (q, $J = 7.1$ Hz, 2H), 5.58 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.3, 27.5, 28.4, 29.2, 29.5, 32.4, 37.9, 47.8, 59.4, 112.7, 163.5, 166.9; IR (neat) 3412, 2948, 2867, 1716, 1648, 1478, 1365, 1247, 1143, 1041, 862 cm^{-1} ; HRMS (CI) Calcd for $\text{C}_{14}\text{H}_{24}\text{O}_2$ ($\text{M}+\text{H}^+$) 225.1855. Found 225.1849.

adamantan-2-ylidene-acetic acid ethyl ester (3f): clear, colorless oil: ^1H NMR (300 MHz, CDCl_3) δ 1.27 (t, $J = 7.1$ Hz, 3H), 1.86 (br s, 6H), 1.93-1.96 (m, 6H), 2.43 (br s, 1H), 4.07 (br s, 1H), 4.13 (q, $J = 7.1$ Hz, 2H), 5.58 (s, 1H); For full characterization refer to L. Griaud et al. *Tetrahedron*, **1998**, 54, 11899-11906.

(E/Z)-3,4,4-trimethyl-pent-2-enoic acid ethyl ester (3g): clear, colorless oil: ^1H NMR (300 MHz, CDCl_3 , E isomer) δ 1.10 (s, 9H), 1.28 (t, $J = 7.1$ Hz, 3H), 2.16 (br d, $J = 1.1$ Hz, 3H), 4.14 (q, $J = 7.1$ Hz, 2H), 5.74 (q, $J = 1.1$ Hz, 1H); ^1H NMR (300 MHz, CDCl_3 , Z isomer) δ 1.20 (s, 9H), 1.28 (t, $J = 7.1$ Hz, 3H), 1.84 (br d, $J = 1.3$ Hz, 3H), 4.14 (q, $J = 7.1$ Hz, 2H), 5.63 (q, $J = 1.3$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3 , E/Z mixture) δ 14.1, 14.3, 15.1, 23.9, 28.5, 29.0, 36.4, 37.9, 59.4, 60.0, 112.9, 116.6, 158.5, 167.2, 167.5, 167.9; IR (neat, E/Z mixture) 2970, 2873, 1719, 1634, 1466, 1372, 1262, 1182, 1123, 1054, 868 cm^{-1} ; HRMS (EI) Calcd for $\text{C}_{10}\text{H}_{18}\text{O}_2$ (M^+) 170.1307. Found 170.1306.

3-butyl-hept-2-enoic acid ethyl ester (3h): clear, colorless oil: ^1H NMR (300 MHz, CDCl_3) δ 0.89-0.94 (m, 6H), 1.27 (t, $J = 7.1$ Hz, 3H), 1.28-1.47 (m, 8H), 2.13 (br t, $J = 7.6$ Hz, 2H), 2.59 (br t, $J = 7.6$ Hz, 2H), 4.13 (q, $J = 7.1$ Hz, 2H), 5.61 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 13.88, 13.92, 14.3, 22.4, 23.0, 29.8, 30.8, 31.9, 38.1, 59.3, 115.0, 164.8, 166.6; IR (neat) 2958, 2931, 2872, 1716, 1642, 1466, 1378, 1190, 1147, 1039, 862 cm^{-1} ; HRMS (CI) Calcd for $\text{C}_{13}\text{H}_{24}\text{O}_2$ ($\text{M}+\text{H}^+$) 213.1855. Found 213.1857.

(4,6,6-trimethyl-bicyclo[3.1.1]hept-3-en-(2E/Z)-ylidene)-acetic acid ester (3i): clear oil; ^1H NMR (300 MHz, CDCl_3 , E isomer) δ 0.86 (s, 3H), 1.28 (t, $J = 7.1$ Hz, 3H), 1.40 (s, 3H), 1.68 (d, $J = 7.9$ Hz, 1H), 1.90 (d, $J = 1.5$ Hz, 3H), 2.20-2.45 (m, 1H), 2.53-2.63 (m, 2H), 4.08-4.20 (m, 2H), 5.32 (s, 1H), 7.13 (s, 1H); ^1H NMR (300 MHz, CDCl_3 , Z isomer) δ 0.84 (s, 3H), 1.26 (t, $J = 7.1$ Hz, 3H), 1.44 (s, 3H), 1.58 (d, $J = 8.8$ Hz, 1H), 1.86 (d, $J = 1.4$, 3H), 2.20-2.45 (m, 1H), 2.53-2.63 (m, 2H), 4.08-4.20 (m, 2H), 5.46 (s, 1H), 5.77 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3 , E/Z mixture) δ 14.3, 14.4, 21.7, 21.8, 23.2, 23.6, 26.4, 26.5, 37.5, 38.1, 45.3, 47.8, 48.2, 49.0, 49.1, 53.1, 59.2, 59.3, 107.6, 110.0, 117.6, 121.6, 156.9, 158.0, 159.6, 161.2, 166.8, 167.4; IR (neat) 2979, 2930, 2870, 1708, 1622, 1466, 1443, 1380, 1370, 1226, 1164, 1040, 874, 705 cm^{-1} ; HRMS (EI) Calcd for $\text{C}_{14}\text{H}_{20}\text{O}_2$ (M^+) 220.1463. Found 220.1462.

(1,7,7-trimethyl-bicyclo[2.2.1]hept-(2E/Z)-ylidene)-acetic acid ethyl ester (3j): clear, colorless oil; ^1H NMR (300 MHz, CDCl_3) diagnostic resonance signals for E and Z isomers appeared at δ 5.58 (br t, $J = 2.3$ Hz, 1H), 5.64 (br t, $J = 2.0$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3 , E/Z mixture) δ 12.7, 13.2, 14.5, 14.6, 19.0, 19.1, 19.8, 20.2, 27.5, 27.8, 34.0, 34.4, 38.6, 40.6, 44.0, 44.6, 48.2, 50.0, 53.9, 54.0, 59.6, 60.0, 108.2, 112.2, 166.9, 167.6, 175.6; IR (neat, E/Z mixture) 2955, 2874, 1714, 1652, 1450, 1368, 1306, 1288, 1200, 1177, 1097, 1048, 873, 850 cm^{-1} ; HRMS (CI) Calcd for $\text{C}_{14}\text{H}_{22}\text{O}_2$ ($\text{M}+\text{H}^+$) 223.1698. Found 223.1695.

(3,3,5,5-tetramethyl-cyclohexylidene)-acetic acid ethyl ester (3k): clear, colorless oil; ^1H NMR (300 MHz, CDCl_3) δ 0.95 (s, 6H), 0.97 (s, 6H), 1.28 (t, $J = 7.1$ Hz, 3H), 1.33 (s, 2H), 1.95 (s, 2H), 2.61 (s, 2H), 4.15 (q, $J = 7.1$ Hz, 2H), 5.69 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.3, 30.8, 30.9, 34.8, 34.9, 42.2, 51.1, 52.5, 59.4, 115.8, 160.4, 166.6; IR (neat) 2953, 2896, 1716, 1646, 1461, 1376, 1229, 1154, 1043, 862 cm^{-1} ; HRMS (EI) Calcd for $\text{C}_{14}\text{H}_{24}\text{O}_2$ (M^+) 224.1776. Found 224.1771.