Oligosaccharide Synthesis in Microreactors

Frédéric R. Carrel, Karolin Geyer, Jeroen D.C. Codée, and Peter H. Seeberger

*Laboratory of Organic Chemistry, Swiss Federal Institute of Technology (ETH) Zürich,
Wolfgang-Pauli-Strasse 10, 8093 Zürich, Switzerland
E-mail: seeberger@org.chem.ethz.ch*

Supporting Information

Experimental Section.....	S2
References.....	S13
¹ H, ¹³ C, ¹⁹ F, ³¹ P - NMR Spectra.....	S14

Experimental Section

General methods. All chemicals were reagent grade and were used as supplied unless otherwise noted. All reactions were performed in oven-dried glassware under an inert atmosphere (nitrogen or argon) unless noted otherwise. Dichloromethane (CH_2Cl_2) and tetrahydrofuran (THF) were purified by a J. C. Meyer Solvent Dispensing System (two packed columns of neutral alumina). Solvents for chromatography and work-up procedures were distilled from commercially available technical grade solvents. Analytical thin-layer chromatography was performed on E. Merck silica gel 60 F₂₅₄ plates (0.25 mm). Compounds were visualized by UV and/or by dipping the plates in a cerium sulphate-ammonium molybdate solution followed by heating. Liquid column chromatography was performed using forced flow of the indicated solvent on Fluka silica gel 60 (40-63 μm). ^1H -, ^{13}C -, ^{19}F -, and ^{31}P -NMR spectra were recorded on a Varian Mercury XL 300 spectrometer. The ^1H -NMR spectra (300 MHz) are expressed in ppm relative to CHCl_3 (7.26 ppm) as internal reference, the coupling constants are reported in Hz. The same is valid for ^{13}C -NMR spectra (75 MHz, internal reference CDCl_3 : 77.0 ppm). For ^{19}F -NMR (282 MHz) and ^{31}P -NMR spectra (121 MHz) spectra, CCl_3F ($\delta = 0$ ppm) and H_3PO_4 ($\delta = 0$ ppm) were used as internal references. Optical rotations $[\alpha]_D$ were recorded on a Jasco DIP-370 spectrometer using a sodium lamp ($\lambda = 589$ nm) at room temperature with a 10cm/1mL cell. The solvent and the concentration are specified, e.g. $c = 1 = 10$ mg/mL. IR spectra were recorded in CHCl_3 on a Perkin-Elmer 1600 FT-IR spectrometer and are expressed in cm^{-1} . LC MS spectra were obtained using a Agilent 1100 LC MSD high-performance liquid chromatograph with a Waters Symmetry[®] C18 column (3.9 x 150 nm, 5 μm), and a gradient of acetonitrile/*iso*-propanol (80/20) and water/*iso*-propanol (80/20) as the mobile phase (flow rate 1 mL min^{-1} ; if necessary, 0.1% TFA was added to the eluents). The spectra were detected at 208, 234, 254 and 280 nm. Preparative HPLC was performed using a Waters HPLC apparatus with a Waters SunFireTM Prep C8 column (10 x 150 mm, 5 μm), and a gradient of acetonitrile/*iso*-propanol (80/20, 0.1% TFA) and water/*iso*-propanol (80/20, 0.1% TFA) as the mobile phase (flow rate 5 $\text{mL}\cdot\text{min}^{-1}$). The spectra were detected at a wavelength of 208 and 254 nm. High-resolution mass spectroscopy (HRMS) was performed by the MS-service at the Laboratory of Organic Chemistry at ETH Zürich; 2,5-dihydroxybenzoic acid (DHB) was used as matrix.

Dibutyl (3,4-di-*O*-benzyl-6-*O*-[fluorenylmethoxycarbonyl]-2-*O*-pivaloyl- β -D-glucopyranosyl) phosphate (1) was synthesized according to a published procedure.¹

8,8,9,9,10,10,11,11,12,12,13,13,14,14,15,15,15-Heptafluoropentadec-4(*Z*)-en-1-ol (2): To a solution of **7** (647 mg, 0.835 mmol) in THF (12.5 mL) was added HCl (1M in H₂O, 4.2 mL, 4.2 mmol) at room temperature. The solution was refluxed for 20 h, and then allowed to cool to room temperature. The reaction mixture was diluted with EtOAc (100 mL) and treated with a saturated aqueous NaHCO₃ solution (40 mL). The aqueous phase was extracted with EtOAc (2 x 50 mL), and the combined organic phases were dried over MgSO₄ and concentrated. The residue was purified by silica gel flash column chromatography (EtOAc/cyclohexane, gradient 1:9 to 1:2) to afford **2** (352 mg, 79%) as a white solid. *R*_f 0.25 (EtOAc/cyclohexane = 1:4); ¹H-NMR (300 MHz, CDCl₃) δ 5.49 (dddd, *J* = 10.9, 7.1, 7.1, 1.3, 1.3 Hz, 1H), 5.37 (ddd, *J* = 10.9, 7.1, 7.1 Hz, 1H), 3.66 (t, *J* = 6.5 Hz, 2H), 2.42–2.31 (m, 2H, CH₂), 2.21–2.01 (m, 4H), 1.64 (quint., *J* = 6.5 Hz, 2H), 1.40 (bs, 1H); ¹³C-NMR (75 MHz, CDCl₃) δ 131.2, 126.8, 62.2, 30.8 (t, *J* = 22 Hz), 32.2, 23.3, 18.1; ¹⁹F-NMR (282 MHz, CDCl₃) δ -80.7 (t, *J* = 9.6 Hz, 3F), -114.4 - -114.7 (quint., 2F), -121.4 - -122.1 (m, 6F), -122.7 (bs, 2F), -123.5 (bs, 2F), -126.1 (bs, 2F); IR (CHCl₃): 3618, 3443, 3008, 2939, 1448, 1200, 1152, 1113, 1060, 973, 876 cm⁻¹; HRMS-ESI Calcd for C₁₅H₁₁F₁₇ 514.0584 (M-H₂O)⁺; Found: 514.0581.

4-Trityloxybutan-1-ol (3). To a vigorously stirred solution of 1,4-butanediol (27.3 g, 300 mmol) in CH₂Cl₂ (300 mL) were added TrCl (8.363 g, 30 mmol) and pyridine (4.83 mL, 60 mmol) at room temperature. After 90 min, the solution was washed with brine (3 x 50 mL). The organic phase was dried over MgSO₄ and the solvents were evaporated in *vacuo*. The residue was purified by silica gel flash column chromatography (EtOAc/cyclohexane = 1:2) to afford **3** (7.90 g, 79%) as a white solid. Characterization data were consistent with the previously reported data.²

1-Iodo-4-trityloxybutane (4). To a solution of **3** (7.9 g, 23.7 mmol) in Et₂O (90 mL) and CH₃CN (30 mL) were added imidazole (4.85 g, 71.3 mmol) and triphenylphosphine (9.35 g, 35.6 mmol) at room temperature. The solution was cooled to 0 °C and iodine (9.05 g, 35.6 mmol) was added in portions over 5 min. After 30 min at 0 °C, the solution was allowed to warm to room temperature overnight. After 16 h at room temperature, the solution was diluted with Et₂O (300 mL). The precipitate was filtered off and washed with Et₂O (2 x 50 mL). The

filtrate was washed with a saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ solution (2 x 50 mL), and dried over MgSO_4 . The solvents were evaporated in *vacuo* and the residue was purified by silica gel flash column chromatography (EtOAc/cyclohexane = 1:19) to afford **4** (8.66 g, 82%) as a white solid. Characterization data were consistent with the previously reported data.²

(4-Trityloxybutan)-triphenylphosphonium iodide (5). To a solution of **4** (8.66 g, 19.58 mmol) in benzene (80 mL) was added triphenylphosphine (10.27 g, 39.16 mmol) at room temperature. The solution was heated under reflux for 17 h, and then allowed to cool to room temperature. The phosphonium salt was filtered off, washed with benzene (3 x 30 mL), and dried under vacuum to afford **5** (9.392 g, 68%) as a white solid. ^1H -NMR (300 MHz, CDCl_3) δ 7.80–7.60 (m, 15H), 7.30–7.16 (m, 15H), 3.60–3.46 (m, 2H), 3.12 (t, J = 5.6 Hz, 2H), 2.00 (quint., J = 5.9 Hz, 2H), 1.86–1.74 (m, 2H); ^{13}C -NMR (75 MHz, CDCl_3) δ 143.7, 135.0, 133.5 (d, J = 10 Hz), 130.4 (d, J = 10 Hz), 128.4, 127.7, 126.9, 117.9 (d, J = 85 Hz), 86.5, 61.8, 29.8 (d, J = 16 Hz), 22.4 (d, J = 51 Hz), 19.7; ^{31}P -NMR (121 MHz, CDCl_3) δ 24.3; IR (CHCl_3): 3391, 3087, 3060, 3007, 2942, 2876, 1969, 1818, 1589, 1488, 1439, 1390, 1316, 1200, 1152, 1113, 1072, 1034, 998, 900 cm^{-1} ; HRMS-MALDI Calcd for $\text{C}_{41}\text{H}_{38}\text{OP}$ 577.2655 (M-I)⁺; Found: 577.2648.

4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11,11-Heptadecafluoro-undecan-1-al (6). To a solution of 3-(perfluorooctyl)propan-1-ol (4.78 g, 10 mmol) in CH_2Cl_2 (200 mL) was added Dess-Martin periodinane (4.24 g, 10 mmol) at room temperature. After 30 min, further Dess-Martin periodinane (2.12 g, 5 mmol) was added and the solution was stirred for another 30 min. The solution was poured into a saturated aqueous NaHCO_3 / $\text{Na}_2\text{S}_2\text{O}_3$ solution (150 mL) and stirred for 30 min. The aqueous phase was extracted with CH_2Cl_2 (200 mL). The combined organic phases were dried over MgSO_4 and silica gel (20 g) was added. The solvents were evaporated in *vacuo* and the adsorbed residue was purified by silica gel flash column chromatography (EtOAc/cyclohexane = 1:2) to afford **6** (3.502 g, 73%) as a colorless liquid. Characterization data were consistent with the previously reported data.³

1,1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8-Heptadecafluoro-15-trityloxy-pentadec-11(Z)-ene (7). To a solution of **3** (7.47 g, 10.6 mmol) in THF (50 mL) was added BuLi (1.6 M in hexane, 6.6 mL, 10.6 mmol) at -30 °C. The orange solution was stirred for 15 min at -30 °C and a solution of **4** (2.52 g, 5.3 mmol) in THF (10 mL) was added. The resulting solution was stirred for 15 min at -30 °C and then allowed to warm to room temperature for 30 min. The solution was

diluted with CH₂Cl₂ (200 mL) and treated with a saturated aqueous NaHCO₃ solution (50 mL). The aqueous phase was extracted with CH₂Cl₂ (2 x 100 mL). The combined organic phases were dried over MgSO₄ and the solvents were evaporated in *vacuo*. The residue was purified by silica gel flash column chromatography (CH₂Cl₂ pure) to afford **7** (3.68 g, 90%) as a yellowish oil. R_f 0.78 (EtOAc/cyclohexane = 1:9); ¹H-NMR (300 MHz, CDCl₃) δ 7.50–7.44 (m, 6H), 7.36–7.23 (m, 9H), 5.46 (dddd, *J* = 11.0, 7.1, 7.1, 1.3, 1.3 Hz, 1H), 5.33 (ddd, 1H, *J* = 11.0, 7.1, 7.1 Hz, 1H), 3.17 (t, *J* = 6.6 Hz, 2H), 2.38–2.28 (m, 2H), 2.23–1.98 (m, 4H), 1.77–1.66 (m, 2H); ¹³C-NMR (75 MHz, CDCl₃) δ 144.3, 131.6, 128.6, 127.6, 126.8, 126.4, 86.3, 62.9, 30.8 (t, *J* = 22 Hz), 29.8, 23.9, 18.1; ¹⁹F-NMR (282 MHz, CDCl₃) δ -80.7 (t, *J* = 9.9 Hz, 3F), -114.4- -114.6 (quint., 2F), -121.4- -122.0 (m, 6F), -122.6 (bs, 2F), -123.4 (bs, 2F), -126.0 (bs, 2F); IR (CHCl₃): 3032, 2943, 1601, 1490, 1448, 1200, 1152 cm⁻¹; HRMS-ESI Calcd for C₂₈H₂₂F₁₇O 697.1594 (M-C₆H₅)⁺; Found: 697.1590.

Optimization of Glycosylation Conditions Using the Microreactor: General Procedure.

Prior to introduction of the reagents, the microreactor was rinsed with 20 - 50 reactor volumes of each anhydrous tetrahydrofuran and the anhydrous solvent used in the reaction. Reagent solutions were prepared by azeotroping the building blocks separately with toluene and drying under high-vacuum. The reagents were diluted with anhydrous solvent to the desired concentrations (Table 2). The gastight syringes were flushed with anhydrous solvents before loading the reagents. Inlet 1 was used for the nucleophile (2.5 mL syringe), inlet 2 for the glycosylating agent **1** (2.5 mL syringe), inlet 3 for the activator TMSOTf (1.0 mL syringe) and inlet 4 for a mixture of piperidine/DMF (1:4), also containing methyl 2,3,4,6-tetra-*O*-benzyl-α-D-glucopyranose [0.06 M] for LC MS analysis and TBAF. Inlet 3 was preconditioned with 1 mL activator solution. After loading, the syringes were connected to the device and the microreactor was set to the desired temperature by an ice bath. A pre-quenched volume of 100 μL was discarded at the desired flow rate to flush the device, followed by the collection of 3 aliquots of 25 μL of reaction mixture diluted with 1 mL acetonitrile for analysis. Seven flow rates were screened (469.8 μL·min⁻¹, 234.9 μL·min⁻¹, 156.6 μL·min⁻¹, 78.3 μL·min⁻¹, 39.15 μL·min⁻¹, 15.66 μL·min⁻¹ and 7.8 μL·min⁻¹), corresponding to reaction times of 10 s, 20 s, 30 s, 60 s, 120 s, 300 s and 600 s. The results depicted in Graphs 2-8 are summarized in Table 1.

Table 1

Entry	Nucleophile (1 equiv.)	Building Block 1 (equiv.); 0.24 M	TMSOTf (equiv.); 0.6M	Solvent	Temp [°C]	TBAF (equiv.); 0.08M	Time [s]	Product
1	2 ; 0.12M	2	2	TFT	0	-	60	8*
2	2 ; 0.12M	2	2	TFT	20	-	30	8*
3	8 ; 0.12M	2	2	CH ₂ Cl ₂	20	-	20	9*
4a	9 ; 0.12M	2	2	CH ₂ Cl ₂	20	-	30	10* , residual 9
4b	9 ; 0.12M	2	2	CH ₂ Cl ₂	20	2**	30	10 , residual 9
5	9 ; 0.08M	3	3	CH ₂ Cl ₂	20	2	60	10
6	10 ; 0.08M	3	3	CH ₂ Cl ₂	20	2	60	11

* Contaminated with the C(6)-TMS derivative. ** TBAF was added to the combined aliquots before FSPE

Graph 1: 1st glycosylation at temperature of 0 °C and 20 °C

Graph2

Graph3

Graph4

Graph5

Graph6

Graph7

(only one data point, since small amounts)

General Procedure for Scale-Out Using the Microreactor.

Microreactor and gastight syringes were prepared according to the screening procedure. The building blocks were coevaporated with toluene and dried under high-vacuum overnight. Both reactants were dissolved in the solvent of the reaction to give solutions with defined concentrations (Table 2, optimized reaction conditions). TMSOTf was dissolved in the solvent of the reaction to the corresponding defined solution, TBAF was dissolved in a mixture of piperidine/DMF = 1:4 (v/v). After flushing the microreactor and the syringes, the gastight glass syringes were filled with the corresponding reagent solutions and connected to the microreactor system. The reaction was carried out at the optimal reaction temperature and reaction time, whereby the first 300 μ L of pre-quenched volume were discarded, until the stock solutions were consumed. The solvents were evaporated in *vacuo* and residual DMF and piperidine were coevaporated with toluene (3 x 5 mL). The residue was purified by FSPE (“Si-Tridecafluoro, Silicycle”, loading with DMF/H₂O = 9:1), using the eluent MeOH/H₂O = 8:2 for non-fluorinated fractions and MeOH for the fluorinated fractions. The methanol fractions were evaporated in *vacuo* and residual solvents were coevaporated with toluene/methanol in *vacuo*. The residue was dissolved in CH₂Cl₂ and dried over MgSO₄. After evaporation of the solvents in *vacuo*, the corresponding glycoside was obtained as a white foam and was used for the next glycosylation.

Table 2

Entry	Nucleophile (1 equiv.)	Building Block 1 (equiv.); 0.24M	TMSOTf (equiv.); 0.6M	Solvent	TBAF (equiv.)	Temp [°C]	time [s]	Product (yield)
1	2 ; 0.12M	2	2	TFT	1.5; 0.09M	20	30	8 (99%)
2	8 ; 0.12M	2	2	CH ₂ Cl ₂	1.5; 0.09M	20	20	9 (97%)
3	9 ; 0.08M	3	3	CH ₂ Cl ₂	2; 0.08M	20	60	10 (90%)
4	10 ; 0.08M	3	3	CH ₂ Cl ₂	2; 0.08M	20	60	11 (95%)

[8,8,9,9,10,10,11,11,12,12,13,13,14,14,15,15,15-Heptadecafluoropentadec-4(Z)-en-1-yl] 3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranoside (8). General procedure using building blocks **2** (133 mg, 0.25 mmol, 0.12M in TFT), **1** (430 mg, 0.50 mmol, 0.24M in TFT), TMSOTf (90 μL, 0.50 mmol, 0.6M in TFT) and TBAF (1M in THF, 0.375 mmol, 0.09M in piperidine/DMF = 1:4) at +20 °C and 30 s reaction time gave a total pre-quenched volume of 4.28 mL (0.214 mmol) which afforded **8** (203 mg, 99%) as a white foam. *R_f* 0.63 (EtOAc:cyclohexane = 1:2); $[\alpha]_D^{25} = -6.6^\circ$ (*c* = 1, CHCl₃); ¹H-NMR (300 MHz, CDCl₃) δ 7.36–7.22 (m, 10H), 5.50–5.31 (m, 2H), 5.04 (dd, *J* = 9.1, 8.0 Hz, 1H), 4.82 (d, *J* = 11.0 Hz, 1H), 4.78 (d, *J* = 11.0 Hz, 1H), 4.71 (d, *J* = 11.0 Hz, 1H), 4.63 (d, *J* = 11.0 Hz, 1H), 4.43 (d, *J* = 8.0 Hz, 1H), 3.89 (dd, *J* = 12.4, 2.8 Hz, 1H), 3.85 (ddd, *J* = 9.3, 6.6, 6.6 Hz, 1H), 3.74 (dd, *J* = 9.0, 9.0 Hz, 1H), 3.73 (dd, *J* = 12.4, 4.7 Hz, 1H), 3.67 (dd, *J* = 9.0, 9.0 Hz, 1H), 3.50–3.38 (m, 2H), 2.40–2.29 (m, 2H), 2.21–2.01 (m, 4H), 1.90 (dd, *J* = 7.4, 6.1 Hz, 1H), 1.64 (quint., *J* = 6.8 Hz, 2H), 1.19 (s, 9H); ¹³C-NMR (75 MHz, CDCl₃) δ 176.6, 137.9, 137.7, 131.0, 128.4, 128.3, 128.0, 127.3, 127.9, 127.6, 126.8, 101.2, 82.9, 77.7, 75.2, 75.0, 74.9, 72.9, 69.2, 61.8, 38.7, 30.8 (t, *J* = 22 Hz), 29.3, 27.0, 23.5, 18.1; ¹⁹F-NMR (282 MHz, CDCl₃) δ -80.6 (t, *J* = 9.6 Hz, 3F), -114.3- -114.6 (m, 2F), -121.4- -122.0 (m, 6F), -122.6 (bs, 2F), -123.4 (bs, 2F), -126.0 (m, 2F); IR (CHCl₃): 3600, 3065, 3008, 2961, 2874, 1737, 1602, 1497, 1479, 1455, 1397, 1365, 1200, 1152, 1087, 1039, 866 cm⁻¹; HRMS-MALDI Calcd for C₄₀H₄₃F₁₇O₇Na 981.2630 (M+Na)⁺; Found: 981.2627.

[8,8,9,9,10,10,11,11,12,12,13,13,14,14,15,15,15-Heptadecafluoropentadec-4(Z)-en-1-yl] 6-O-[3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranosyl]-3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranoside (9). General procedure using building blocks **8** (194 mg, 0.20 mmol, 0.12M in CH₂Cl₂), **1** (348 mg, 0.40 mmol, 0.24M in CH₂Cl₂), TMSOTf (73 μL, 0.40 mmol, 0.6M in CH₂Cl₂) and TBAF (1M in THF, 0.30 mmol, 0.09M in piperidine/DMF = 1:4) at +20 °C and 20 s reaction time gave a total pre-quenched volume of 3.53 mL (0.176 mmol) which afforded **9** (238 mg, 97%) as a white foam. *R_f* 0.67 (EtOAc/cyclohexane = 1:2); $[\alpha]_D^{25} = -7.2^\circ$ (*c* = 0.5, CHCl₃); ¹H-NMR (300 MHz, CDCl₃) δ 7.37–7.22 (m, 20H), 5.51–5.30 (m, *J* = 10.7, 7.1 Hz, 2H), 5.06 (dd, *J* = 9.0, 7.8 Hz, 1H), 5.02 (dd, *J* = 9.3, 8.1 Hz, 1H), 4.80 (d, *J* = 11.2 Hz, 2H), 4.77 (d, *J* = 11.2 Hz, 1H), 4.75 (d, *J* = 11.2 Hz, 1H), 4.69 (d, *J* = 11.2 Hz, 1H), 4.69 (d, *J* = 11.2 Hz, 1H), 4.62 (d, *J* = 11.2 Hz, 1H), 4.59 (d, *J* = 11.2 Hz, 1H), 4.51 (d, *J* = 7.8 Hz, 1H), 4.35 (d, *J* = 8.1 Hz, 1H), 3.95–3.36 (m, 12H), 2.40–2.30 (m, 2H), 2.20–1.99 (m, 5H), 1.68–1.57 (quint., *J* = 7.2 Hz, 2H), 1.19 (s, 9H), 1.18 (s, 9H);

^{13}C -NMR (75 MHz, CDCl_3) δ 176.5, 137.9, 137.7, 137.6, 131.0, 128.4, 128.3, 128.0, 127.9, 127.8, 127.5, 127.3, 127.2, 126.8, 101.1, 100.9, 83.0, 77.9, 75.6, 75.3, 75.0, 74.9, 74.8, 72.9, 69.1, 68.3, 61.9, 38.8, 30.9 (t, $J = 22$ Hz), 29.5, 27.2, 23.7, 18.3; ^{19}F -NMR (282 MHz, CDCl_3) δ -80.6 (t, $J = 9.6$ Hz, 3F), -114.0- -114.3 (m, 2F), -121.1- -121.7 (m, 6F), -122.3 (bs, 2F), -123.0 (bs, 2F), -125.3 (m, 2F); IR (CHCl_3): 3600, 3436, 3008, 2970, 1737, 1602, 1524, 1479, 1454, 1397, 1361, 1200, 1146, 1090, 1042, 932, 874 cm^{-1} ; HRMS-MALDI Calcd for $\text{C}_{65}\text{H}_{73}\text{F}_{17}\text{O}_{13}\text{Na}$ 1407.4672 ($\text{M}+\text{Na}$) $^{+}$; Found: 1407.4703.

8,8,9,9,10,10,11,11,12,12,13,13,14,14,15,15,15-Heptafluoropentadec-4(Z)-en-1-yl 6-O-{6-O-[3,4-di-O-benzyl-2-O-pivaloyl- β -D-glucopyranosyl]-3,4-di-O-benzyl-2-O-pivaloyl- β -D-glucopyranosyl}-3,4-di-O-benzyl-2-O-pivaloyl- β -D-glucopyranoside (10).

General procedure using building blocks **9** (221 mg, 0.16 mmol, 0.08M in CH_2Cl_2), **1** (411 mg, 0.479 mmol, 0.24M in CH_2Cl_2), TMSOTf (86 μL , 0.479 mmol, 0.6M in CH_2Cl_2) and TBAF (1M in THF, 0.319 mmol, 0.08M in piperidine/DMF = 1:4) at +20 $^{\circ}\text{C}$ and 60 s reaction time gave a total pre-quenched volume of 3.06 mL (0.102 mmol) which afforded **10** (166 mg, 90%) as a white foam. $R_f = 0.74$ (EtOAc/cyclohexane = 1:2); $[\alpha]_{\text{D}}^{25} = -7.8^{\circ}$ ($c = 0.5$, CHCl_3); ^1H -NMR (300 MHz, CDCl_3) δ 7.37–7.22 (m, 30H), 5.47 (ddd, $J = 10.7, 7.2, 7.2$ Hz, 1H), 5.35 (ddd, $J = 10.7, 7.2, 7.2$ Hz, 1H), 5.10–4.99 (m, 3H), 4.81 (d, $J = 11.0$ Hz, 1H), 4.79 (d, $J = 11.0$ Hz, 2H), 4.76 (d, $J = 11.0$ Hz, 1H), 4.75 (d, $J = 11.0$ Hz, 2H), 4.71 (d, $J = 11.0$ Hz, 1H), 4.71 (d, $J = 11.0$ Hz, 1H), 4.68 (d, $J = 11.0$ Hz, 1H), 4.63 (d, $J = 11.0$ Hz, 1H), 4.61 (d, $J = 11.0$ Hz, 1H), 4.60 (d, $J = 11.0$ Hz, 1H), 4.50 (d, $J = 7.7$ Hz, 1H), 4.44 (d, $J = 7.7$ Hz, 1H), 4.39 (d, $J = 8.0$ Hz, 1H), 3.98–3.33 (m, 17H), 2.48 (dd, $J = 6.3, 6.3$ Hz, 1H), 2.39–2.29 (m, 2H), 2.20–1.99 (m, 4H), 1.68–1.57 (quint., $J = 7.0$ Hz, 2H), 1.19 (s, 9H), 1.19 (s, 9H), 1.17 (s, 9H); ^{13}C -NMR (75 MHz, CDCl_3) δ 176.7, 176.5, 176.4, 138.0, 137.9, 137.8, 137.7, 131.1, 128.3, 128.2, 128.1, 127.9, 127.8, 127.5, 127.3, 127.2, 126.7, 101.9, 101.1, 100.7, 83.1, 82.9, 77.9, 77.6, 75.7, 75.0, 74.7, 73.0, 72.8, 69.3, 68.8, 68.3, 61.7, 38.7, 30.7 (t, $J = 22$ Hz), 29.4, 27.0, 23.6, 18.2; ^{19}F -NMR (282 MHz, CDCl_3) δ -80.6, -80.7 (2t, $J = 9.6$ Hz, 3F), -114.3- -114.4, -115.9- -116.1 (2m, 2F), -121.3- -122.0 (m, 6F), -122.4- -123.0 (m, 2F), -123.4 (bs, 2F), -125.8- -126.2 (m, 2F); IR (CHCl_3): 3450, 3008, 2967, 2873, 1737, 1604, 1497, 1479, 454, 1398, 1363, 1200, 1144, 1088 cm^{-1} ; HRMS-MALDI Calcd for $\text{C}_{90}\text{H}_{103}\text{F}_{17}\text{O}_{19}\text{Na}$ 1833.6714 ($\text{M}+\text{Na}$) $^{+}$, Found: 1833.6730.

8,8,9,9,10,10,11,11,12,12,13,13,14,14,15,15,15-Heptadecafluoropentadec-4(Z)-en-1-yl 6-O-{6-O-[6-O-(3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranosyl)-3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranosyl]-3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranosyl}-3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranoside (11). General procedure using building blocks **10** (166 mg, 0.09 mmol, 0.08M in CH₂Cl₂), **1** (236 mg, 0.275 mmol, 0.24M in CH₂Cl₂), TMSOTf (50 μL, 0.275 mmol, 0.6M in CH₂Cl₂) and TBAF (1M in THF, 0.183 mmol, 0.08M in piperidine/DMF = 1:4) at +20 °C and 60 s reaction time gave a total pre-quenched volume of 2.19 mL (0.073 mmol) which afforded **11** (156 mg, 95%) as a white foam. R_f 0.75 (EtOAc/cyclohexane = 1:2); [α]_D²⁵ = -8.0° (c = 0.5, CHCl₃); ¹H-NMR (300 MHz, CDCl₃) δ 7.36–7.20 (m, 40H), 5.47 (ddd, J = 10.7, 7.1, 7.1 Hz, 1H), 5.36 (ddd, J = 10.7, 7.1, 7.1 Hz, 1H), 5.13–5.00 (m, 4H), 4.84–4.58 (m, 16H), 4.52 (d, J = 11.0 Hz, 1H), 4.44 (d, J = 7.7 Hz, 1H), 4.43 (d, J = 7.7 Hz, 1H), 4.39 (d, J = 8.2 Hz, 1H), 4.01 (d, J = 10.0 Hz, 1H), 3.92–3.33 (m, 21H), 2.66–2.55 (bs, 1H), 2.40–2.30 (m, 2H), 2.21–2.00 (m, 4H), 1.63 (quint., J = 6.9 Hz, 2H), 1.22 (s, 9H), 1.20 (s, 9H), 1.20 (s, 9H), 1.19 (s, 9H); ¹³C-NMR (75 MHz, CDCl₃) δ 176.8, 176.6, 176.5, 176.4, 138.0, 137.9, 137.9, 137.7, 131.1, 128.4, 128.3, 128.2, 127.9, 127.9, 127.8, 127.7, 127.4, 127.2, 127.2, 126.7, 102.0, 101.6, 100.7, 83.1, 82.9, 78.3, 77.7, 77.2, 75.7, 75.0, 74.9, 74.7, 74.3, 73.1, 72.9, 72.5, 69.4, 69.0, 68.7, 67.8, 61.7, 38.7, 30.7 (t, J = 22 Hz), 29.4, 27.1, 27.0, 27.0, 23.6, 18.1; ¹⁹F-NMR (282 MHz, CDCl₃) δ -80.6 (t, J = 9.6 Hz, 3F), -114.2- -114.5 (quint., 2F), -121.4- -122.0 (m, 6F), -122.5 (s, 2F), -123.3 (s, 2F), -125.8- -126.1 (m, 2F); IR (CHCl₃): 3066, 3008, 2967, 2935, 2874, 1737, 1602, 1497, 1479, 1454, 1397, 1363, 1277, 1200, 1144, 1085 cm⁻¹; HRMS-MALDI Calcd for C₁₁₅H₁₃₃F₁₇O₂₅Na 2259.8757 (M+Na)⁺; Found: 2259.8818.

8,8,9,9,10,10,11,11,12,12,13,13,14,14,15,15,15-Heptadecafluoropentadec-4(Z)-en-1-yl 6-O-{6-O-[6-O-(6-O-acetyl-3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranosyl)-3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranosyl]-3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranosyl}-3,4-di-O-benzyl-2-O-pivaloyl-β-D-glucopyranoside (12). To a solution of **11** (29 mg, 12.9 μmol) in pyridine (0.5 mL) was added acetic anhydride (0.25 mL) and DMAP (some crystals) at room temperature. After 4 h, the solution was diluted with CH₂Cl₂ (5 mL) and treated with ice-water (5 mL). The aqueous phase was extracted with CH₂Cl₂ (3 x 5 mL). The combined organic phases were washed with brine (3 mL) and dried over MgSO₄. The solvents were evaporated in *vacuo* and the residue was purified by silica gel flash column chromatography (EtOAc/hexane = 1:2) to afford **13** (27 mg, 92%) as a colorless

oil. R_f 0.72 (EtOAc/hexane = 1:2); $[\alpha]_D^{25} = -13.0^\circ$ ($c = 1.55$, CHCl_3); $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.38–7.16 (m, 40H), 5.47 (ddd, $J = 10.6, 7.0, 7.0$ Hz, 1H), 5.35 (ddd, $J = 10.6, 6.7, 6.7$ Hz, 1H), 5.15–4.98 (m, 4H), 4.79–4.55 (m, 16H), 4.50–4.34 (m, 4H), 4.29 (dd, $J = 11.2, 1.6$ Hz, 1H), 4.19 (dd, $J = 11.2, 5.2$ Hz, 1H), 4.06–3.34 (m, 20H), 2.41–2.28 (m, 2H), 2.21–2.00 (m, 4H), 1.99 (s, 3H), 1.71–1.55 (m, 2H), 1.19 (s, 36H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ 176.7, 176.4, 170.5, 138.1, 137.8, 131.5, 131.2, 128.4, 128.3, 128.2, 128.0, 127.9, 127.8, 127.6, 127.4, 127.2, 127.1, 127.1, 126.7, 101.4, 101.2, 100.8, 83.4, 83.1, 78.5, 78.3, 74.8, 72.9, 72.7, 68.7, 63.2, 38.7, 32.0, 29.6, 27.1, 27.0, 23.6, 20.8, 18.0; $^{19}\text{F-NMR}$ (282 MHz, CDCl_3) δ -80.5 (t, $J = 9.6$ Hz, 3F), -114.4 (quint., 2F), -121.7 (m, 6F), -122.5 (s, 2F), -123.3 (s, 2F), -125.9 (m, 2F); IR (CHCl_3): 3008, 2929, 1737, 1497, 1479, 1454, 1364, 1278, 1144, 1086 cm^{-1} ; HRMS-MALDI Calcd for $\text{C}_{117}\text{H}_{135}\text{F}_{17}\text{O}_{26}\text{Na}$ 2301.8868 ($\text{M}+\text{Na}$) $^+$; Found 2301.8814.

Pent-4(Z)-en-1-yl 6-O-{6-O-[6-O-(3,4-di-O-benzyl-2-O-pivaloyl- β -D-glucopyranosyl)-3,4-di-O-benzyl-2-O-pivaloyl- β -D-glucopyranosyl]-3,4-di-O-benzyl-2-O-pivaloyl- β -D-glucopyranosyl}-3,4-di-O-benzyl-2-O-pivaloyl- β -D-glucopyranoside (13). To a solution of **11** (4.7 mg, 2.1 μmol) in CH_2Cl_2 (2 mL) was added Grubbs' 2nd generation catalyst (0.2 mg, 5 mol%, 0.3 μmol) at room temperature. The solution was degassed with vacuum/argon (3 times) and ethylene (excess, 1 atm.) was bubbled into the solution for one minute. The reaction mixture was stirred at room temperature under ethylene atmosphere for 12 h, the TFT (0.5 mL) and Grubbs' catalyst (0.2 mg) were added and ethylene was bubbled into the solution for another minute. After 12 h under ethylene atmosphere, Grubbs' catalyst (0.2 mg) were added and ethylene was bubbled into the solution for 1 min. After another 12 h under ethylene atmosphere, the black solution was evaporated to dryness in *vacuo*. The residue was purified using HPLC to afford **13** (r_t 65 min, 2.1 mg, 56%) and **11** (r_t 69 min, 2.1 mg, 44%) as white solids. R_f 0.41 (EtOAc/cyclohexane = 1:3); $[\alpha]_D^{25} = -14.4^\circ$ ($c = 0.5$, CHCl_3); $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.34–7.18 (m, 40H), 5.78 (dddd, $J = 17.1, 10.2, 6.6, 6.6$ Hz, 1H), 5.10–4.92 (m, 6H), 4.80–4.56 (m, 16H), 4.50 (d, $J = 11.0$ Hz, 1H), 4.42 (d, $J = 7.7$ Hz, 2H), 4.35 (d, $J = 8.0$ Hz, 1H), 4.02–3.30 (m, 22H), 2.53 (t, $J = 6.3, 6.3$ Hz, 1H), 2.12–2.02 (m, 2H), 1.69–1.58 (m, 2H), 1.19 (s, 9H), 1.19 (s, 9H), 1.18 (s, 9H), 1.16 (s, 9H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3) δ 176.7, 176.6, 176.4, 138.1, 138.0, 138.0, 137.9, 137.8, 137.7, 128.4, 128.4, 128.3, 128.2, 128.0, 127.9, 127.9, 127.8, 127.5, 127.3, 127.2, 114.9, 102.0, 101.7, 100.8, 83.3, 83.2, 83.0, 78.5, 77.8, 77.3, 75.8, 75.3, 75.1, 74.9, 74.8, 74.7, 74.5, 73.2,

73.1, 72.7, 69.4, 69.1, 68.9, 67.9, 61.9, 38.9, 30.2, 28.9, 27.1, 27.1; IR (CHCl₃): 3064, 3038, 2966, 2875, 1737, 1605, 1497, 1479, 1454, 1397, 1363, 1278, 1261, 1200, 1139, 1086, 1038, 914 cm⁻¹; HRMS-MALDI Calcd for C₁₀₅H₁₃₀O₂₅Na 1813.8793 (M+Na)⁺; Found: 1813.9023.

***O*-(6-*O*-{6-*O*-[6-*O*-(6-*O*-Acetyl-3,4-di-*O*-benzyl-2-*O*-pivaloyl-β-D-glucopyranosyl)-3,4-di-*O*-benzyl-2-*O*-pivaloyl-β-D-glucopyranosyl]-3,4-di-*O*-benzyl-2-*O*-pivaloyl-α-D-glucopyranosyl)trichloroacetimidate (14).** To a solution of **12** (14.2 mg, 6.23 μmol) in CH₃CN (+1% H₂O, 630 μL) was added NBS (3.3 mg, 18.7 mmol) at room temperature. After 30 min, the solution was treated with saturated aqueous Na₂S₂O₃ solution (0.6 mL) and water (0.6 mL) and extracted with Et₂O (3 x 5 mL). The combined organic phase was dried over MgSO₄ and the solvents were evaporated in *vacuo* to afford the crude lactol [(23 mg, α/β ~1:1, R_f (α) 0.27, R_f (β) 0.15 (EtOAc/cyclohexane = 1:3)] as yellowish oil. The residue was dissolved in CH₂Cl₂ (600 μL) and trichloroacetonitrile (20 μL, 32 eq, 0.2 mmol) was added at room temperature. The solution was cooled to 0 °C and DBU (1 μL, 6.7 μmol) was added. After 10 min, the solution was allowed to warm to room temperature and was stirred for further 30 min. The solvents were evaporated in *vacuo* and the residue was purified by silica gel flash column chromatography (EtOAc/cyclohexane = 1:4, +1%NEt₃) to afford **14** (11.7 mg, 98%, α/β = 6:1) as colorless oil. R_f (α) 0.35 (EtOAc/cyclohexane = 1:4); [α]_D²⁵ = 4.7° (c = 0.5, CHCl₃); ¹H-NMR (300 MHz, CDCl₃, α-isomer) δ 8.67 (s, 1H), 7.35–7.19 (m, 40H), 6.50 (d, J = 3.6 Hz, 1H), 5.11 (dd, J = 9.1, 7.7 Hz, 1H), 5.09 (dd, J = 9.3, 7.1 Hz, 1H), 5.07 (dd, J = 10.2, 3.6 Hz, 1H), 5.01 (dd, J = 8.5, 8.3 Hz, 1H), 4.82–4.46 (m, 19H), 4.30 (dd, J = 11.2, 2.2 Hz, 1H), 4.16 (dd, J = 12.1, 5.1 Hz, 1H), 4.13–3.42 (m, 18H), 1.93 (s, 3H), 1.17 (s, 9H), 1.15 (s, 9H), 1.14 (s, 9H), 1.10 (s, 9H); IR (CHCl₃): 3038, 2965, 2872, 1737, 1674, 1602, 1497, 1479, 1455, 1397, 1363, 1262, 1200, 1138, 1090, 910 cm⁻¹; HRMS-MALDI Calcd for C₁₀₄H₁₂₄Cl₃NO₂₆Na 1930.7369 (M+Na)⁺; Found: 1930.7317.

4-Oxobutan-1-yl 6-*O*-{6-*O*-[6-*O*-(6-*O*-acetyl-3,4-di-*O*-benzyl-2-*O*-pivaloyl-β-D-glucopyranosyl)-3,4-di-*O*-benzyl-2-*O*-pivaloyl-β-D-glucopyranosyl]-3,4-di-*O*-benzyl-2-*O*-pivaloyl-β-D-glucopyranosyl}-3,4-di-*O*-benzyl-2-*O*-pivaloyl-β-D-glucopyranoside (15). To a solution of **12** (12.0 mg, 5.3 μmol) in CH₂Cl₂ (1.5 mL) was bubbled O₃ (3% in O₂) at -78 °C for 3 min. The light-blue solution was treated with PPh₃ (2 mg, 7.6 μmol) and was allowed to warm to room temperature. After 3 h, the solution was treated with saturated aqueous

NH₄Cl solution (5 mL) and extracted with CH₂Cl₂ (3 x 5 mL). The combined organic phase was washed with brine (3 mL) and dried over MgSO₄. The solvents were evaporated in *vacuo* and the residue was purified by silica gel flash column chromatography (EtOAc/hexane = 1:3) to afford **15** (6.5 mg, 68%) as colorless oil. R_f 0.32 (EtOAc/hexane = 1:3); [α]_D²⁵ = -13.3° (c = 0.61, CHCl₃); ¹H-NMR (300 MHz, CDCl₃) δ 9.76 (t, *J* = 1.2 Hz, 1H), 7.32–7.14 (m, 40H), 5.10 (dd, *J* = 7.8, 6.8 Hz, 1H), 5.07 (dd, *J* = 7.7, 6.2 Hz, 1H), 5.03 (dd, *J* = 7.8, 7.8 Hz, 1H), 5.00 (dd, *J* = 8.1, 8.1 Hz, 1H), 4.76–4.53 (m, 16H), 4.44–4.22 (m, 5H), 4.18 (dd, *J* = 11.8, 5.0 Hz, 1H), 3.98–3.34 (m, 20H), 2.57–2.48 (m, 2H), 2.00 (s, 3H), 1.91–1.80 (m, 2H), 1.20 (s, 9H), 1.18 (s, 9H), 1.18 (s, 9H), 1.17 (s, 9H). ¹³C-NMR (75 MHz, CDCl₃) δ 201.9; 176.6, 176.2, 170.4, 138.1, 138.0, 137.8, 137.7, 137.5, 128.3, 128.2, 128.2, 128.2, 127.8, 127.5, 127.3, 127.2, 127.0, 101.6, 101.0, 100.5, 83.1, 77.2, 74.9, 74.6, 74.2, 73.2, 73.0, 69.1, 68.5, 63.4, 40.6, 38.9, 29.7, 27.3, 27.2, 22.2, 21.0; IR (CHCl₃): 3008, 2960, 1737, 1455, 1364, 1278, 1139, 1067, 909 cm⁻¹; HRMS-MALDI Calcd for C₁₀₆H₁₃₀O₂₇Na 1857.8697 (M+Na)⁺; Found: 1857.8649.

References

- (1) Carrel, F.; Seeberger, P.H. *J. Carbohydr. Chem.*, **2007**, *in press*.
- (2) Markidis, T.; Kokotos, G. *J. Org. Chem.*, **2002**, *67*, 1685-1688.
- (3) Rocaboy, C.; Bauer, W.; Gladysz, J. A. *Eur. J. Org. Chem.*, **2000**, *14*, 2621-2628.

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 01/18/07 13:06:44 USER:pfcar GROUP:seeber

^1H -NMR of 2

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 06/15/06 10:24:41 USER:pfarr GROUP:seeber

^{13}C -NMR of 2

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 06/15/06 10:06:24 USER:pfcar GROUP:seeber

¹⁹F-NMR of 2

LOC ETHZ NMR Mercury-vx 300MHz Nr.6 11/09/06 13:07:14 USER:plcarr GROUP:seeber

LOC ETHZ NMR Mercury-vx 300MHz Nr.6 11/09/06 13:38:32 USER:ptcarr GROUP:seeber

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 11/10/06 12:49:52 USER:pfarr GROUP:seeber

LOC ETHZ NMR Mercury-vx 300MHz Nr.6 01/18/07 14:25:21 USER:ptfcar GROUP:seeber

¹H-NMR of 7

LOC ETHZ NMR Mercury-Vx 300MHz Nr.5 06/01/06 17:55:43 USER:ptcarr GROUP:seeber

¹³C-NMR of 7

LOC ETHZ NMR Mercury-ix 300MHz Nr.5 06/01/06 17:52:50 USER:pfcaTr GROUP:seeber

¹⁹F-NMR of 7

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 01/18/07 14:07:40 USER:pfarrt GROUP:seeber

¹H-NMR of 8

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 08/02/06 14:13:11 USER:pfcarr GROUP:seeber

¹⁹F-NMR of 8

LOC ETHZ NMR Mercury-vx 300MHz Nr.6 01/18/07 14:13:58 USER:ptcarr GROUP:seeber

¹H-NMR of 9

¹³C-NMR of 9

¹⁹F-NMR of 9

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 01/18/07 14:20:55 USER:pfcarr GROUP:seeber

¹H-NMR of 10

¹⁹F-NMR of 10

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 10/24/06 13:13:39 USER:pfearr GROUP:seeber

¹H-NMR of 11

¹³C-NMR of 11

¹⁹F-NMR of 11

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 01/18/07 16:52:52 USER:pgeyer GROUP:seeber

¹H-NMR of 12

LOC ETHZ NMR Mercury-vx 300MHz Nr.5 01/18/07 16:40:51 USER:ppg/yer GROUP:seeber

¹⁹F-NMR of 12

¹³C-NMR of 13

LOC ETHZ NMR Mercury-vx 300MHz Nr.6 01/18/07 17:01:51 USER:pgeyer GROUP:seeber

¹H-NMR of 15

LOC ETHZ NMR Mercury-vx 300MHz Nr.6 01/18/07 17:11:05 USER:pgeyer GROUP:seeber

¹³C-NMR of 15