Supporting information

for

**σ -antibonding between valence s-orbitals as a primary cause for band gaps in
compounds of diamond-like structures**

Jürgen Köhler* and Shuiquan Deng

Max-Planck-Institut für Festkörperforschung, Heisenbergstr. 1, D-70569 Stuttgart,
Germany.

Changhoon Lee and Myung-Hwan Whangbo*

Department of Chemistry, North Carolina State University, Raleigh, North Carolina
27695-8204, USA

Figure S.1. COHP plot calculated for the Au-Sn bond on the basis of first principles TB-LMTO calculations for ScAuSn.

Figure S.2. Dispersion relations of the bands around the Fermi level calculated for (a) CaAuBi and (b) YAuPb by using the self-consistent full-potential linearized augmented-plane-wave method embodied in the WIEN2k code. The fat band representation is given for the valence s-orbital contributions of all the atoms.

Figure S.3. Schematic representation of the σ -antibonding orbital of AML at Γ , which is made up of the valence ns orbitals of M and L, around each electropositive atom A (a yellow circle) located at the center of an M_4 tetrahedron (shaded large circles) and an L_6 octahedron (unshaded medium circles). The ns orbitals of the M atoms make antibonding interactions with those of the adjacent L atoms such that the four ns orbitals of the M_4 tetrahedron make a group orbital of a_1 symmetry, and those of the L_6 octahedron a group orbital of a_{1g} symmetry.