Supporting Information

One-Pot Amidation of Olefins through Pd-Catalyzed Coupling of Alkylboranes and Carbamoyl Chlorides

Yoshizumi Yasui, Sayo Tsuchida, Hideto Miyabe, and Yoshiji Takemoto*

Graduate School of Pharmaceutical Sciences,
Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan

Table of contents

General methods	S2
Preparation and analytical data of compound 3a	S2
Preparation and analytical data of compound 3d	S3
Preparation and analytical data of compound 4aa	S3
Preparation and analytical data of compound 4ba	S3
Preparation and analytical data of compound 4ca	S4
Preparation and analytical data of compound 4da	S4
Preparation and analytical data of compound 4ea	S5
Preparation and analytical data of compound 4fa	S5
Preparation and analytical data of compound 4ab	S5
Preparation and analytical data of compound 4ac	S6
Preparation and analytical data of compound 4ad	S6
Preparation and analytical data of compound 5	S6
Preparation and analytical data of compound 6	S7
Preparation and analytical data of compound 7	S7
Copies of ¹ HNMR and ¹³ CNMR spectrum	S8–S35

General methods. All reactions were performed under argon. Pd(PPh₃)₄ was prepared by using the reported protocol.¹ Cs₂CO₃ was intensively dried by heating under vacuum followed by cooling to room temperature under argon before use. *N,N*-dimethylchloroformamide (**3b**) and morpholine-4-carbonyl chloride (**3c**) were distilled before use. 3-methoxymethoxy-3-phenylprop-1-ene (**1e**) and 3-methylbut-3-enyl benzoate (**1f**) were synthesized from corresponding commercially available 1-phenylprop-2-en-1-ol and 3-methylbut-3-en-1-ol. Other commercially available reagents and solvents were used as received without further purification.

Analytical thin-layer chromatography was performed with Merck silica gel 60 F₂₅₄. Flash silica gel column chromatography was performed with Kanto silica gel 60 (particle size 0.063–0.210 mm). Proton nuclear magnetic resonance (¹H NMR) spectra were recorded at 500 MHz and chemical shifts are reported relative to Me₄Si (δ 0.00). Multiplicity is indicated by one or more of the following: s (singlet); d (doublet); t (triplet); m (multiplet); br (broad). Carbon nuclear magnetic resonance (¹³C NMR) spectra were recorded at 125 MHz and chemical shifts are reported relative to CDCl₃ (δ 77.2). Infrared spectra were recorded as KBr pellets (KBr) or thin film on NaCl plates (thin film). Only characteristic absorption bands are reported. Only charged molecular ion data are reported for both low-resolution mass spectra (LRMS) and high-resolution mass spectra (HRMS).

Experimental Procedures.

***N,N*-Dibenzylchloroformamide (**3a**)²:** To a solution of triphosgene (2.70 g, 9.14 mmol) in CH₂Cl₂ (80 mL) was added triethylamine (5.40 mL, 39.2 mmol) followed by dibenzylamine (5.00 mL, 26.1 mmol) at –78 °C. The reaction mixture was warmed to room temperature, and water was added at 0 °C, and the product was extracted with EtOAc. The combined organic extracts were washed with 1 M HCl, water, saturated aqueous NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered, and concentrated. Purification by silica gel column chromatography (hexane/EtOAc = 95/5) afforded **3a** (6.59 g, 97%) as a white solid: R_f 0.55 (hexane/Et₂O = 8/2); mp. 39.5–40.0 °C; ¹H NMR (δ, CDCl₃) 7.38–7.30 (m, 6H), 7.23 (d, 4H, *J* = 7.3 Hz), 4.61 (s, 2H), 4.50 (s, 2H); ¹³C NMR (δ, CDCl₃) 150.5, 135.4, 135.2, 129.1, 129.0, 128.5, 128.3, 128.2, 127.4, 53.0, 51.4; IR (KBr) 1725 cm^{–1}; HRMS (EI) Calcd for C₁₅H₁₄ClNO (M⁺) 259.0764, found 259.0773; Anal. Calcd for C₁₅H₁₄ClNO: C, 69.36; H, 5.43; N, 5.39. Found: C, 69.43; H, 5.43; N, 5.52.

Methyl 1-(chlorocarbonyl)pyrrolidine-2-carboxylate (3d**)³:** To a solution of triphosgene (945 mg, 3.18 mmol) in THF (15 mL) was added a suspension of methyl D,L-proline hydrochloride (1.51 g, 9.15 mmol) and triethylamine (2.00 mL, 26.8 mmol) in THF (15 mL) at –78 °C. The reaction mixture was warmed to room temperature, and water was added at 0 °C, and the

¹ Coulson, D. R. *Inorg. Synth.* **1972**, 13, 121.

² Jorand-Lebrun, C.; Valognes, D.; Halazy, S. *Synth. Commun.* **1998**, 28, 1189.

³ Drauz, K.; Kleemann, A.; Martens, J.; Scherberich, P. *J. Org. Chem.* **1986**, 51, 3494.

product was extracted with EtOAc. The combined organic extracts were washed with 1 M HCl, water, saturated aqueous NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered, and concentrated. Purification by silica gel column chromatography (hexane/EtOAc = 95/5) afforded **3d** (1.61 g, 93%) as a colorless oil, which was distilled before use: R_f 0.36 (hexane/EtOAc = 8/2); ¹H NMR (δ, CDCl₃) 4.58* (dd, *J*₁ = 3.0 Hz, *J*₂ = 8.8 Hz), 4.46 (dd, 1H, *J*₁ = 3.4 Hz, *J*₂ = 8.6 Hz), 3.79* (s), 3.76 (s, 3H), 3.74–3.56 (m, 2H), 2.38–2.29 (m, 1H), 2.16–1.99 (m, 3H); ¹³C NMR (δ, CDCl₃) 171.7*, 171.1, 147.4, 146.5*, 62.3*, 60.6, 52.7*, 52.5, 50.5, 49.1*, 30.1, 23.5*, 23.4; IR (thin film) 1744 cm⁻¹; LRMS (EI) *m/z* = 191 (M⁺); Anal. Calcd for C₇H₁₀ClNO₃: C, 43.88; H, 5.26; N, 7.31. Found: C, 43.60; H, 5.18; N, 7.03. (* Peaks of minor rotamer)

***N,N*-Dibenzyl-3-phenylpropanamide (4aa)**⁴: To styrene (**1a**) (220 μL, 1.92 mmol) at 0 °C was added 9-BBN-H (0.5 M solution in THF, 4.20 mL, 2.10 mmol). The reaction mixture was warmed to room temperature and stirred for 2 h. *N,N*-Dibenzylchloroformamide (**3a**) (343 mg, 1.32 mmol) in anhydrous THF (3 mL), Cs₂CO₃ (636 mg, 1.95 mmol) and Pd(PPh₃)₄ (77.0 mg, 0.0666 mmol) were added sequentially at 0 °C. After stirring for 16 h at reflux, water was added at 0 °C, and the product was extracted with EtOAc. The combined organic extracts were washed with brine, dried over MgSO₄, filtered, and concentrated. Purification by silica gel column chromatography (hexane/EtOAc = 9/1→5/5), followed by trituration from hexane, and drying under vacuum gave **4aa** (339 mg, 78%) as a white solid: R_f 0.48 (hexane/Et₂O = 5/5); mp. 104.5–105.0 °C; ¹H NMR (δ, CDCl₃) 7.35–7.24 (m, 8H), 7.20–7.17 (m, 5H), 7.07 (d, 2H, *J* = 7.4 Hz), 4.60 (s, 2H), 4.37 (s, 2H), 3.05 (t, 2H, *J* = 3.1 Hz), 2.72 (t, 2H, *J* = 3.1 Hz); ¹³C NMR (δ, CDCl₃) 173.0, 141.4, 137.5, 136.6, 129.1, 128.8, 128.7, 128.6, 128.5, 127.8, 127.6, 126.5, 126.3, 50.0, 48.4, 35.1, 31.7; IR (KBr) 1633 cm⁻¹; HRMS (EI) Calcd for C₂₃H₂₃NO (M⁺) 329.1779, found 329.1773; Anal. Calcd for C₂₃H₂₃NO: C, 83.85; H, 7.04; N, 4.25. Found: C, 84.15; H, 7.20; N, 4.24.

***N,N*-Dibenzylheptanamide (4ba)**: Amide **4ba** (299 mg, 72%) was obtained as a colorless oil as described for the synthesis of compound **4aa** but using 1-hexene (250 μL, 2.02 mmol), 9-BBN-H (0.5 M solution in THF, 4.40 mL, 2.20 mmol), *N,N*-dibenzylchloroformamide (**3a**) (356 mg, 1.37 mmol), Cs₂CO₃ (664 mg, 2.04 mmol) and Pd(PPh₃)₄ (79.0 mg, 0.0684 mmol). Purification was carried out by silica gel column chromatography (hexane/Et₂O = 10/0→8/2), followed by drying under vacuum (10 mmHg, 180 °C) to remove the trace 9-BBN residue: R_f 0.43 (hexane/Et₂O = 7/3); ¹H NMR (δ, CDCl₃) 7.37 (dd, 2H, *J*₁ = *J*₂ = 7.0 Hz), 7.33–7.25 (m, 4H), 7.23 (d, 2H, *J* = 7.0 Hz), 7.15 (d, 2H, *J* = 7.0 Hz), 4.60 (s, 2H), 4.45 (s, 2H), 2.42 (t, 2H, *J* = 7.7 Hz), 1.74–1.68 (m, 2H), 1.36–1.26 (m, 6H), 0.87 (t, 3H, *J* = 6.7 Hz); ¹³C NMR (δ, CDCl₃) 173.9, 137.7, 136.8, 129.0, 128.7, 128.4, 127.7, 127.4, 126.5, 49.9, 48.1, 33.3, 31.7, 29.1, 25.5, 22.5, 14.1; IR (thin film) 1650 cm⁻¹; LRMS (EI) *m/z* = 309 (M⁺); Anal. Calcd for C₂₁H₂₇NO: C, 81.51; H, 8.79; N, 4.53. Found: C, 81.30; H, 8.76; N, 4.39.

⁴ Aoyagi, Y.; Asakura, R.; Kondoh, N.; Yamamoto, R.; Kuromatsu, T.; Shimura, A.; Ohta, A. *Synthesis* **1996**, 970.

***N,N*-Dibenzyl-4-trimethylsilylbutanamide (4ca):** Amide **4ca** (320 mg, 74%) was obtained as a white solid as described for the synthesis of compound **4aa** but using allyltrimethylsilane (300 μ L, 1.89 mmol), 9-BBN-H (0.5 M solution in THF, 4.00 mL, 2.00 mmol), *N,N*-dibenzylchloroformamide (**3a**) (328 mg, 1.27 mmol), Cs₂CO₃ (589 mg, 1.81 mmol) and Pd(PPh₃)₄ (72.8 mg, 0.0630 mmol). Purification was carried out by silica gel column chromatography (hexane/Et₂O = 10/0→8/2), followed by drying under vacuum (10 mmHg, 180 °C) to remove the trace 9-BBN residue. The initially obtained colorless oil changed into a white solid by standing at room temperature for a few days: R_f 0.39 (hexane/Et₂O = 7/3); mp. 43.0–43.5 °C; ¹H NMR (δ , CDCl₃) 7.38–7.15 (m, 10H), 4.61 (s, 2H), 4.45 (s, 2H), 2.45 (t, 2H, *J* = 7.6 Hz), 1.75–1.69 (m, 2H), 0.54–0.51 (m, 2H), –0.02 (s, 9H); ¹³C NMR (δ , CDCl₃) 173.9, 137.8, 136.9, 129.1, 128.8, 128.5, 127.8, 127.5, 126.6, 50.0, 48.2, 37.1, 20.2, 16.8, –1.67; IR (KBr) 1648 cm^{–1}; LRMS (EI) *m/z* = 399 (M⁺); Anal. Calcd for C₂₁H₂₉NOSi: C, 74.28; H, 8.61; N, 4.13. Found: C, 74.40; H, 8.64; N, 4.17.

***N,N*-Dibenzyl-4-benzyloxybutanamide (4da):** Amide **4da** (327 mg, 80%) was obtained as a colorless oil as described for the synthesis of compound **4aa** but using allylbenzylether (250 μ L, 1.62 mmol), 9-BBN-H (0.5 M solution in THF, 3.60 mL, 1.80 mmol), *N,N*-dibenzylchloroformamide (**3a**) (285 mg, 1.10 mmol), Cs₂CO₃ (528 mg, 1.62 mmol) and Pd(PPh₃)₄ (62.4 mg, 0.0540 mmol). Purification was carried out by silica gel column chromatography (hexane/Et₂O = 95/5), followed by drying under vacuum (10 mmHg, 180 °C) to remove the trace 9-BBN residue: R_f 0.38 (hexane/Et₂O = 5/5); ¹H NMR (δ , CDCl₃) 7.36–7.24 (m, 11H), 7.21 (d, 2H, *J* = 6.7 Hz), 7.14 (d, 2H, *J* = 7.3 Hz), 4.61 (s, 2H), 4.46 (s, 2H), 4.44 (s, 2H), 3.55 (t, 2H, *J* = 6.1 Hz), 2.55 (t, 2H, *J* = 7.0 Hz), 2.05 (tt, 2H, *J*₁ = 6.1 Hz, *J*₂ = 7.0 Hz); ¹³C NMR (δ , CDCl₃) 173.5, 138.7, 137.7, 136.8, 129.1, 128.8, 128.5, 128.4, 127.8, 127.8, 127.7, 127.6, 126.6, 73.0, 69.6, 50.0, 48.3, 29.8, 25.6; IR (thin film) 1647 cm^{–1}; LRMS (FAB) *m/z* = 374 (M+H⁺); Anal. Calcd for C₂₅H₂₇NO₂: C, 80.40; H, 7.29; N, 3.75. Found: C, 80.59; H, 7.32; N, 3.60.

***N,N*-Dibenzyl-4-methoxymethoxy-4-phenylbutanamide (4ea):** Amide **4ea** (370 mg, 86%) was obtained as a colorless oil as described for the synthesis of compound **4aa** but using 3-methoxymethoxy-3-phenylprop-1-ene (**1e**) (286 mg, 1.62 mmol), 9-BBN-H (0.5 M solution in THF, 3.40 mL, 1.70 mmol), *N,N*-dibenzylchloroformamide (**3a**) (278 mg, 1.07 mmol), Cs₂CO₃ (528 mg, 1.62 mmol) and Pd(PPh₃)₄ (62.0 mg, 0.0536 mmol). Purification was carried out by silica gel column chromatography (hexane/Et₂O = 95/5→8/2), followed by drying under vacuum (10 mmHg, 180 °C) to remove the trace 9-BBN residue: R_f 0.37 (hexane/Et₂O = 7/3); ¹H NMR (δ, CDCl₃) 7.34–7.20 (m, 13H), 7.09 (d, 2H, *J* = 7.6 Hz), 4.68 (t, 1H, *J* = 5.8 Hz), 4.61 (s, 2H), 4.50 (s, 2H), 4.37 (s, 2H), 3.25 (s, 3H), 2.50 (t, 2H, *J* = 8.3 Hz), 2.25–2.11 (m, 2H); ¹³C NMR (δ, CDCl₃) 173.1, 141.7, 137.6, 136.7, 129.0, 128.7, 128.5, 128.4, 127.8, 127.7, 127.5, 126.9, 126.6, 94.4, 77.4, 55.6, 50.0, 48.3, 33.3, 29.4; IR (thin film) 1647 cm⁻¹; LRMS (ESI) *m/z* = 404 (M+H⁺); Anal. Calcd for C₂₆H₂₉NO₃: C, 77.39; H, 7.24; N, 3.47. Found: C, 77.43; H, 7.36; N, 3.35.

5-Dibenzylamino-3-methyl-5-oxopentyl benzoate (4fa): Amide **4fa** (384 mg, 77%) was obtained as a colorless oil as described for the synthesis of compound **4aa** but using 3-methylbut-3-enyl benzoate (**1f**) (367 mg, 1.80 mmol), 9-BBN-H (0.5 M solution in THF, 3.80 mL, 1.90 mmol), *N,N*-dibenzylchloroformamide (**3a**) (310 mg, 1.20 mmol), Cs₂CO₃ (587 mg, 1.80 mmol) and Pd(PPh₃)₄ (69.3 mg, 0.0600 mmol). Purification was carried out by silica gel column chromatography (hexane/Et₂O = 10/0→8/2), followed by drying under vacuum (10 mmHg, 180 °C) to remove the trace 9-BBN residue: R_f 0.38 (hexane/Et₂O = 5/5); ¹H NMR (δ, CDCl₃) 8.02 (d, 2H, *J* = 7.9 Hz), 7.53 (t, 1H, *J* = 7.4 Hz), 7.41–7.21 (m, 10H), 7.13 (d, 2H, *J* = 7.4 Hz), 4.63 (d, 1H, *J* = 15.0 Hz), 4.60 (d, 1H, *J* = 15.0 Hz), 4.45 (s, 2H), 4.41–4.33 (m, 2H), 2.48 (dd, 1H, *J*₁ = 5.5 Hz, *J*₂ = 14.7 Hz), 2.44–2.37 (m, 1H), 2.34 (dd, 1H, *J*₁ = 7.7 Hz, *J*₂ = 14.7 Hz), 1.95–1.88 (m, 1H), 1.68–1.61 (m, 1H), 1.07 (d, 3H, *J* = 6.4 Hz); ¹³C NMR (δ, CDCl₃) 172.6, 166.8, 137.7, 136.7, 133.0, 130.5, 129.7, 129.1, 128.8, 128.5, 128.4, 127.8, 127.5, 126.5, 63.2, 50.0, 48.2, 40.4, 35.5, 27.8, 19.9; IR (thin film) 1714, 1643 cm⁻¹; LRMS (FAB) *m/z* = 416 (M+H⁺); Anal. Calcd for C₂₇H₂₉NO₃: C, 78.04; H, 7.03; N, 3.37. Found: C, 77.91; H, 7.10; N, 3.33.

***N,N*-Dimethyl-3-phenylpropanamide (4ab)**⁵: Amide **4ab** (399 mg, 87%) was obtained as a colorless oil as described for the synthesis of compound **4aa** but using styrene (440 μL, 3.84 mmol), 9-BBN-H (0.5 M solution in THF, 8.40 mL, 4.20 mmol), *N,N*-dimethylchloroformamide (**3b**) (240 μL, 2.60 mmol), Cs₂CO₃ (1.26 g, 3.87 mmol) and Pd(PPh₃)₄ (149 mg, 0.129 mmol). Purification was carried out by silica gel column chromatography (hexane/Et₂O = 10/0→0/10): R_f 0.60 (Et₂O); ¹H NMR (δ, CDCl₃) 7.28 (dd, 2H, *J*₁ = *J*₂ = 7.6 Hz), 7.22 (d, 2H, *J* = 7.6 Hz), 7.19 (t, 1H, *J* = 7.6 Hz), 2.96 (t, 2H, *J* = 7.9 Hz), 2.94 (s, 3H), 2.91 (s, 3H), 2.61 (t, 2H, *J* = 7.9 Hz); ¹³C NMR (δ, CDCl₃) 172.3, 141.6, 128.6, 128.5,

⁵ Hall, L. R.; Iwamoto, R. T.; Hanzlik, R. P. *J. Org. Chem.* **1989**, *54*, 2446.

126.2, 37.1, 35.3, 35.3, 31.4; IR (thin film) 1645 cm^{-1} ; LRMS (EI) $m/z = 177$ (M^+); Anal. Calcd for $\text{C}_{11}\text{H}_{15}\text{NO}$: C, 74.54; H, 8.53; 7.90. Found: C, 74.68; H, 8.67; N, 7.95.

1-Morpholino-3-phenylpropan-1-one (4ac)⁶: Amide **4ac** (239 mg, 62%) was obtained as a colorless oil as described for the synthesis of compound **4aa** but using styrene (300 μL , 2.62 mmol), 9-BBN-H (0.5 M solution in THF, 5.60 mL, 2.80 mmol), morpholine-4-carbonyl chloride (**3c**) (262 mg, 1.74 mmol), Cs_2CO_3 (862 mg, 2.64 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (104 mg, 0.0900 mmol). Purification was carried out by silica gel column chromatography (hexane/ $\text{Et}_2\text{O} = 10/0 \rightarrow 0/10$): Rf 0.67 (Et_2O); ^1H NMR (δ , CDCl_3) 7.29 (t, 2H, $J = 8.5$ Hz), 7.22–7.20 (m, 3H), 3.62 (br s, 4H), 3.51 (t, 2H, $J = 4.6$ Hz), 3.35 (t, 2H, $J = 4.6$ Hz), 2.98 (t, 2H, $J = 7.7$ Hz), 2.61 (t, 2H, $J = 7.7$ Hz); ^{13}C NMR (δ , CDCl_3) 171.1, 141.3, 128.8, 128.7, 126.5, 67.0, 66.6, 46.1, 42.1, 34.9, 31.6; IR (thin film) 1645 cm^{-1} ; LRMS (EI) $m/z = 219$ (M^+); Anal. Calcd for $\text{C}_{13}\text{H}_{17}\text{NO}_2$: C, 71.21; H, 7.81; N, 6.39. Found: C, 71.00; H, 7.95; N, 6.20.

Methyl 1-(3-phenylpropanoyl)pyrrolidine-2-carboxylate (4ad)⁷: Amide **4ad** (345 mg, 75%) was obtained as a colorless oil as described for the synthesis of compound **4aa** but using styrene (300 μL , 2.62 mmol), 9-BBN-H (0.5 M solution in THF, 5.60 mL, 2.80 mmol), methyl 1-(chlorocarbonyl)pyrrolidine-2-carboxylate (**3d**) (335 mg, 1.76 mmol), Cs_2CO_3 (860 mg, 2.64 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (102 mg, 0.0883 mmol). Purification was carried out by silica gel column chromatography (hexane/ $\text{Et}_2\text{O} = 10/0 \rightarrow 0/10$): Rf 0.50 (Et_2O); ^1H NMR (δ , CDCl_3) 7.28–7.16 (m, 5H), 4.46 (dd, 1H, $J_1 = 4.9$ Hz, $J_2 = 8.6$ Hz), 4.25* (dd, $J_1 = 3.1$ Hz, $J_2 = 8.2$ Hz), 3.71 (s, 3H), 3.68* (s), 3.65–3.60* (m), 3.56–3.51 (m, 1H), 3.39–3.34 (m, 1H), 3.01–2.91 (m, 2H), 2.66–2.54 (m, 2H), 2.50–2.38* (m), 2.16–1.81 (m, 4H); ^{13}C NMR (δ , CDCl_3) 172.8, 172.5*, 171.0*, 170.9, 141.2, 128.3 (2C), 125.9, 59.1*, 58.4, 52.3*, 51.9, 46.7, 46.1*, 36.1, 36.0*, 31.1*, 31.0*, 30.5, 29.0, 24.5, 22.3*; IR (thin film) 1745, 1647 cm^{-1} ; LRMS (EI) $m/z = 261$ (M^+); Anal. Calcd for $\text{C}_{15}\text{H}_{19}\text{NO}_3$: C, 68.94; H, 7.33; N, 5.36. Found: C, 68.65; H, 7.38; N, 5.27. (* Peaks of minor rotamer)

N,N-Dibenzylpropanamide (5)⁸: To a suspension of *N,N*-dibenzylchloroformamide (**3a**) (518 mg, 2.00 mmol) and Cs_2CO_3 (978 mg, 3.00 mmol) in THF (4 mL) was added triethylborane (1.01 M in hexane, 3.00 mL, 3.03 mmol) followed by $\text{Pd}(\text{PPh}_3)_4$ (116 mg, 0.100 mmol). After stirring for 16 h at reflux, water was added at 0 $^\circ\text{C}$, and the product was extracted with EtOAc .

⁶ Tamaru, Y.; Yamada, Y.; Yoshida, Z.-i. *Synthesis* **1983**, 474.

⁷ Purushothaman, K. K.; Sarada, A.; Connolly, J. D.; Akinniyi, J. A. *J. Chem. Soc., Perkin Trans. I* **1979**, 3171.

⁸ Moffett, R. B.; Strube, R. E.; Skaletzky, L. *J. Med. Chem.* **1971**, 14, 1088.

The combined organic extracts were washed with brine, dried over MgSO_4 , filtered, and concentrated. Purification by silica gel column chromatography (hexane/EtOAc = 8/2), followed by distillation afforded amide **5** (371 mg, 73%) as a colorless oil: Rf 0.32 (hexane/Et₂O = 7/3); ¹H NMR (δ , CDCl_3) 7.36 (t, 2H, J = 7.3 Hz), 7.32–7.21 (m, 6H), 7.14 (d, 2H, J = 7.7 Hz), 4.61 (s, 2H), 4.44 (s, 2H), 2.44 (q, 2H, J = 7.3 Hz), 1.20 (t, 3H, J = 7.3 Hz); ¹³C NMR (δ , CDCl_3) 174.5, 137.7, 136.8, 129.1, 128.7, 128.5, 127.7, 127.5, 126.5, 49.9, 48.3, 26.5, 9.7; IR (thin film) 1650 cm^{-1} ; LRMS (FAB) m/z = 254 ($\text{M}+\text{H}^+$); Anal. Calcd for $\text{C}_{17}\text{H}_{19}\text{NO}$: C, 80.60; H, 7.56; N, 5.53. Found: C, 80.79; H, 7.67; N, 5.54.

***N,N*-Dibenzylbenzamide (6)⁹**: To a suspension of *N,N*-dibenzylchloroformamide (**3a**) (310 mg, 1.20 mmol), Cs_2CO_3 (585 mg, 1.79 mmol) and phenylboronic acid (221 mg, 1.81 mmol) in THF (7.2 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (69.3 mg, 0.0600 mmol). After stirring for 16 h at reflux, water was added at 0 °C, and the product was extracted with EtOAc. The combined organic extracts were washed with brine, dried over MgSO_4 , filtered, and concentrated. Purification by silica gel column chromatography (hexane/EtOAc = 8/2) afforded amide **6** (340 mg, 94%) as a white solid: Rf 0.21 (hexane/Et₂O); mp. 114.5–115.0 °C; ¹H NMR (δ , CDCl_3) 7.50–7.49 (m, 2H), 7.37–7.27 (m, 11H), 7.13 (br d, 2H, J = 4.9 Hz), 4.70 (s, 2H), 4.39 (s, 2H); ¹³C NMR (δ , CDCl_3) 172.5, 137.1, 136.6, 136.4, 129.8, 129.0, 128.9, 128.7, 128.6, 127.8, 127.7, 127.2, 126.9, 51.6, 46.9; IR (KBr) 1639 cm^{-1} ; LRMS (EI) m/z = 301 (M^+); Anal. Calcd for $\text{C}_{21}\text{H}_{19}\text{NO}$: C, 83.69; H, 6.35; N, 4.65. Found: C, 83.69; H, 6.44; N, 4.64.

***(E)*-*N,N*-Dibenzyl-3-phenylprop-2-enamide (7)¹⁰**: To a suspension of *N,N*-dibenzylchloroformamide (**3a**) (308 mg, 1.19 mmol), Cs_2CO_3 (590 mg, 1.81 mmol) and *(E)*-styrylboronic acid (266 mg, 1.80 mmol) in THF (7.2 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (69.3 mg, 0.0600 mmol). After stirring for 16 h at reflux, water was added at 0 °C, and the product was extracted with EtOAc. The combined organic extracts were washed with brine, dried over MgSO_4 , filtered, and concentrated. Purification by silica gel column chromatography (hexane/EtOAc = 8/2) afforded amide **6** (373 mg, 96%) as a white solid: Rf 0.48 (hexane/Et₂O = 5/5); mp. 131.5–132.0 °C; ¹H NMR (δ , CDCl_3) 7.86 (d, 1H, J = 15.3 Hz), 7.47–7.22 (m, 15H), 6.90 (d, 1H, J = 15.3 Hz), 4.72 (s, 2H), 4.61 (s, 2H); ¹³C NMR (δ , CDCl_3) 167.4, 144.0, 137.6, 136.9, 135.4, 130.0, 129.2, 129.0, 128.8, 128.6, 128.1, 127.9, 127.6, 126.8, 117.4, 50.2, 48.9; IR (KBr) 1642, 1594 cm^{-1} ; LRMS (EI) m/z = 327 (M^+); Anal. Calcd for $\text{C}_{23}\text{H}_{21}\text{NO}$: C, 84.37; H, 6.46; N, 4.28. Found: C, 84.39; H, 6.43; N, 4.13.

⁹ Flynn, D. L.; Crich, J. Z.; Devraj, R. V.; Hockerman, S. L.; Parlow, J. J.; South, M. S.; Woodard, S. J. *Am. Chem. Soc.* **1997**, *119*, 4874.

¹⁰ Kojima, S.; Inai, H.; Hidaka, T.; Fukuzaki, T.; Ohkata, K. *J. Org. Chem.* **2002**, *67*, 4093.

4ad