**One-pot Asymmetric Synthesis of Either Diastereomer of *tert*-Butanesulfinyl-protected
Amines from Ketones**

Jessica Tanuwidjaja, Hillary M. Peltier, and Jonathan A. Ellman*

Department of Chemistry, University of California, Berkeley, CA 94720

jellman@berkeley.edu

SUPPORTING INFORMATION

TABLE OF CONTENTS

General Experimental Methods	S2
General Procedure A for L-Selectride® Reductions	S2-S3
General Procedure B for NaBH₄ Reductions	S3
Synthesis, isolation, and characterization of sulfinamides 5a-j and 4d, f-j	S3-S13
References	S13
¹H NMR spectra for compounds 5b, e, i and 4j	S14-S17

General Experimental Methods. All reagents were obtained from commercial suppliers and were used without further purification. Tetrahydrofuran (THF) was passed through a column of activated alumina (type A2, 12 x 32, Purify Co.) under nitrogen pressure prior to use. All reactions were carried out in flame- or oven-dried glassware under a nitrogen atmosphere. Chromatography was carried out using 60 230-400 mesh silica gel or an HPFC purification system with prepacked silica columns and automatic gradient generator. NMR spectra (^1H and ^{13}C) were obtained at room temperature in CDCl_3 with TMS as an internal standard. Chemical shifts are reported in ppm, and coupling constants are reported in Hz. For diastereomeric mixtures, only peaks corresponding to the major diastereomer are listed. IR spectra were recorded on an FTIR spectrometer equipped with an attenuated total reflectance accessory, and only partial data are listed. Diastereomeric ratios were determined either by HPLC analysis with a silica normal phase column (Microsorb Si 100 A packing) and a Chiralpak AD-H column or by GC analysis with an Ultra-II column. MTPA amides were synthesized using a previously described method.¹ Absolute configuration of stereocenters was determined by correlation with known amines.² Optical rotation values were recorded for the amine hydrochloride salts of isolated **4a-g** and **5a-g** as previously described in the literature¹ using a polarimeter in MeOH at 25 °C and 589 nm. All other optical rotations are given for the major diastereomer of the sulfinamide products **4h-j** and **5h-j** in CHCl_3 . Products were isolated as diastereomeric mixtures unless otherwise stated. Characterization data is given for the major diastereomer.

General procedure A for L-Selectride® reductions. Ketone was added to a solution of (*R*)-**1** and $\text{Ti}(\text{OEt})_4$ in THF at rt. The reaction mixture was heated to the temperature indicated in

Table 1, and the reaction conversion was followed by TLC. Once the reaction was determined to be complete by TLC, the mixture was cooled to room temperature and then to -48 °C. L-Selectride® (1 M solution in THF) was added dropwise. The reaction mixture was allowed to warm to rt. Once the reduction was determined to be complete by TLC, the reaction mixture was cooled to 0 °C and MeOH was added dropwise until gas evolution was no longer observed. The crude reaction mixture was poured into an equal volume of brine while rapidly stirring. The resulting suspension was filtered through a plug of celite and the filter cake was washed with EtOAc. The filtrate was washed with brine and the brine layer was extracted with EtOAc (3x). The combined organic portions were dried (Na₂SO₄), filtered, and concentrated. Sulfinamides **5a-g** were purified by silica gel chromatography (hexanes/EtOAc).

General procedure B for NaBH₄ reductions. Ketone was added to a solution of (*R*)-**1** and Ti(OEt)₄ in THF at rt. The reaction mixture was heated to the temperature indicated in Table 1, and the reaction conversion was followed by TLC. Once the reaction was determined to be complete by TLC, the mixture was cooled to room temperature and then to -48 °C. The reaction mixture was added dropwise via cannula to a -48 °C suspension of NaBH₄ in a minimum amount of THF. The vessel was then rinsed with THF (2x). The reaction mixture was allowed to warm to rt. Once the reduction was determined to be complete by TLC, the reaction mixture was cooled to 0 °C and MeOH was added dropwise until gas evolution was no longer observed. The crude reaction mixture was poured into an equal volume of brine while rapidly stirring. The resulting suspension was filtered through a plug of celite and the filter cake was washed with EtOAc. The filtrate was washed with brine and the brine layer was extracted with EtOAc (3x). The combined organic portions were dried (Na₂SO₄), filtered, and concentrated. Sulfinamides **4d**, **f**, and **g** were purified by silica gel chromatography (hexanes/EtOAc).

Large scale preparation of (*R,S*)-*N*-(1-Phenylethyl)-2-methylpropylsulfonamide (4a**).** General procedure B was followed with a 0.5 M solution of (*R*)-**1** (5.43 g, 44.8 mmol, 1 equiv), Ti(OEt)₄ (21.7 mL, 89.6 mmol, 2 equiv), acetophenone (6.34 mL, 53.8 mmol, 1.2 equiv), and NaBH₄ (6.77 g, 179 mmol, 4 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 9.40 g (93%) of **4a**² as a pale yellow oil. The diastereomeric ratio was determined by HPLC analysis of crude **4a** (Microsorb-100 Å Si column, 90:10 hexanes/IPA; 1.0 L/min; 222 nm; **4a** *t_R* = 6.2 min (major diastereomer), **5a** *t_R* = 7.4 min (minor diastereomer).

(*R,S*)-*N*-(1-Phenylethyl)-2-methylpropylsulfonamide (5a**).** General procedure A was followed with a 0.5 M solution of (*R*)-**1** (61.6 mg, 0.509 mmol, 1 equiv), Ti(OEt)₄ (0.247 mL, 1.02 mmol, 2 equiv), acetophenone (0.0707 mL, 0.611 mmol, 1.2 equiv), and L-Selectride® (1.02 mL, 1.02 mmol, 2 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 102 mg (89%) of **5a** as a pale yellow crystalline solid. The diastereomeric ratio was determined by HPLC analysis of crude **5a** (Microsorb-100 Å Si column, 90:10 hexanes/IPA; 1.0 L/min; 222 nm; **4a** *t_R* = 6.3 min (minor diastereomer), **5a** *t_R* = 7.4 min (major diastereomer). ¹H NMR (400 MHz) δ 1.19 (s, 9H), 1.53 (d, 3H, *J* = 6.8 Hz), 3.41 (br d, 1H, *J* = 2.8 Hz), 4.54-4.60 (m, 1H), 1.24-7.35 (m, 5H); ¹³C (100 MHz) δ 22.6, 25.2, 54.7, 55.6, 126.9, 127.5, 128.5, 143.5. All analytical data agrees with the literature data.¹

(*R,S*)-*N*-(1-Methylpentyl)-2-methylpropylsulfonamide (5b**).** General procedure A was followed with a 0.5 M solution of (*R*)-**1** (59.0 mg, 0.487 mmol, 1 equiv), Ti(OEt)₄ (0.236 mL, 0.974 mmol, 2 equiv), 2-hexanone (0.0730 mL, 0.584 mmol, 1.2 equiv) and L-Selectride® (1.46 mL, 1.46 mmol, 3 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 72.2 mg (72%) of **5b** as a yellow oil. The diastereomeric ratio was determined by GC analysis of (*R*)- and (*S*)-MTPA derivatives of the amine hydrochloride salts of crude **5b** (Ultra II column, 100-250 °C, 2

°C/min, 20 psi; (*R*)-MTPA derivative of **4b** t_R = 34.9 min (minor diastereomer), **5b** t_R = 35.9 min (major diastereomer). Characterization data for major diastereomer: $[\alpha]_D^{23}$ -4.2° (*c* 1.5, MeOH, for amine hydrochloride); IR 1054, 1363, 1458, 1662, 2871, 2931, 2958, 3211 cm^{-1} ; ^1H NMR (400 MHz) δ 0.89 (t, 3H, J = 6.6 Hz), 1.20 (s, 9H), 1.26 (d, 3H, J = 6.4 Hz), 1.29-1.49 (m, 6H), 2.87 (br s, 1H), 3.31-3.37 (m, 1H); ^{13}C (100 MHz) δ 14.1, 22.6, 22.7, 23.3, 28.1, 37.9, 52.7, 55.7; MS (FAB): m/z $[\text{M} + \text{H}]^+$ 206. HRMS (FAB) m/z 206.1583 $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{10}\text{H}_{24}\text{NOS}$ 206.1579.

(*R,S*)-*N*-(1,2-Dimethylpropyl)-2-methylpropylsulfonamide (5c). General procedure A was followed with a 0.5 M solution of (*R*)-**1** (63.3 mg, 0.523 mmol, 1 equiv), $\text{Ti}(\text{OEt})_4$ (0.254 mL, 1.05 mmol, 2 equiv), 3-methyl-2-butanone (0.0672 mL, 0.628 mmol, 1.2 equiv), and L-Selectride® (1.46 mL, 1.46 mmol, 3 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 76.4 mg (76%) of **5c** as a pale yellow oil. The diastereomeric ratio was determined by GC analysis of (*R*)- and (*S*)- MTPA derivatives of the amine hydrochloride salts of crude **5c** (Ultra II column, 100-250 °C, 2 °C/min, 20 psi; (*R*)-MTPA derivative of **4c** t_R = 29.7min (minor diastereomer), **5c** t_R = 30.4 min (major diastereomer). ^1H NMR (400 MHz) δ 0.77 (d, 3H, J = 6.0 Hz), 0.79 (d, 3H, J = 6.8 Hz), 1.11 (d, 3H, J = 6.8 Hz), 1.11 (s, 9H), 1.57-1.65 (m, 1H), 2.76 (br d, 1H, J = 7.2 Hz), 3.05-3.15 (m, 1H); ^{13}C (100 MHz) δ 18.1, 18.2, 19.7, 22.7, 33.8, 55.8, 57.7. All analytical data agrees with the literature data.¹

(*R,S*)-*N*-(1,2,2-Trimethylpropyl)-2-methylpropylsulfonamide (5d). General procedure A was followed with a 0.5 M solution of (*R*)-**1** (59.0 mg, 0.487 mmol, 1 equiv), $\text{Ti}(\text{OEt})_4$ (0.236 mL, 0.974 mmol, 2 equiv), pinacolone (0.0745 mL, 0.584 mmol, 1.2 equiv) and L-Selectride® (1.46 mL, 1.46 mmol, 3 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 83.0 mg (83%) of **5d** as a white crystalline solid. The diastereomeric ratio was determined by GC

analysis of (*R*)- and (*S*)- MTPA derivatives of the amine hydrochloride salts of crude **5d** (Ultra II column, 100-250 °C, 2 °C/min, 20 psi; (*R*)-MTPA derivative of **4d** t_R = 31.8 min (minor diastereomer), **5d** t_R = 32.8 min (major diastereomer). mp 72-75 °C; $[\alpha]^{23}_D$ +2.3° (*c* 1.0, MeOH, for amine hydrochloride); IR 1016, 1055, 1377, 1652, 2958, 2978, 3128, 3249 cm^{-1} ; ^1H NMR (400 MHz) δ 0.89 (s, 9H), 1.23 (s, 9H), 1.25 (d, 3H, J = 6.8 Hz), 2.76 (br d, 1H, J = 9.2 Hz), 3.02-3.09 (m, 1H); ^{13}C (100 MHz) δ 18.9, 23.0, 26.4, 35.5, 56.4, 62.3; MS (FAB): m/z $[\text{M} + \text{H}]^+$ 206. Anal. Calcd for $\text{C}_{10}\text{H}_{23}\text{NOS}$: C, 58.49; H, 11.29; N, 6.82. Found: C, 58.71; H, 11.56; N, 6.81.

(*R,S*)-*N*-(1,2,2-Trimethylpropyl)-2-methylpropylsulfonamide (4d). General procedure B was followed with a 0.5 M solution of (*R*)-**1** (118 mg, 0.974 mmol, 1 equiv), $\text{Ti}(\text{OEt})_4$ (0.472 mL, 1.95 mmol, 2 equiv), pinacolone (0.150 mL, 1.17 mmol, 1.2 equiv) and NaBH_4 (147 mg, 3.90 mmol, 4 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 124 mg (62%) of **4d** as a white crystalline solid. The diastereomeric ratio was determined by GC analysis of (*R*)- and (*S*)- MTPA derivatives of the amine hydrochloride salts of crude **4d** (Ultra II column, 100-250 °C, 2 °C/min, 20 psi; (*R*)-MTPA derivative of **4d** t_R = 31.8 min (major diastereomer), **5d** t_R = 32.8 min (minor diastereomer). mp 55-58 °C; $[\alpha]^{23}_D$ -2.3° (*c* 1.0, MeOH, for amine hydrochloride); IR 1057, 1364, 1474, 1653, 2870, 2958, 3251 cm^{-1} ; ^1H NMR (400 MHz) δ 0.88 (s, 9H), 1.09 (d, 3H, J = 6.8 Hz), 1.16 (s, 9H), 3.05-3.11 (m, 1H), 3.21 (br s, 1H); ^{13}C (100 MHz) δ 16.0, 22.8, 26.3, 34.5, 55.5, 59.0; MS (FAB): m/z $[\text{M} + \text{H}]^+$ 206. Anal. Calcd for $\text{C}_{10}\text{H}_{23}\text{NOS}$: C, 58.49; H, 11.29; N, 6.82. Found: C, 58.14; H, 11.39; N, 7.00.

(*R,S*)-*N*-(1,3-Dimethylbutyl)-2-methylpropylsulfonamide (5e). General procedure A was followed with a 0.5 M solution of (*R*)-**1** (59.0 mg, 0.487 mmol, 1 equiv), $\text{Ti}(\text{OEt})_4$ (0.236 mL, 0.974 mmol, 2 equiv), 4-methyl-2-pentanone (0.0735 mL, 0.584 mmol, 1.2 equiv) and L-

Selectride® (1.46 mL, 1.46 mmol, 3 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 70.0 mg (70%) of **5e** as colorless oil. The diastereomeric ratio was determined by GC analysis of (*R*)- and (*S*)- MTPA derivatives of the amine hydrochloride salts of crude **5e** (Ultra II column, 100-200 °C, 1 °C/min, 20 psi; (*R*)-MTPA derivative of **4e** t_R = 48.3 min (minor diastereomer), **5e** t_R = 50.7 min (major diastereomer). $[\alpha]_D^{23}$ -6.1° (*c* 0.85, MeOH, for amine hydrochloride); IR 941, 1052, 1158, 1364, 1470, 1661, 2869, 2957, 3211 cm⁻¹; ¹H NMR (400 MHz) δ 0.89 (d, 3H, *J* = 6.4 Hz), 0.89 (d, 3H, *J* = 6.8 Hz), 1.15-1.23 (m, 1H), 1.20 (s, 9H), 1.27 (d, 3H, *J* = 6.4 Hz), 1.36-1.43 (m, 1H), 1.67-1.76 (m, 1H), 2.81 (br d, 1H, *J* = 8.0 Hz), 3.36-3.43 (m, 1H); ¹³C (100 MHz) δ 22.2, 22.7, 22.9, 24.0, 24.8, 47.8, 51.2, 55.8; MS (FAB): *m/z* [M + H]⁺ 206. HRMS (FAB) *m/z* 206.1581 [M + H]⁺ Calcd for C₁₀H₂₄NOS 206.1579.

(*R,S*)-*N*-(2,3-dihydro-1H-inden-1-yl)-2-methylpropylsulfonamide (5f). General procedure A was followed with a 1.0 M solution of (*R*)-**1** (103 mg, 0.850 mmol, 2 equiv), Ti(OEt)₄ (0.236 mL, 0.974 mmol, 2 equiv), 1-indanone (56.7 mg, 0.425 mmol, 1 equiv), and L-Selectride® (1.28 mL, 1.28 mmol, 3 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 26.7 mg (27%) of **5f** as a light brown solid. The diastereomeric ratio was determined by GC analysis of (*R*)- and (*S*)- MTPA derivatives of the amine hydrochloride salts of crude **5f** (Ultra II column, 100-250 °C, °C/min, 20 psi; (*R*)-MTPA derivative of **4f** t_R = 58.0 min (minor diastereomer), **5f** t_R = 58.7 min (major diastereomer). mp 88-93 °C; $[\alpha]_D^{23}$ -2.9° (*c* 1.0, MeOH, for amine hydrochloride); IR 1052, 1459, 1475, 1625, 2957, 3216 cm⁻¹; ¹H NMR (400 MHz) δ 1.27 (s, 9H), 1.94-2.03 (m, 1H), 2.67-2.77 (m, 1H), 2.81-2.89 (m, 1H), 2.96-3.03 (m, 1H), 3.35 (br d, 1H, *J* = 9.6 Hz), 4.81-4.87 (m, 1H), 7.21-7.26 (m, 3H), 7.34-7.36 (m, 1H); ¹³C NMR (100 MHz) δ 22.9, 30.4, 36.7, 56.2, 62.5, 124.5, 125.0, 126.7, 128.2, 143.1, 143.9; MS (FAB): *m/z* [M

+ H]⁺ 238. Anal. Calcd for C₁₃H₁₉NOS: C, 65.78; H, 8.07; N, 5.90. Found: C, 66.06; H, 8.32; N, 5.57.

(*R*_S,*R*)- *N*-(2,3-dihydro-1H-inden-1-yl)-2-methylpropylsulfinamide (4f). General procedure B was followed with a 2.0 M solution of (*R*)-**1** (515 mg, 4.25 mmol, 2 equiv), Ti(OEt)₄ (1.18 mL, 4.87 mmol, 2 equiv), 1-indanone (284 mg, 2.13 mmol, 1 equiv) and NaBH₄ (322 mg, 8.50 mmol, 4 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 28.0 mg (28%) of **4f** as a light brown crystalline solid. The diastereomeric ratio was determined by GC analysis of (*R*)- and (*S*)- MTPA derivatives of the amine hydrochloride salts of crude **4f** (Ultra II column, 100-250 °C, °C/min, 20 psi; (*R*)-MTPA derivative of **4f** *t*_R = 58.0 min (major diastereomer), **5f** *t*_R = 58.7 min (minor diastereomer). mp 123-126 °C; [α]_D²³ +2.8° (*c* 1.4, MeOH, for amine hydrochloride); IR 1020, 1042, 1365, 1458, 1476, 1605, 2928, 2946, 3236 cm⁻¹; ¹H NMR (400 MHz) δ 1.23 (s, 9H), 1.95-2.04 (m, 1H), 2.43-2.51 (m, 1H), 2.79-2.87 (m, 1H), 2.97-3.04 (m, 1H), 3.45 (br d, 1H, *J* = 6.4 Hz), 4.89-4.94 (m, 1H), 7.24-7.26 (m, 3H), 7.52-7.55 (m, 1H); ¹³C (100 MHz) δ 22.7, 30.1, 35.0, 55.7, 61.5, 124.8, 124.9, 126.9, 128.1, 143.5, 143.7; MS (FAB): *m/z* [M + H]⁺ 238. Anal. Calcd for C₁₃H₁₉NOS: C, 65.78; H, 8.07; N, 5.90. Found: C, 65.39; H, 8.14; N, 5.89.

(*R*_S,*S*)-*N*-(1,2,3,4-tetrahydro-1-naphthalenyl)-2-methylpropylsulfinamide (5g). General procedure A was followed with a 4.0 M solution of (*R*)-**1** (193 mg, 1.59 mmol, 2 equiv), Ti(OEt)₄ (0.386 mL, 1.59 mmol, 2 equiv), α -tetralone (0.108 mL, 0.796 mmol, 1 equiv) and L-Selectride® (2.39 mL, 2.39 mmol, 3 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 165 mg (83%) of **5g** as a pale yellow oil. The diastereomeric ratio was determined by GC analysis of (*R*)- and (*S*)- MTPA derivatives of the amine hydrochloride salts of crude **5g** (Agilent Ultra II column, 100-250 °C, 2 °C/min, 20 psi; (*R*)-MTPA derivative of **4g** *t*_R = 62.8 min (minor

diastereomer), **5g** t_R = 63.3 min (major diastereomer). $[\alpha]^{23}_D$ -2.6° (c 1.3, MeOH, for amine hydrochloride); IR 1053, 1454, 1630, 2865, 2946, 3212 cm^{-1} ; ^1H NMR (400 MHz) δ 1.27 (s, 9H), 1.79-2.01 (m, 3H), 2.29-2.37 (m, 1H), 2.70-2.86 (m, 2H), 3.39-3.42 (br d, 1H, J = 10 Hz), 4.45-4.50 (m, 1H), 7.09-7.11 (m, 1H), 7.15-7.20 (m, 2H), 7.40-7.42 (m, 1H); ^{13}C NMR (100 MHz) δ 20.1, 23.0, 29.3, 33.4, 56.1, 56.6, 126.1, 127.5, 129.0, 129.4, 137.7 (2); MS (FAB): m/z $[\text{M} + \text{H}]^+$ 252. Anal. Calcd for $\text{C}_{14}\text{H}_{21}\text{NOS}$: C, 66.89; H, 8.42; N, 5.57. Found: C, 66.65; H, 8.71; N, 5.39.

(*R,S*)-*N*-(1,2,3,4-tetrahydro-1-naphthalenyl)-2-methylpropylsulfonamide (4g).

General procedure B was followed with a 4.0 M solution of (*R*)-**1** (386 mg, 3.18 mmol, 2 equiv), $\text{Ti}(\text{OEt})_4$ (0.772 mL, 3.18 mmol, 2 equiv), α -tetralone (0.216 mL, 1.59 mmol, 1 equiv) and NaBH_4 (241 mg, 6.37 mmol, 4 equiv). Chromatography (60:40 EtOAc/hexanes) afforded 320 mg (80%) of **4g** as an off-white crystalline solid. The diastereomeric ratio was determined by GC analysis of (*R*)- and (*S*)- MTPA derivatives of the amine hydrochloride salts of crude **4g** (Agilent Ultra II column, 100-250 °C, 2 °C/min, 20 psi; (*R*)-MTPA derivative of **4g** t_R = 62.8 min (major diastereomer), **5g** t_R = 63.3 min (minor diastereomer). Characterization data for major diastereomer: mp 115-118 °C; $[\alpha]^{23}_D$ +2.6° (c 1.2, MeOH, for amine hydrochloride); IR 1045, 1453, 1625, 2861, 2938, 3200 cm^{-1} ; ^1H NMR (400 MHz) δ 1.22 (s, 9H), 1.74-1.79 (m, 1H), 1.85-2.04 (m, 3H), 2.69-2.84 (m, 2H), 3.22-3.23 (br d, 1H, J = 2.8 Hz), 4.57-4.59 (m, 1H), 7.09-7.11 (m, 1H), 7.17-7.22 (m, 2H), 7.44-7.46 (m, 1H); ^{13}C (100 MHz) δ 18.3, 22.9, 29.3, 30.8, 52.9, 55.6, 126.8, 127.8, 129.5, 129.9, 137.1, 137.9; MS (FAB): m/z $[\text{M} + \text{H}]^+$ 252. Anal. Calcd for $\text{C}_{14}\text{H}_{21}\text{NOS}$: C, 66.89; H, 8.42; N, 5.57. Found: C, 66.50; H, 8.56; N, 5.66.

(*R_S*)-*N*-(*S*)-6-Methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-methylpropane-2-sulfonamide (5h). General procedure A was followed with a 4.0 M solution of (*R*)-**1** (388 mg,

3.20 mmol, 2 equiv), Ti(OEt)₄ (0.776 mL, 3.20 mmol, 2 equiv), 6-Methoxy-1-tetralone (282 mg, 1.60 mmol, 1 equiv) and L-Selectride® (4.80 mL, 4.80 mmol, 3 equiv). Chromatography (10:90 to 100:0 EtOAc/hexanes) afforded 406 mg (79%) of **5h** as a tan solid. The diastereomeric ratio was determined by HPLC analysis crude **5h** (Chiralpak AD-H column, 95:5 hexanes/IPA; 1.0 L/min; 250 nm; **5h** t_R = 9.9 min (major diastereomer), **4h** t_R = 13.4 min (minor diastereomer). ¹H NMR (500 MHz) δ 1.24 (s, 9H), 1.72-1.84 (m, 1H), 1.84-1.99 (m, 2H), 2.21-2.34 (m, 1H), 2.64-2.73 (m, 1H), 2.73-2.84 (m, 1H), 3.43 (br d, 1H, J = 9.4 Hz), 3.76 (s, 3H), 4.37-4.45 (m, 1H), 6.58 (d, 1H, J = 2.6 Hz), 6.72 (dd, 1H, J = 8.8, 2.6 Hz), 7.29 (d, 1H, J = 10.0 Hz). All analytical data agrees with the literature data.³

(*R_S*)-*N*-(*R*)-6-Methoxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-methylpropane-2-sulfonamide (4h). General procedure B was followed with a 4.0 M solution of (*R*)-**1** (194 mg, 1.60 mmol, 2 equiv), Ti(OEt)₄ (0.388 mL, 1.60 mmol, 2 equiv), 6-Methoxy-1-tetralone (141 mg, 0.80 mmol, 1 equiv) and NaBH₄ (121 mg, 3.20 mmol, 4 equiv). Chromatography (10:90 to 100:0 EtOAc/hexanes) afforded 199 mg (89%) of **4h** as a colorless solid. The diastereomeric ratio was determined by HPLC analysis crude **4h** (Chiralpak AD-H column, 95:5 hexanes/IPA; 1.0 L/min; 250 nm; **5h** t_R = 8.9 min (minor diastereomer), **4h** t_R = 12.5 min (major diastereomer). ¹H NMR (500 MHz) δ 1.20 (s, 9H), 1.69-1.77 (m, 1H), 1.81-1.93 (m, 2H), 1.96-2.02 (m, 1H), 2.66-2.74 (m, 1H), 2.74-2.82 (m, 1H), 3.15 (br d, 1H, J = 3.0 Hz), 3.77 (s, 3H), 4.50-4.55 (m, 1H), 6.62 (d, 1H, J = 2.6 Hz), 6.76 (dd, 1H, J = 8.4, 2.5 Hz), 7.36 (d, 1H, J = 8.3 Hz). All analytical data agrees with the literature data.³

(*R_S*)-*N*-(*S*)-6-Nitro-1,2,3,4-tetrahydronaphthalen-1-yl)-2-methylpropane-2-sulfonamide (5i). General procedure A was followed with a 1.5 M solution of (*R*)-**1** (26.0 mg, 0.215 mmol, 1 equiv), Ti(OEt)₄ (0.100 mL, 0.429 mmol, 2 equiv), **2i** (41.0 mg, 0.214 mmol, 1

equiv), and L-Selectride® (0.236 mL, 0.236 mmol, 1.1 equiv) at -78 °C for the reduction. The reaction was quenched by dropwise addition of methanol at -78 °C and then was worked up according to general procedure A. Chromatography (10:90 to 100:0 EtOAc/hexanes) afforded 45.2 mg of **5i** as an orange solid. The diastereomeric ratio was determined by HPLC analysis crude **5i** (Chiralpak AD-H column, 93:7 hexanes/IPA; 1.0 L/min; 254 nm; **5i** t_R = 10.6 min (major diastereomer), **4i** t_R = 20.4 min (minor diastereomer). Characterization data for major diastereomer: mp 133-135 °C; $[\alpha]^{23}_D +14.6^\circ$ (c 1.1, CHCl₃); IR 1065, 1337, 1511, 2956, 3280 cm⁻¹; ¹H NMR (400 MHz) δ 1.28 (s, 9H), 1.82-2.04 (m, 3H), 2.34-2.47 (m, 1H), 2.79-2.98 (m, 2H), 3.40 (br d, 1H, J = 10.0 Hz), 4.47-4.59 (m, 1H), 7.58 (d, 1H, J = 8.4 Hz), 7.94-8.06 (m, 2H); ¹³C (100 MHz) δ 20.0, 23.0, 29.4, 32.8, 55.8, 56.8, 121.0, 124.3, 130.0, 139.3, 145.0, 147.2; MS (ESI): m/z [M + H]⁺ 297.

(*R*_S)-*N*-(*R*)-6-Nitro-1,2,3,4-tetrahydronaphthalen-1-yl)-2-methylpropane-2-sulfinamide (4i**).** General procedure B was followed with a 1.5 M solution of (*R*)-**1** (71.2 mg, 0.588 mmol, 2 equiv), Ti(OEt)₄ (0.143 mL, 0.588 mmol, 2 equiv), **2i** (56.2 mg, 0.294 mmol, 1 equiv) and NaBH₄ (25.0 mg, 0.662 mmol, 2.3 equiv) at -78 °C for the reduction. Chromatography (10:90 to 100:0 EtOAc/hexanes) afforded 63.7 mg of **4i** as an orange solid. The diastereomeric ratio was determined by HPLC analysis crude **4i** (Chiralpak AD-H column, 93:7 hexanes/IPA; 1.0 L/min; 254 nm; **5i** t_R = 10.2 min (minor diastereomer), **4i** t_R = 19.5 min (major diastereomer). Characterization data for major diastereomer: mp 122-125 °C; $[\alpha]^{23}_D -8.12^\circ$ (c 0.8, CHCl₃); IR 1032, 1335, 1515, 2935, 3186 cm⁻¹; ¹H NMR (500 MHz) δ 1.23 (s, 9H), 1.78-1.91 (m, 1H), 1.91-2.08 (m, 3H), 2.77-2.97 (m, 2H), 3.30 (br d, 1H, J = 6.4 Hz), 4.59-4.64 (m, 1H), 7.69-7.74 (m, 1H), 7.96-8.05 (m, 2H); ¹³C (125 MHz) δ 18.6, 22.8, 29.4, 30.9, 53.6, 56.0,

121.5, 124.3, 131.0, 139.6, 144.4, 147.2; MS (ESI): m/z $[M + H]^+$ 297. Anal. Calcd for $C_{14}H_{20}N_2O_3S$: C, 56.73; H, 6.80; N, 9.45. Found: C, 56.99; H, 7.15; N, 9.12.

(*R_s*)-*N*-(*S*)-6,7,8,9-Tetrahydro-5H-benzo[7]annulen-5-sulfinamide (5j). General procedure A was followed with a 4.0 M solution of (*R*)-**1** (388 mg, 3.20 mmol, 2 equiv), $Ti(OEt)_4$ (0.776 mL, 3.20 mmol, 2 equiv), 1-benzosuberone (0.239 mL, 1.60 mmol, 1 equiv) and L-Selectride® (4.80 mL, 4.80 mmol, 3 equiv). Chromatography (10:90 to 96:4 EtOAc/hexanes) afforded 208 mg (49%) of **5i** as a colorless crystalline solid. The diastereomeric ratio was determined by HPLC analysis crude **5j** (Chiralpak AD-H column, 97:3 hexanes/IPA; 1.0 L/min; 210 nm; **4j** t_R = 12.8 min (minor diastereomer), **5j** t_R = 15.9 min (major diastereomer). Characterization data for major diastereomer: mp 113-115 °C; $[\alpha]_D^{23}$ -88.0° (*c* 1.1, $CHCl_3$); IR 1041, 1363, 1453, 2925, 3210 cm^{-1} ; 1H NMR (400 MHz) δ 1.23 (s, 9H), 1.60-1.75 (m, 2H), 1.83-2.04 (m, 4H), 2.70-2.80 (m, 1H), 2.92-3.03 (m, 1H), 3.33 (br d, 1H, J = 4.4 Hz), 4.61-4.68 (m, 1H), 7.07-7.13 (m, 1H), 7.14-7.20 (m, 2H), 7.29-7.35 (m, 1H); ^{13}C (100 MHz) δ 22.9, 27.4, 27.8, 35.8 (2), 55.7, 59.5, 126.2, 127.7, 128.2, 130.2, 141.2, 142.3; MS (ESI): m/z $[M + H]^+$ 266. Anal. Calcd for $C_{15}H_{23}NOS$: C, 67.88; H, 8.73; N, 5.28. Found: C, 67.51; H, 9.06; N, 5.16.

(*R_s*)-*N*-(*R*)-6,7,8,9-Tetrahydro-5H-benzo[7]annulen-5-sulfinamide (4j). General procedure B was followed with a 4.0 M solution of (*R*)-**1** (388 mg, 3.20 mmol, 2 equiv), $Ti(OEt)_4$ (0.776 mL, 3.20 mmol, 2 equiv), 1-Benzosuberone (0.239 mL, 1.60 mmol, 1 equiv) and $NaBH_4$ (242 mg, 6.40 mmol, 4 equiv). Chromatography (10:90 to 96:4 EtOAc/hexanes) afforded 436 mg (100%) of **4j**. The diastereomeric ratio was determined by HPLC analysis crude **4j** (Chiralpak AD-H column, 97:3 hexanes/IPA; 1.0 L/min; 210 nm; **4j** t_R = 13.1 min (major diastereomer), **5j** t_R = 16.3 min (minor diastereomer). Characterization data for major diastereomer: $[\alpha]_D^{23}$ -18.5° (*c* 1.1, $CHCl_3$); IR 1053, 1453, 2923, 3221 cm^{-1} ; 1H NMR (500

MHz) δ 1.27 (s, 9H), 1.52-1.56 (m, 1H), 1.79-1.89 (m, 3H), 1.99-2.04 (m, 1H), 2.13-2.27 (m, 1H), 2.68-2.72 (m, 1H), 3.05-3.10 (m, 1H), 3.35 (br s, 1H), 4.26 (br d, 1H, $J = 7.0$ Hz), 7.09-7.19 (m, 3H), 7.25-7.26 (m, 1H); ^{13}C (125 MHz) δ 23.0, 26.0, 28.2, 31.6, 35.8, 56.0, 58.7, 126.5, 128.1, 128.6, 130.5, 141.9, 143.1; MS (ESI): m/z $[\text{M} + \text{H}]^+$ 266. HRMS (FAB) m/z 266.1576 $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{15}\text{H}_{23}\text{NOS}$ 266.1579.

Ketone 2i. An acetone solution of dimethyldioxirane was prepared according to a previously reported procedure.⁴ To the crude DMDO solution (70.0 mL, 0.1 M, 4 equiv) was added 6-amino-1,2,3,4-tetrahydronaphthalen-1-one (282 mg, 1.75 mmol, 1 equiv) and the mixture was stirred at rt until conversion was complete by TLC. The resulting ketone **2i** was purified by silica gel chromatography (2:98 to 50:50 EtOAc/hexanes). mp 117-118 °C; IR 1341, 1515, 1684, 2967, 3102 cm^{-1} ; ^1H NMR (500 MHz) δ 2.19-2.24 (m, 2H), 2.74 (t, 2H, $J = 6.3$ Hz), 3.10 (t, 2H, $J = 6.0$ Hz), 8.11-8.18 (m, 3H); ^{13}C (125 MHz) δ 23.0, 29.9, 39.1, 112.0, 121.6, 124.2, 129.0, 136.8, 145.9, 150.5; MS (EI): m/z 191. Anal. Calcd for $\text{C}_{10}\text{H}_9\text{NO}_3$: C, 62.82; H, 4.74; N, 7.33. Found: C, 62.65; H, 4.90; N, 7.17.

References

1. Cogan, D. A.; Liu, G.; Ellman, J. A. *Tetrahedron* **1999**, *55*, 8883.
2. Borg, G.; Cogan, D. A.; Ellman, J. A. *Tetrahedron Lett.* **1999**, *40*, 6709.
3. Colyer, J. T.; Andersen, N. G.; Tedrow, J. S.; Soukup, T. S.; Faul, M. M. *J. Org. Chem.* **2006**, *71*, 6859.
4. Adam, W.; Bialas, J.; Hadjiarapoglou, L.; *Chem. Ber.* **1991**, *124*, 2377.

AVB-400 jt-1-150diastl/major

4j
major diastereomer

S-14

7.260

Current Data Parameters
 USER jessicat
 NAME jt-1-74co1s13
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20060501
 Time 16.16
 INSTRUM AVO-400
 PROBD 5 mm QNP 1H/13
 PULPROG zg30
 TD 65536
 SOLVENT CDCl3
 NS 8
 DS 0
 SWH 8012.820 Hz
 FIDRES 0.122266 Hz
 AQ 4.0894966 sec
 RG 256
 DW 62.400 usec
 DE 6.00 usec
 TE 293.4 K
 D1 1.00000000 sec
 MCREST 0.00000000 sec
 MCWRK 0.01500000 sec

===== CHANNEL f1 =====

NUC1 1H
 P1 12.80 usec
 PL1 0.00 dB
 SFO1 400.1324700 MHz

F2 - Processing parameters

SI 65536
 SF 400.1300170 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 4.00

5b

86:14 dr

7.605
7.259
6.908

Current Data Parameters
USER jessicat
NAME jt-l-76colsp12
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060504
Time 14.23

INSTRUM av-300

PROBHD 5 mm Dual 13C/

PULPROG zg30

TD 65536

SOLVENT CDCl3

NS 8

DS 0

SWH 6172.839 Hz

FIDRES 0.094190 Hz

AQ 5.3084660 sec

RG 512

DW 81.000 usec

DE 6.00 usec

TE 294.8 K

D1 0.20000000 sec

MCREST 0.00000000 sec

MCWRRK 0.01500000 sec

===== CHANNEL f1 =====

NUC1 1H

P1 11.00 usec

PL1 -3.00 dB

SFO1 300.1318533 MHz

F2 - Processing parameters

SI 32768

SF 300.1300067 MHz

WDW EM

SSB 0

LB 0.30 Hz

GB 0

PC 4.00

3.409
3.383
2.815
1.767
1.744
1.720
1.697
1.675
1.653
1.460
1.432
1.413
1.393
1.386
1.365
1.342
1.280
1.258
1.222
1.199
1.150
0.984
0.933
0.899
0.894
0.877

5e
96:4 dr

8.0
7.5
7.0
6.5
6.0
5.5
5.0
4.5
4.0
3.5
3.0
2.5
2.0
1.5
1.0
0.5
ppm

0.95
0.80
1.05
1.20
13.00
5.71