Supporting Information for

One-pot Synthesis of α -Iodo Substituted α , β -Unsaturated
Aldehydes from Propargyl Alcohol

Shufeng Chen and Jianbo Wang*

*Beijing National Laboratory of Molecular Sciences (BNLMS), Green Chemistry Center (GCC) and
Key Laboratory of Bioorganic Chemistry and Molecular Engineering of Ministry of Education,
College of Chemistry, Peking University, Beijing 100871, China*

Contents

General.....	S1
Spectra data for 2a-k	S1-S2
Spectral for 4, 6, 8, 9, 11	S3
Alternative reaction mechanism.....	S4
References	S4

General All solvents were distilled prior to use. PhCH₃ and THF were distilled over sodium. Aqueous HI (45%, commercial available) was used directly without further purification. For chromatography, 200-300 mesh silica gel was employed. Chemical shifts are reported in ppm using tetramethylsilane as internal standard.

The stereochemistry of **2a** was investigated by NOE experiments. Irradiation of the signal for H_b led to enhancements of 4.3% for H_a.

(Z)-2-Iodo-3-*p*-tolylacrylaldehyde (2b)

Yield 66%; Yellow solid; mp: 64-65 °C; R_f = 0.36 (Petroleum ether/EtOAc, 30:1); IR (neat) 1686, 1588, 1093 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.41 (s, 3H), 7.30 (d, J = 7.8 Hz, 2H), 7.95 (d, J = 8.4 Hz, 2H), 8.06 (s, 1H), 8.75 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 21.6, 104.3, 129.2, 130.6, 131.0, 142.5, 155.8, 189.0; EI-MS (m/z , relative intensity): 272 (M^+ , 74), 257 (97), 145 (29), 115(100). Anal. calcd for C₁₀H₉IO: C, 44.14; H, 3.33. Found: C, 44.19; H, 3.36.

(Z)-2-Iodo-3-*o*-tolylacrylaldehyde (2c)

Yield 70%; Yellow solid; mp: 56-57 °C; R_f = 0.37 (Petroleum ether/EtOAc, 30:1); IR (neat) 1691, 1590, 1091 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.35 (s, 3H), 7.26~7.37 (m, 3H), 7.83 (d, J = 7.8 Hz, 1H), 8.16 (s, 1H), 8.87 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 19.8, 109.6, 125.5, 128.2, 130.3, 130.4, 134.2, 137.0, 155.3, 188.6; EI-MS (m/z , relative intensity): 272 (M^+ , 25), 257 (65), 145 (58), 115(100). HRMS calcd for C₁₀H₉IO: 271.9698. Found: 271.9696.

(Z)-2-Iodo-3-(4-methoxyphenyl)acrylaldehyde (2d)

Yield 61%; Yellow solid; mp: 57-58 °C; R_f = 0.11 (Petroleum ether/EtOAc, 30:1); IR (neat) 1683, 1582, 1509, 1259, 1178, 1093 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 3.89 (s, 3H), 7.02 (d, J = 9.0 Hz, 2H), 8.04 (s, 1H), 8.09 (d, J = 9.0 Hz, 2H), 8.74 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 55.5, 102.3, 114.0, 126.4, 132.9, 155.4, 162.5, 189.0; EI-MS (m/z , relative intensity): 288 (M^+ , 100), 257 (5), 161 (59). HRMS calcd for C₁₀H₉IO₂: 287.9647. Found: 287.9648.

(Z)-3-(4-Fluorophenyl)-2-iodoacrylaldehyde (2e)

Yield 71%; Yellow solid; mp: 47-48 °C; R_f = 0.22 (Petroleum ether/EtOAc, 30:1); IR (neat) 1688, 1596, 1505, 1087 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.18~7.23 (m, 2H), 8.04~8.09 (m, 3H), 8.79 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 105.5, 115.9 (d, J_{C-F} = 22.0 Hz), 130.2, 132.8 (d, J_{C-F} = 8.9 Hz), 154.4, 164.4 (d, J_{C-F} = 252.8 Hz), 188.8; EI-MS (m/z , relative intensity): 276 (M^+ , 100), 149 (37). Anal. calcd for C₉H₆FIO: C, 39.16; H, 2.19. Found: C, 39.22; H, 2.25.

(Z)-3-(4-Chlorophenyl)-2-iodoacrylaldehyde (2f)

Yield 53%; Yellow solid; mp: 68-69 °C; R_f = 0.20 (Petroleum ether/EtOAc, 30:1); IR (neat) 1689, 1587, 1088 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.48 (d, J = 8.7 Hz, 2H), 7.97 (d, J = 8.7 Hz, 2H), 8.06 (s, 1H), 8.80 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 106.5, 128.9, 131.6, 132.4, 137.6, 154.2, 188.7; EI-MS (m/z , relative intensity): 292 (M^+ , 83), 257 (64), 165 (33), 101(100). HRMS calcd for $\text{C}_9\text{H}_6\text{ClIO}$: 291.9152.

Found: 291.9156.

(Z)-3-(2-Bromophenyl)-2-iodoacrylaldehyde (2g)

Yield 58%; Yellow solid; mp: 51-52 °C; R_f = 0.42 (Petroleum ether/EtOAc, 30:1); IR (neat) 1692, 1600, 1089 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.35 (dt, J = 7.5, 1.8 Hz, 1H), 7.45 (dt, J = 7.5, 1.2 Hz, 1H), 7.69 (dd, J = 11.1, 1.2 Hz, 1H), 7.98 (dd, J = 7.8, 1.5 Hz, 1H), 8.19 (s, 1H), 8.90 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 110.4, 124.5, 127.1, 130.4, 131.8, 133.0, 135.1, 155.0, 188.5; EI-MS (m/z , relative intensity): 336 (M^+ , 2), 257 (100), 209 (2). Anal. calcd for $\text{C}_9\text{H}_6\text{BrIO}$: C, 32.08; H, 1.79. Found: C, 32.19; H, 2.09.

(Z)-3-(4-Phenyl)-2-iodoacrylaldehyde (2h)

Yield 54%; Yellow solid; mp: 111-112 °C; R_f = 0.14 (Petroleum ether/EtOAc, 30:1); IR (neat) 1679, 1585, 1088 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.40~7.51 (m, 3H), 7.64~7.67 (m, 2H), 7.71~7.75 (m, 2H), 8.11~8.14 (m, 3H), 8.79 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 105.3, 120.5, 127.1, 128.2, 129.0, 131.1, 132.8, 140.0, 144.3, 155.2, 189.0; EI-MS (m/z , relative intensity): 334 (M^+ , 100), 257 (7), 207 (44).

HRMS calcd for $\text{C}_{15}\text{H}_{11}\text{IO}$: 333.9855. Found: 333.9849.

(2Z, 4E)-2-Iodo-5-phenylpenta-2, 4-dienal (2i)

Yield 50%; Yellow solid; mp: 104-105 °C; R_f = 0.24 (Petroleum ether/EtOAc, 30:1); IR (neat) 1676, 1607, 1576, 1095 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.27~7.35 (m, 2H), 7.40~7.44 (m, 3H), 7.58~7.66 (m, 3H), 8.76 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 108.3, 128.0, 128.5, 129.0, 130.4, 135.3, 145.7, 154.9, 187.8; EI-MS (m/z , relative intensity): 284 (M^+ , 33), 157 (52), 128

(100). Anal. calcd for $\text{C}_{11}\text{H}_9\text{IO}$: C, 46.51; H, 3.19. Found: C, 46.48; H, 3.24.

(Z)-3-(Furan-2-yl)-2-iodoacrylaldehyde (2j)

Yield 40%; Yellow solid; mp: 106-107 °C; R_f = 0.28 (Petroleum ether/EtOAc, 30:1); IR (neat) 1670, 1593, 1465, 1100 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 6.68~6.69 (m, 1H), 7.75~7.79 (m, 2H), 8.10 (s, 1H), 8.70 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 100.8, 112.9, 118.8, 142.3, 146.6, 150.5, 187.9; EI-MS (m/z , relative intensity): 248 (M^+ , 100), 121 (43). HRMS calcd for $\text{C}_7\text{H}_5\text{IO}_2$: 247.9334. Found: 247.9336.

3-(5-Bromothiophen-2-yl)-2-iodoacrylaldehyde (2k)

This compound was isolated as a mixture of *E* and *Z* isomer. Yield 48%; Yellow solid; mp: 158-159 °C; R_f = 0.36 (Petroleum ether/EtOAc, 15:1); IR (neat) 1669, 1573, 1409, 1102 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.25~7.32 (m, 1H),

7.43~7.44 (m, 1H), 8.31 (s, 1H), 8.76(s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 84.1, 102.8, 103.4, 120.8, 130.5, 137.1, 137.3, 137.8, 139.9, 144.2, 147.3, 147.7, 187.7; EI-MS (m/z , relative intensity): 342 (M^+ , 28), 263 (86), 108 (100). HRMS calcd for $\text{C}_7\text{H}_4\text{BrIOS}$: 341.8211. Found: 341.8204.
2, 75.9, 82.3, 123.7, 127.3, 146.7, 147.7.

(E)-2,3-Diiodo-1-(4-nitrophenyl)prop-2-en-1-ol (4): Yield 60%; R_f = 0.26 (Petroleum ether/EtOAc, 10:1); IR (neat) 3505, 1573, 1604, 1517, 1343 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.76 (d, J = 5.4 Hz, 1H), 5.53 (d, J = 5.4 Hz, 1H), 7.28 (s, 1H), 7.67 (d, J = 9.0 Hz, 2H), 8.25 (d, J = 9.0 Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 76.7, 81.5, 109.8, 123.7, 126.7, 146.7, 147.7; EI-MS (m/z , relative intensity): 431 (M^+ , 65), 304 (52), 177 (24), 152 (100); HRMS calcd for $\text{C}_9\text{H}_7\text{I}_2\text{NO}_3$: 430.8516. Found: 430.8518.

2-Iodo-3-methylbut-2-enal (6):¹ Yield 30%; ^1H NMR (300 MHz, CDCl_3) δ 2.32 (s, 3H), 2.46 (s, 3H), 9.14 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 22.2, 34.2, 107.1, 164.5, 184.6.

(E)-2-Iodo-3, 5-dimethylhex-2-enal (8):² This compound was isolated as a mixture of *E* and *Z* isomer. ^1H NMR (300 MHz, CDCl_3) δ 0.95 (s, 3H), 0.97 (s, 3H), 1.92~2.01 (m, 1H), 2.29 (s, 3H), 2.75 (d, J = 7.2 Hz, 2H), 9.08 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 22.2, 28.3, 33.0, 44.8, 109.1, 168.2, 184.1.

(Z)-2-Bromo-3-phenylacrylaldehyde (9)¹

Yield 23%; ^1H NMR (300 MHz, CDCl_3) δ 7.49~7.53 (m, 3H), 7.91, (s, 1H), 8.00~8.03 (m, 2H), 9.36 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 124.2, 128.7, 130.9, 131.5, 132.9, 149.1, 187.0.

1,2-Diiodo-3-phenylprop-2-en-1-ol (11)

Yield 57%; White solid; mp: 58-59 $^{\circ}\text{C}$; R_f = 0.43 (Petroleum ether/EtOAc, 10:1); IR (neat) 3353, 1565, 1493, 1451, 698 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.48 (d, J = 5.1 Hz, 1H), 5.17 (d, J = 5.1 Hz, 1H), 7.35~7.39 (m, 5H), 7.82 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 81.7, 92.4, 126.5, 128.0, 128.5, 128.6, 139.7; EI-MS (m/z , relative intensity): 386 (M^+ , 38), 259 (22), 132 (24), 107 (100).

HRMS calcd for $\text{C}_9\text{H}_8\text{I}_2\text{O}$: 385.8665. Found: 385.8658.

One of the reviewers has suggested following alternative pathways that may explain the stereochemistry of the products.

References

- (1) Antonioletti, R.; D'Auria, M.; Piancatelli, G.; Scettri, A. *Tetrahedron Lett.* **1981**, 22, 1041-1042.
- (2) Ma, S.; Hao, X.; Huang, X. *Org. Lett.* **2003**, 5, 1217-1219.