Supporting information for

One-Pot Transformation of Trichloroacetamide into Readily Deprotectable Carbamates

Toshio Nishikawa,^{a,b*} Daisuke Urabe,^a Miho Tomita,^a Takashi Tsujimoto,^b Tomoko Iwabuchi,^a
and Minoru Isobe^{a*}

^aLaboratory of Organic Chemistry, School of Bioagricultural Sciences, Nagoya University, Chikusa, Nagoya 464-8601, Japan.

FAX (+)81-52-789-4111; E-mail; nisikawa@agr.nagoya-u.ac.jp

^bPRESTO, Japan Science and Technology Agency (JST), Honcho 4-1-8, Kawaguchi, Saitama, 332-0012, Japan

Contents

1. Preparation of substrates
2. Transformation of trichloroacetamide to carbamate

1. Preparation of substrates,

Reagent and Conditions: (a) H_5IO_6 , AcOEt, rt; (b) $NaBH_4$, EtOH, 0 °C (90% in 2 steps); (c) TBSCl, imid. DMF (96%); (d) PvCl, DMAP, pyridine (97%); (e) BzCl, pyridine (96%).

Alcohol 20. Mp 95-96 °C. $[\alpha]_D^{25} +58.7$ (c 1.04, $CHCl_3$); IR (film) ν_{max} 3464, 3250, 2992, 1717, 1523, 1376, 1217, 1084, 1008 cm^{-1} . 1H NMR (300 MHz, $CDCl_3$) δ 1.36 (3H, s, acetonide), 1.55 (3H, s, acetonide), 3.94 (1H, d, $J = 3.5$ Hz, H-4), 3.96 (1H, dd, $J = 13$ Hz, H-5), 4.18 (1H, dd, $J = 13, 3.5$ Hz, H-5'), 5.06 (1H, d, $J = 3.5$ Hz, H-2), 5.37 (1H, d, $J = 18$ Hz, $-CH=CH_A H_B$), 5.42 (1H, d, $J = 11$ Hz, $-CH=CH_A H_B$), 5.91 (1H, d, $J = 3.5$ Hz, H-1), 6.08 (1H, dd, $J = 18, 11$ Hz, $-CH=CH_2$), 9.57 (1H, br s, $-NH$). ^{13}C NMR (75 MHz, $CDCl_3$) δ 26.2, 26.7, 58.1, 70.0, 78.3, 82.5, 92.8, 104.5, 112.9, 117.4, 130.0, 161.8. HRMS (FAB⁺) calcd. for $C_{12}H_{17}O_5NCl_3$ [M+H] 360.0173, found. 360.0187.

TBS-ether 9. $[\alpha]_D^{25} +53$ (c 0.93, $CHCl_3$). IR (KBr) ν_{max} 3289, 2932, 2886, 2859, 1721, 1514, 1473, 1375, 1255, 1217, 1166, 1086, 1013, 925, 830 cm^{-1} . 1H NMR (400 MHz, $CDCl_3$) δ 0.15 (3H, s, $-SiMe$), 0.17 (3H, s, $-SiMe$), 0.92 (9H, s, t -Bu), 1.35 (3H, s, acetonide), 1.54 (3H, s, acetonide), 3.85 (1H, d, $J = 3$ Hz, H-4), 4.02 (1H, d, $J = 12.5$ Hz, H-5), 4.15 (1H, dd, $J = 12.5, 3$ Hz, H-5'), 4.99 (1H, d, $J = 3.5$ Hz, H-2), 5.35 (1H, d, $J = 17.5$ Hz, $-CH=CH_A H_B$), 5.39 (1H, d, $J = 11$ Hz, $-CH=CH_A H_B$), 5.88 (1H, d, $J = 3.5$ Hz, H-1), 6.08 (1H, dd, $J = 17.5, 11$ Hz, $-CH=CH_A H_B$), 9.18 (1H, br s, $-NH$). ^{13}C NMR (100 MHz, $CDCl_3$) δ -5.4, -4.7, 19.0, 26.1, 26.2, 26.7, 59.7, 70.0, 78.3, 82.6, 93.0, 104.4, 112.4, 116.8, 130.6, 161.4. HRMS (FAB⁺) calcd. for $C_{18}H_{31}NO_5SiCl_3$ [M+H] 474.1038, found 474.1045.

Pivalate 15. $[\alpha]_D^{25} +45$ (c 0.66, $CHCl_3$). IR (KBr) ν_{max} 3392, 2982, 1729, 1501, 1376, 1218, 1084, 1007 cm^{-1} . 1H NMR (400 MHz, $CDCl_3$) δ 1.22 (9H, s, t -Bu), 1.35 (3H, s, acetonide), 1.57 (3H, s, acetonide), 4.12 (1H, dd, $J = 12.5, 4$ Hz, H-5), 4.27 (1H, dd, $J = 5.5, 4.5$ Hz, H-4), 4.63 (1H, dd, $J = 12.5, 5.5$ Hz, H-5'), 5.08 (1H, d, $J = 3.5$ Hz, H-2), 5.34 (1H, d, $J = 17.5$ Hz, $-CH=CH_A H_B$), 5.40 (1H, d, $J = 11$ Hz, $-CH=CH_A H_B$), 5.92 (1H, d, $J = 3.5$ Hz, H-1), 6.06 (1H, dd, $J = 17.5, 11$ Hz, $-CH=CH_2$), 7.30 (1H, br s, $-NH$). ^{13}C NMR (100 MHz, $CDCl_3$) δ 26.2, 26.6, 27.2, 38.9, 59.3, 69.2, 78.4, 83.4, 92.9, 104.7, 112.8, 117.4, 130.5, 161.3, 178.1. HRMS (FAB⁺) calcd. for $C_{17}H_{25}O_6NCl_3$ [M+H] 444.0748, found. 444.0746.

Benzoate 16. $[\alpha]_D^{25} +49.1$ (c 1.59, $CHCl_3$). IR (KBr) ν_{max} 3391, 2990, 1728, 1603, 1508, 1376, 1270, 1165, 1084, 1025, 822 cm^{-1} . 1H NMR (300 MHz, $CDCl_3$) δ 1.37 (3H, s, acetonide), 1.59 (3H, s, acetonide), 4.33 (1H, dd, $J = 4.5, 3$ Hz, H-4), 4.56 (1H, dd, $J = 13, 3$ Hz, H-5), 4.80 (1H, dd, $J = 13, 4.5$ Hz, H-5'), 5.05 (1H, d, $J = 3.5$ Hz, H-2), 5.35 (1H, d, $J = 17.5$ Hz, $-CH=CH_A H_B$), 5.42 (1H, d, $J = 11$ Hz, $-CH=CH_A H_B$), 5.97 (1H, d, $J = 3.5$ Hz, H-1), 6.10 (1H, dd, $J = 17.5, 11$ Hz, $-CH=CH_2$), 7.43-7.51 (2H, m, Ar), 7.26 (1H, tt, $J = 7.5, 1.5$ Hz -Ar), 7.73 (1H, br s, $-NH$),

8.04 (2H, m, -Ar). ^{13}C NMR (100 MHz, CDCl_3) δ 26.2, 26.6, 59.8, 69.4, 77.8, 82.9, 92.6, 104.5, 112.8, 117.7, 128.5, 129.1, 129.8, 129.9, 133.6, 161.2, 166.1. HRMS (FAB+) calcd. for $\text{C}_{19}\text{H}_{21}\text{O}_6\text{NCl}_3$ $[\text{M}+\text{H}]$ 464.0435, found. 464.0432.

2. Transformation of trichloroacetamide to carbamate

Dry DMF was purchased as dehydrated DMF from Kanto Chemical Co., Inc. CuCl was dried under vacuum after washing with EtOH and Et_2O .¹

Experimental procedure for equation 1.

Trichloroacetamide **1** (50 mg, 0.13 mmol) and Na_2CO_3 (29 mg, 0.26 mmol) were placed in a reaction vessel, which was filled with nitrogen. DMF (2.0 ml) and benzyl alcohol (70 μl , 0.66 mmol) were added, and the mixture was stirred at reflux temperature for 2 h. After cooling to rt, the mixture was diluted with Et_2O (20 ml), and washed with H_2O (x1) and brine (x1), dried over anhydrous Na_2SO_4 , and concentrated. The residue was purified by flash chromatography to give benzyl carbamate **2a** (47 mg, 96%).

Typical experimental procedure exemplified for **1** to **2a** (entry 1 in Table 1).

A mixture of trichloroacetamide **1** (50 mg, 0.13 mmol) and Na_2CO_3 (29 mg, 0.26 mmol) in dry DMF (2.0 ml) was heated at reflux temperature for 40 min. (Completion of the reaction was monitored by silica gel TLC). After cooling to rt, $n\text{-Bu}_4\text{NCl}$ (75 mg, 0.27 mmol), benzyl alcohol (140 μl , 1.32 mmol) and CuCl (27 mg, 0.27 mmol) were added and the resulting mixture was stirred at rt for 45 min. and then filtered through the pad of Super-Cel. The filtrate was diluted with Et_2O , and the solution was washed with H_2O , sat. NH_4Cl solution and brine, dried over anhydrous Na_2SO_4 , and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give benzyl carbamate **2a** (41 mg, 83%).

2a (R = Bn): $[\alpha]_{\text{D}}^{27} +55.1$ (c 1.02, CHCl_3). IR (KBr) ν_{max} 3379, 2985, 2908, 1733, 1518, 1255, 1063 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 1.38 (3H, s, CH_3 of acetonide), 1.42 (3H, s, CH_3 of acetonide), 1.61-1.69 (5H, m), 2.01 (1H, m), 2.28 (1H, br d, $J = 18$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.22 (1H, br d, $J = 18$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.60 (1H, m), 3.94-4.06 (2H, m), 5.07 (2H, br s, $-\text{CH}_2\text{-Ph}$), 5.22 (1H, dd, $J = 17.5$, 1 Hz, $-\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.26 (1H, br d, $J = 11$ Hz, $-\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.37 (1H, br, $-\text{CH}=\text{C}-\text{CH}_3$), 5.85 (1H, dd, $J = 17.5$, 11 Hz, $-\text{CH}=\text{CH}_2$), 7.18 (1H, br s, $-\text{NH}$), 7.27-7.38 (5H, m, *Ph*). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.6, 26.1, 26.5, 30.2, 36.8, 44.2, 58.0, 65.6, 68.7, 109.7, 115.3, 119.5, 127.7, 128.4, 130.8, 136.2, 137.4, 155.0. EI-MS m/z 371 (M^+), 356 ($\text{M}-15$). Anal. Calcd for $\text{C}_{22}\text{H}_{29}\text{O}_4\text{N}_1$: C, 71.13; H, 7.87; N, 3.77. Found: C, 70.93; H, 7.97; N, 3.81.

2b (R = CH_2CCl_3): $[\alpha]_{\text{D}}^{28} +43.5$ (c 1.75, CHCl_3). IR (KBr) ν_{max} 3363, 2986, 2908, 1752, 1523, 1256, 1238, 1109, 1066 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 1.38 (3H, s, CH_3 of acetonide), 1.44 (3H, s, CH_3 of acetonide), 1.62-1.72 (5H, m), 2.02 (1H, m), 2.28 (1H, br d, $J = 18$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.18 (1H, br dd, $J = 18$, 6 Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.62 (1H, t, $J = 8$ Hz), 3.94-4.09 (2H, m), 4.57 (1H, d, $J =$

¹ Perrin, D. D.; Armarego, W. L. F. *Purification of Laboratory Chemicals*, 3rd ed., Pergamon Press, 1988, pp 322.

12 Hz, $-\text{CH}_\text{A}\text{H}_\text{B}\text{CCl}_3$), 4.77 (1H, d, $J = 12$ Hz, $-\text{CH}_\text{A}\text{H}_\text{B}\text{CCl}_3$), 5.28 (1H, dd, $J = 18$, 1 Hz, $-\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.29 (1H, br d, $J = 10$ Hz, $-\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.38 (1H, br, $-\text{CH}=\text{C}-\text{CH}_3$), 5.84 (1H, dd, $J = 18$, 10 Hz, $-\text{CH}=\text{CH}_2$), 7.59 (1H, br s, $-\text{NH}$). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.7, 26.0, 26.5, 30.2, 36.6, 44.2, 58.2, 68.8, 73.8, 76.7, 109.7, 115.6, 119.2, 130.9, 135.5, 152.9. EI-MS m/z 415 (M^+), 413 (M^+), 411 (M^+), 400 ($\text{M}-15$), 398 ($\text{M}-15$), 396 ($\text{M}-15$). HRMS (EI) Calcd for $\text{C}_{17}\text{H}_{24}\text{NO}_4\text{Cl}_3$ (M^+) 411.0771. Found 411.0736.

2c ($\text{R} = \text{CH}_2\text{CH}=\text{CH}_2$): $[\alpha]_\text{D}^{27} +60$ (c 0.81, CHCl_3). IR (KBr) ν_max 3384, 2986, 2907, 1733, 1521, 1255, 1066 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 1.39 (3H, s, CH_3 of acetonide), 1.43 (3H, s, CH_3 of acetonide), 1.61-1.72 (5H, m), 2.00 (1H, m), 2.27 (1H, br d, $J = 18$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.22 (1H, br d, $J = 18$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.60 (1H, m), 3.94-4.07 (2H, m), 4.52 (2H, br d, $J = 5$ Hz, $-\text{CH}_2-\text{CH}=\text{CH}_2$), 5.15-5.34 (4H, m, $-\text{CH}=\text{CH}_2 \times 2$), 5.37 (1H, br d, $J = 4$ Hz, $-\text{CH}=\text{C}-\text{CH}_3$), 5.85 (1H, dd, $J = 17$, 11 Hz, $-\text{C}-\text{CH}=\text{CH}_2$), 5.77-6.01 (1H, m, $-\text{CH}_2-\text{CH}=\text{CH}_2$), 7.15 (1H, br s, $-\text{NH}$). ^{13}C NMR (CDCl_3 , 75 MHz) δ 22.6, 26.1, 26.5, 30.2, 36.8, 44.2, 57.9, 64.6, 68.7, 109.7, 115.3, 116.8, 119.5, 130.7, 133.6, 136.2, 154.9. EI-MS m/z 321 (M^+), 306 ($\text{M}-15$). HRMS (EI) Calcd for $\text{C}_{18}\text{H}_{27}\text{NO}_4$ (M^+) 321.1940. Found 321.1927.

2d ($\text{R} = t\text{-Bu}$): $[\alpha]_\text{D}^{29} +21.1$ (c 2.46, CH_2Cl_2). IR (KBr) ν_max 3394, 2980, 2902, 1727, 1515, 1367, 1254, 1174, 1066, 862 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 1.38 (3H, s, CH_3), 1.43 (9H, s, $t\text{-Bu}$), 1.60-1.71 (8H, m, $\text{CH}_3 \times 2$, $\text{C}-\text{CH}_2-\text{CH}$), 1.92-2.06 (1H, m, $\text{CH}-\text{CH}-\text{O}$), 2.27 (1H, br d, $J = 17$ Hz, $\text{C}=\text{CH}-\text{CH}_\text{A}\text{H}_\text{B}$), 3.18 (1H, br d, $J = 17$ Hz, $\text{CH}=\text{CH}-\text{CH}_\text{A}\text{H}_\text{B}$), 3.59 (1H, td, $J = 7$, 1.5 Hz, $-\text{O}-\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}-\text{O}-$), 3.93-4.06 (2H, m, $-\text{O}-\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}-\text{O}-$), 5.24 (1H, dd, $J = 17$, 1.5 Hz, $\text{CH}=\text{H}_\text{A}\text{H}_\text{B}$), 5.26 (1H, br d, $J = 10.5$ Hz, $\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.33-5.40 (1H, m, $\text{CH}=\text{C}$), 5.84 (1H, dd, $J = 17$, 10.5 Hz, $\text{CH}=\text{CH}_2$), 6.88 (1H, br s, NH). ^{13}C NMR (CDCl_3 , 100 MHz) δ 22.7, 26.2, 26.6, 28.5, 30.3, 37.1, 44.2, 57.8, 68.7, 78.3, 109.5, 114.9, 119.6, 130.6, 136.6, 154.9. Anal. Calcd for $\text{C}_{19}\text{H}_{31}\text{O}_4\text{N}$: C, 67.63; H, 9.26; N, 4.15. Found: C, 67.63; H, 9.26; N, 4.35.

2e ($\text{R} = \text{CH}_2\text{CH}_2\text{TMS}$). $[\alpha]_\text{D}^{29} +54.1$ (c 1.38, CHCl_3). IR (KBr) ν_max 3386, 2956, 1733, 1525, 1372, 1255, 1063, 844 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 0.04 (9H, s, TMS), 0.96 (2H, t, $J = 8$ Hz, $\text{Si}-\text{CH}_2$), 1.38 (3H, s, CH_3), 1.42 (3H, s, CH_3), 1.50-1.72 (2H, m, CH_2), 1.63 (3H, s, CH_3), 1.91-2.03 (1H, m, CH), 2.24 (1H, br d, $J = 18$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.24 (1H, br d, $J = 18$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.59 (1H, td, $J = 7.5$, 1 Hz, $\text{O}-\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}-\text{O}$), 3.92-4.20 (4H, m, $\text{COO}-\text{CH}_2$, $\text{O}-\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}-\text{O}$), 5.24 (1H, dd, $J = 17$, 1.2 Hz, $\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.27 (1H, dd, $J = 10.5$, 1.5 Hz, $\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.33-5.40 (1H, m, $\text{C}=\text{CH}$), 5.84 (1H, dd, $J = 17$, 10.5 Hz, $\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 7.06 (1H, br s, NH). ^{13}C NMR (CDCl_3 , 75 MHz) δ -1.4, 17.3, 22.6, 26.2, 26.5, 30.2, 36.9, 44.3, 57.8, 62.0, 68.7, 109.7, 115.2, 119.6, 130.7, 136.3, 155.4. Anal. Calcd for $\text{C}_{20}\text{H}_{35}\text{O}_4\text{NSi}$: C, 62.95; H, 9.25; N, 3.67. Found: C, 62.96; H, 9.26; N, 3.58.

2f ($\text{R} = \text{CH}_2\text{C}_{12}\text{H}_8$). $[\alpha]_\text{D}^{28} +50.6$ (c 2.13, CHCl_3). IR (KBr) ν_max 3379, 2898, 1736, 1526, 1450, 1255 cm^{-1} . ^1H NMR (CDCl_3 , 300 MHz) δ 1.44 (3H, br s, CH_3), 1.45 (3H, br s, CH_3), 1.64 (3H, br s, CH_3), 1.54-1.76 (2H, m), 2.00 (1H, m), 2.28 (1H, br d, $J = 17$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.24 (1H, br d, $J = 17$ Hz, $\text{CH}_\text{A}\text{H}_\text{B}$), 3.62 (1H, br t, $J = 7.5$ Hz, $\text{O}-\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}-\text{O}$), 3.94-4.10 (2H, m, $\text{O}-\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}-\text{O}$), 4.14-4.26 (2H, m, $\text{CO}-\text{O}-\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}$), 4.46 (1H, br dd, $J = 13.0$, 9 Hz, $\text{CO}-\text{O}-\text{CH}_\text{A}\text{H}_\text{B}-\text{CH}$), 5.18 (1H, br d, $J = 17$ Hz, $\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.25 (1H, br d, $J = 10.5$ Hz, $\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 5.38 (1H, m, $\text{C}=\text{CH}$), 5.84 (1H, dd, $J = 17$, 10.5 Hz, $\text{CH}=\text{CH}_\text{A}\text{H}_\text{B}$), 7.30 (2H, td, $J = 7.5$, 1 Hz, aromatic), 7.34-7.44 (1H, br, NH), 7.39 (2H, td, $J = 7.5$, 0.8 Hz, aromatic), 7.63 (2H, br d, $J = 7.5$ Hz, aromatic), 7.76 (2H, d, $J = 7.5$ Hz, aromatic). ^{13}C NMR (CDCl_3 , 100 MHz) δ 22.7, 26.3,

26.6, 30.3, 36.9, 44.4, 47.4, 58.0, 65.7, 68.9, 76.8, 109.6, 115.4, 119.4, 119.9, 125.1, 126.9, 127.5, 130.7, 135.9, 141.4, 144.2, 144.3, 154.7. Anal. Calcd for $C_{29}H_{33}O_4N$: C, 75.79; H, 7.24; N, 3.05. Found: C, 75.80; H, 7.45; N, 3.01.

Urea 4. $[\alpha]_D^{28} +124$ (*c* 0.83, $CHCl_3$). IR (KBr) ν_{max} 3388, 2984, 2910, 1699, 1668, 1645, 1521, 1371, 1067 cm^{-1} . 1H NMR ($CDCl_3$, 300 MHz) δ 1.38 (3H x 2, s, CH_3 of acetonide), 1.40 (3H x 2, s, CH_3 of acetonide), 1.56-1.78 (5H x 2, m), 2.01 (1H x 2, m), 2.22 (1H x 2, br d, $J = 18$ Hz, CH_AH_B), 3.30 (1H x 2, br dd, $J = 18$, 5.5 Hz, CH_AH_B), 3.57 (1Hx2, m), 3.93-4.04 (2Hx2, m), 5.28 (1Hx2, dd, $J = 10.5$, 1.5 Hz, $-CH=CH_AH_B$), 5.30-5.38 (1Hx2, br, $-CH=C-CH_3$), 5.32 (1Hx2, dd, $J = 17.5$, 1.5 Hz, $-CH=CH_AH_B$), 5.85 (1Hx2, dd, $J = 17.5$, 10.5 Hz, $-CH=CH_2$), 6.38 (1Hx2, br s, $-NH$). ^{13}C NMR ($CDCl_3$, 75 MHz) δ 22.6, 26.4, 26.8, 30.3, 37.4, 44.6, 58.0, 68.7, 76.4, 109.5, 115.5, 119.7, 130.7, 137.2, 156.4. EI-MS m/z 500 (M^+), 485 ($M-15$). HRMS (EI) Calcd for $C_{29}H_{44}N_2O_5$ (M^+) 500.3250. Found 500.3206.

Compound 10. IR (KBr) ν_{max} 3363, 2972, 1729, 1507, 1238, 1073, 914 cm^{-1} . 1H NMR ($CDCl_3$, 300 MHz) δ 1.41 (3H, s, $NH-C-CH_3$), 1.57 (3H, br s, $CH=C-CH_3$), 1.67 (3H, d, $J = 1.0$ Hz, $CH=C-CH_3$), 1.67-1.74 (2H, m, CH_2), 1.89-2.00 (2H, m, CH_2), 4.85 (1H, s, NH), 5.04-5.14 (1H, m, $C=CH-CH_2$), 5.06 (2H, s, CH_2-Ph), 5.09 (1H, d, $J = 17.5$ Hz, $CH=CH_ACH_B$), 5.11 (1H, d, $J = 10.5$ Hz, $CH=CH_ACH_B$), 5.90 (1H, dd, $J = 17.5$, 10.5 Hz, $CH=CH_ACH_B$), 7.27-7.40 (5H, m, Ph). ^{13}C NMR ($CDCl_3$, 100 MHz) δ 17.6, 22.5, 24.5, 25.6, 39.6, 56.5, 66.1, 112.4, 123.8, 128.0, 128.1, 128.5, 131.9, 136.7, 143.1, 154.5. FAB-MS m/z 288 ($M+H$)⁺. Anal. Calcd for $C_{18}H_{25}O_2N$: C, 75.22; H, 8.77; N, 4.87. Found: C, 75.23; H, 8.67; N, 5.08.

Compound 11. $[\alpha]_D^{27} -33.8$ (*c* 1.82, $CHCl_3$). IR (KBr) ν_{max} 3343, 2956, 2928, 2857, 1726, 1510, 1308 cm^{-1} . 1H NMR ($CDCl_3$, 300 MHz) δ 0.06 (6H, s, $-OSi(CH_3)_2$), 0.89 (9H, s, $C(CH_3)_3$), 1.21 (3H, t, $J = 7$ Hz, $-OCH_2CH_3$), 3.49-3.65 (2H, m), 3.71 (1H, dd, $J = 10$, 6 Hz), 3.81 (1H, dq, $J = 10$, 7 Hz, CH_AH_B), 4.12 (1H, m), 4.45 (1H, m), 4.94 (1H, d, $J = 4$ Hz), 5.10-5.17 (1H, br), 5.12 (2H, br s, $-CH_2-Ph$), 5.62 (1H, br d, $J = 10.5$ Hz, $CH_A=CH_B$), 5.84 (1H, br d, $J = 10.5$ Hz, $CH_A=CH_B$), 7.28-7.40 (5H, m, Ph). ^{13}C NMR ($CDCl_3$, 75 MHz) δ -5.5, -5.4, 14.9, 18.2, 25.8, 47.3, 63.7, 65.4, 66.8, 68.8, 95.6, 125.3, 128.0, 128.2, 128.6, 136.5, 156.1. Anal. Calcd for $C_{22}H_{35}O_5NSi$: C, 62.68; H, 8.37; N, 3.32. Found: C, 62.68; H, 8.50; N, 3.41.

Compound 12. $[\alpha]_D^{27} +23.9$ (*c* 1.50, $CHCl_3$). IR (KBr) ν_{max} 3330, 2922, 1717, 1695, 1522, 1251 cm^{-1} . 1H NMR ($CDCl_3$, 300 MHz) δ 0.75 (3H, s, CH_3), 1.17 (1H, d, $J = 10$ Hz), 1.26 (3H, s, CH_3), 1.79 (1H, br dt, $J = 14.5$, 3 Hz), 2.01 (1H, m), 2.40-2.55 (3H, m), 4.53 (1H, br), 4.83 (1H, br s, $C=CH_AH_B$), 4.93 (1H, br, NH), 5.02 (1H, t, $J = 1.5$ Hz, $C=CH_AH_B$), 5.11 (2H, br s, $-CH_2-Ph$), 7.27-7.42 (5H, m, Ph). ^{13}C NMR ($CDCl_3$, 75 MHz) δ 22.0, 25.8, 29.7, 34.8, 39.8, 40.3, 46.2, 51.3, 66.6, 111.7, 128.1, 128.6, 136.6, 153.4, 156.0. EI-MS m/z 285 (M^+), 194 ($M-91$). Anal. Calcd for $C_{18}H_{23}O_2Si$: C, 75.76; H, 8.12; N, 4.91. Found: C, 75.67; H, 8.32; N, 5.03.

Benzyl Carbamate 13. $[\alpha]_D^{27} +41$ (*c* 0.54, $CHCl_3$). IR (KBr) ν_{max} 3358, 2989, 2938, 2897, 1736, 1509, 1374, 1216, 1166, 1074, 875 cm^{-1} . 1H NMR (400 MHz, $CDCl_3$) δ 1.33 (6H, s, acetonide), 1.43 (3H, s, acetonide), 1.54 (3H, s, acetonide), 4.02 (2H, d, $J = 7$ Hz, H-6), 4.08 (1H, d, $J = 5$ Hz, H-4), 4.39 (1H, td, $J = 7$, 5 Hz, H-5), 5.05 (1H, d, $J = 12.5$ Hz, $-CH_AH_BPh$), 5.13 (1H, d, $J = 12.5$ Hz, $-CH_AH_BPh$), 5.23 (1H, br s, H-2), 5.34 (1H, d, $J = 11$ Hz, $-CH=CH_AH_B$), 5.34 (1H, d, $J = 17$ Hz, $-CH=CH_AH_B$), 5.92 (1H, d, $J = 3.5$ Hz, H-1), 6.04 (1H, dd, $J = 17$, 11 Hz, $-CH=CH_2$), 6.33 (1H, br s, $-NH$), 7.29-7.39 (5H, m, -Ar). ^{13}C NMR (100 MHz, $CDCl_3$) δ 24.8, 26.2, 26.2, 26.8, 66.0, 66.6, 68.3, 74.7, 80.0, 84.8, 104.4, 110.1, 112.1, 116.9,

127.9, 128.1, 128.5, 133.2, 136.4, 155.2. HRMS (FAB⁺) calcd. for C₂₂H₃₀NO₇ [M+H] 420.2022, found 420.2024.

Benzyl Carbamate 14. [α]_D²⁷ +56 (c 0.98, CHCl₃). IR (KBr) $\nu_{\max}$ 3354, 2931, 1732, 1509, 1374, 1255, 1216, 1059, 1016, 836 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 0.08 (3H, s, -SiMe), 0.10 (3H, s, -SiMe), 0.87 (9H, s, *t*-Bu), 1.33 (3H, s, acetonide), 1.52 (3H, s, acetonide), 3.79 (1H, br d, *J* = 2.5 Hz, H-4), 3.92 (1H, br d, *J* = 12 Hz, H-5), 4.05 (1H, dd, *J* = 12, 3 Hz, H-5'), 5.03 (1H, d, *J* = 12.5 Hz, -CH_AH_BPh), 5.08 (1H, d, *J* = 3 Hz, H-2), 5.09 (1H, d, *J* = 12.5 Hz, -CH_AH_BPh), 5.30 (1H, d, *J* = 17.5 Hz, -CH=CH_AH_B), 5.35 (1H, d, *J* = 11 Hz, -CH=CH_AH_B), 5.88 (1H, d, *J* = 3 Hz, H-1), 6.07 (1H, dd, *J* = 17.5, 11 Hz, -CH=CH₂), 7.25-7.40 (5H, m, Ar), 7.64 (1H, br s, -NH). ¹³C NMR (100 MHz, CDCl₃) δ -5.8, -5.6, 18.1, 25.6, 26.2, 26.7, 59.3, 66.3, 68.7, 78.7, 83.4, 104.5, 112.1, 116.8, 127.0, 127.9, 128.4, 132.5, 136.7, 155.2. HRMS (FAB⁺) calcd. for C₂₄H₃₈NO₆Si [M+H] 464.2468, found 464.2490.

Amine (pivalate) 17. [α]_D²⁵ +69 (c 0.64, CHCl₃). IR (film) $\nu_{\max}$ 3603, 3381, 2981, 1732, 1644, 1482, 1374, 1285, 1217, 1166, 1073, 1017, 876 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 1.18 (9H, s, -COCMe₃), 1.32 (3H, s, acetonide), 1.55 (3H, s, acetonide), 4.13 (1H, dd, *J* = 11.5, 6.5 Hz, H-5), 4.18 (1H, d, *J* = 3.5 Hz, H-2), 4.23 (1H, dd, *J* = 6.5, 5 Hz, H-4), 4.30 (1H, dd, *J* = 11.5, 5 Hz, H-5'), 5.34 (1H, dd, *J* = 11, 1 Hz, -CH=CH_AH_B), 5.45 (1H, dd, *J* = 18, 1 Hz, -CH=CH_AH_B), 5.93 (1H, d, *J* = 3.5 Hz, H-1), 6.09 (1H, dd, *J* = 18, 11 Hz, -CH=CH₂). ¹³C NMR (100 MHz, CDCl₃) δ 26.3, 26.9, 27.1, 38.7, 61.6, 65.0, 80.0, 89.2, 104.5, 112.5, 116.5, 135.9, 178.3. HRMS (FAB⁺) calcd. for C₁₅H₂₆NO₅ [M+H] 300.1811, found 300.1811.

Amine (benzoate) 18. [α]_D²⁵ +43.8 (c 0.51, CHCl₃). IR (film) $\nu_{\max}$ 3380, 2989, 1721, 1603, 1453, 1374, 1317, 1275, 1216, 1166, 1100, 1071, 1018, 931, 876, 805 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 1.33 (3H, s, acetonide), 1.58 (3H, s, acetonide), 4.20 (1H, d, *J* = 3.5 Hz, H-2), 4.36-4.44 (2H, m, H-4, H-5), 4.54 (1H, dd, *J* = 14, 7 Hz, H-5'), 5.32 (1H, dd, *J* = 11, 1 Hz, -CH=CH_AH_B), 5.47 (1H, dd, *J* = 17.5, 1 Hz, -CH=CH_AH_B), 5.99 (1H, d, *J* = 4 Hz, H-1), 6.14 (1H, dd, *J* = 17.5, 11 Hz, -CH=CH₂), 7.40-7.46 (2H, m, Ar), 7.56 (1H, tt, *J* = 7, 1.5 Hz, Ar), 8.00-8.06 (2H, m, Ar). ¹³C NMR (75 MHz, CDCl₃) δ 26.2, 26.8, 62.2, 65.1, 80.0, 89.2, 104.7, 112.6, 116.5, 128.4, 129.8, 133.1, 135.9, 166.5. HRMS (FAB⁺) calcd. for C₁₇H₂₂NO₅ [M+H] 320.1498, found 320.1491.

Urea (benzoate) 19. [α]_D²⁶ +26.1 (c 0.75, CHCl₃). IR (film) $\nu_{\max}$ 3379, 2990, 1722, 1603, 1535, 1454, 1375, 1274, 1166, 1073, 1004, 928, 877 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 1.34 (6H, s, acetonide), 1.56 (6H, s, acetonide), 4.32-4.41 (4H, m, H-4 & H5), 4.64 (2H, dd, *J* = 8, 6.5 Hz, H-5'), 5.10 (2H, d, *J* = 3.5 Hz, H-2), 5.33-5.41 (2H, br, -NH), 5.35 (2H, d, *J* = 17.5 Hz, -CH=CH_AH_B), 5.37 (2H, d, *J* = 11 Hz, -CH=CH_AH_B), 5.90 (2H, d, *J* = 3.5 Hz, H-1), 6.14 (2H, dd, *J* = 17.5, 11 Hz, -CH=CH₂), 7.44 (4H, br t, *J* = 7.5, -Ar), 7.56 (2H, tt, *J* = 7.5, 1.5 Hz, -Ar), 8.00 (4H, m, -Ar). ¹³C NMR (100 MHz, CDCl₃) δ 26.3, 26.8, 61.0, 68.0, 78.9, 84.1, 104.8, 112.6, 117.4, 128.6, 129.4, 129.7, 132.5, 133.5, 155.8, 166.3. HRMS (FAB⁺) calcd. for C₃₅H₄₁N₂O₁₁ [M+H] 665.2711, found 665.2728.

Cyclic carbamate 21. [α]_D²³ +88 (c 0.25, CHCl₃). IR (film) $\nu_{\max}$ 3229, 3020, 1707, 1560, 1508, 1465, 1425, 1322, 1216, 1080, 999 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 1.35 (3H, s, acetonide), 1.57 (3H, s, acetonide), 4.16 (1H, dd, *J* = 12.5, 1 Hz, H-5), 4.22 (1H, br s, H-4), 4.38 (1H, d, *J* = 3 Hz, H-2), 4.43 (1H, dd, *J* = 12.5, 1.5 Hz, H-5'), 5.49 (1H, d, *J* = 10.5 Hz, CH=CH_ACH_B), 5.52 (1H, d, *J* = 17 Hz, CH=CH_ACH_B), 5.97 (1H, d, *J* = 3 Hz, H-1), 6.13 (1H, dd, *J* = 17, 10.5 Hz,

$CH=CH_ACH_B$) 6.90 (1H, br s, -NH). ^{13}C NMR (75 MHz, $CDCl_3$) δ 26.4, 27.0, 63.6, 66.9, 73.2, 86.8, 105.2, 113.3, 119.6, 133.5, 153.3. HRMS (FAB $^+$) calcd. for $C_{11}H_{16}NO_5$ [M+H] 242.1028, found 242.1023.