Appendix A. Supplementary data for
Oxidation of amitriptyline and nortriptyline by ferrate(VI): Efficiency and
reaction pathways

Yiqun Chen, Mengying Xiao, Zongping Wang^{*}, Wei Jiang, Yizhou Guo, Zizheng Liu
School of Environmental Science and Engineering, Huazhong University of Science
and Technology, Wuhan, 430074, China

Tel. +86 27 87792406; emails: yqchen@hust.edu.cn (Y. Chen),

xiao_mengy@163.com (M. Xiao), zongpingw@hust.edu.cn (Z. Wang),

jw0707@163.com (W. Jiang), gyz8824@163.com (Y. Guo), lzz@hust.edu.cn (Z. Liu)

^{*} Corresponding author

Fig. S1. Total ion chromatogram of the main oxidation products of AMI in Fe(VI) solution at pH 9.0.

Fig. S2. Full-scan (+)-ESI-MS/MS spectra of OP-102 (a), OP-294 (b), and OP-209 (c) of AMI in Fe(VI) solutions at pH 9.0.

Fig. S3. Total ion chromatogram of the main oxidation products of NOR in Fe(VI) solution at pH 9.0.

Fig. S4. Full-scan (+)-ESI-MS/MS spectra of OP-280 (a) and OP-209 (b) of NOR in Fe(VI) solutions at pH 9.0.