Supporting Information

Oxidative Coupling of Benzenes with α,β -Unsaturated Aldehydes by Pd(OAc)₂ / HPMoV /O₂ System

Tomoyuki Yamada, Satoshi Sakaguchi, and Yasutaka Ishii*

Department of Applied Chemistry, Faculty of Engineering, Kansai University, Suita, Osaka 564-8680, Japan E-mail: ishii@ipcku.kansai-u.ac.jp

Table of contents

S2: General Experiment

¹H and ¹³C-NMR, IR and MS spectral data of **4**

S3: Copies of ¹H and ¹³C-NMR spectra of **6**

S4: Copies of ¹H and ¹³C-NMR spectra of **8**

S5: ¹H and ¹³C-NMR, IR and MS spectral data of **12**

S6: ¹H and ¹³C-NMR, IR, MS spectra, and HRMS spectral data of **14**

S7: ¹H and ¹³C-NMR, IR, MS spectra, and HRMS spectral data of **16**

¹H and ¹³C-NMR, IR, MS spectra, and HRMS spectral data of **17**

IR and MS spectra, and HRMS spectral data of **18**

S8: Copies of ¹H and ¹³C-NMR spectra of **18'**

S9: IR and MS spectra, and HRMS of **20**

S10: Copies of ¹H and ¹³C-NMR spectra of **20'**

S11: ¹H and ¹³C-NMR, IR, MS spectra, and HRMS spectral data of **22**

S12: ¹H and ¹³C-NMR, IR, MS spectra, and HRMS spectral data of **23**

¹H and ¹³C-NMR, IR, MS spectra, and HRMS spectral data of **24**

General Experimental

All solvents and reagents were purchased from commercial sources and used without further treatment except for α,β -unsaturated aldehydes which were used after the distillation. TLC was performed on TLC plastic sheets F₂₅₄ silica gel 60, using UV light and I₂. GLC analysis was performed on GC equipped with a flame ionization detector using a 0.22 mm x 25 m capillary column. Mass spectra were determined at ionization energy of 70 eV on GCMS. ¹H and ¹³C-NMR were measured at 270 and 400 MHz, respectively, in CDCl₃ with Me₄Si as the internal standard.

β -Phenylcinnamaldehyde (4), [1210-39-5].

¹H-NMR (CDCl₃) δ 6.59 (d, J =7.83 Hz, 1 H), 7.27-7.67 (m, 10 H), 9.51 (d, J =7.83 Hz, 1 H);

¹³C-NMR (CDCl₃) δ 127.1, 128.2, 128.4, 128.5, 129.3, 130.3, 130.6, 136.5, 139.5, 161.9, 193.1; IR (NaCl) 699, 736, 765, 865, 1030, 1115, 1338, 1387, 1445, 1493, 1597, 1690, 2748, 2836, 3070; MS 102, 178, 207, 208.

Unfortunately, the compound **6** could not be isolated as a pure form. The NMR spectra of these mixtures have been presented.

^1H -NMR spectra of a mixture of *o*-, *m*- and *p*-6

^{13}C -NMR spectra of a mixture of *o*-, *m*- and *p*-6

2-Methylcinnamaldehyde (*o*-6), [4549-82-0].

IR (GC-IR) 747, 971, 1119, 1292, 1481, 1017, 1708, 2730, 3037; MS 73, 91, 103, 115, 131, 146.

3-Methylcinnamaldehyde (*m*-6), [66223-54-9].

IR (GC-IR) 688, 773, 970, 1116, 1281, 1482, 1629, 1707, 2726, 3039; MS 73, 91, 103, 115, 131, 146.

4-Methylcinnamaldehyde (*p*-6), [1504-75-2].

IR (GC-IR) 794, 971, 1117, 1416, 1512, 1629, 1707, 2728, 3031; MS 65, 77, 91, 103, 115, 131, 145.

Unfortunately, the compound **8** could not be isolated as a pure form. The NMR spectra of these mixtures have been presented.

¹H-NMR spectra of a mixture of *m*- and *p*-**8**

^{13}C -NMR spectra of a mixture of *m*- and *p*-6

3-*tert*-Butylcinnamaldehyde (*m*-8), [New Compound].

IR (GC-IR) 696, 783, 970, 1114, 1483, 1631, 1705, 2722, 2802, 2926; MS 131, 145, 173, 188; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{16}\text{O}$ $[\text{M}]^+$ 188.1201. Found 188.1214.

4-*tert*-Butylcinnamaldehyde (*p*-8), [84434-23-1].

IR (GC-IR) 811, 968, 1105, 1269, 1628, 1707, 2725, 2806, 2970; MS 131, 145, 173, 188; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{16}\text{O}$ $[\text{M}]^+$ 188.1201. Found 188.1215.

2,3-Dimethoxy-Cinnamaldehyde (12), [14755-98-7].

^1H -NMR (CDCl_3) δ 3.85 (s, 6 H), 6.53 (d,d, J =15.6, 7.81 Hz, 1 H), 6.82 (d, J =8.29 Hz, 1 H), 7.00 (d, J =1.95 Hz, 1 H), 7.08 (d,d, J =8.29, 1.95 Hz, 1 H), 7.33 (d, J =15.6 Hz, 1 H), 9.57 (d, J =7.81 Hz, 1 H); ^{13}C -NMR (CDCl_3) δ 55.8, 55.9, 109.8, 111.0, 123.3, 126.6, 126.9, 149.3, 151.9, 152.7, 193.4; IR (KBr) 582, 744, 774, 790, 984, 1017, 1128, 1165, 1225, 1273, 1426, 1512, 1597, 1619, 1668, 2727, 2837, 2930; MS 77, 149, 161, 177, 192.

2,3-Methylenedioxcinnamaldehyde (14a), [119491-07-5].

$^1\text{H-NMR}$ (CDCl_3) δ 6.08 (s, 2 H), 6.84-6.97 (m, 4 H), 7.41 (d, $J=15.6$ Hz, 1 H), 9.67 (d, $J=7.81$ Hz, 1 H); $^{13}\text{C-NMR}$ (CDCl_3) δ 101.6, 110.6, 116.9, 121.1, 130.8, 146.6, 146.6, 146.8, 148.1, 194.0; IR (KBr) 448, 545, 582, 603, 801, 928, 963, 1036, 118, 1260, 1449, 1493, 1598, 1625, 1670, 2708, 2835, 2919; MS 63, 89, 118, 146, 176; HRMS (EI) calcd for $\text{C}_{10}\text{H}_8\text{O}_3$ $[\text{M}]^+$ 176.0473. Found 176.0474.

3,4-Methylenedioxcinnamaldehyde (14b), [14756-00-4].

$^1\text{H-NMR}$ (CDCl_3) δ 5.96 (s, 2 H), 6.47 (d, $J=15.6$, 7.81 Hz, 1 H), 6.77 (d, $J=8.29$ Hz, 1 H), 6.97 (s, 1 H), 6.98 (d, $J=8.29$ Hz, 1 H), 7.29 (d, $J=15.6$ Hz, 1 H), 9.56 (d, $J=7.81$ Hz, 1 H); $^{13}\text{C-NMR}$ (CDCl_3) δ 101.7, 106.7, 108.6, 125.1, 126.7, 128.3, 148.4, 150.3, 152.3, 193.3; IR (KBr) 448, 545, 582, 603, 801, 928, 963, 1036, 118, 1260, 1449, 1493, 1598, 1625, 1670, 2708, 2835, 2919; MS 63, 89, 118, 147, 176; HRMS (EI) calcd for $\text{C}_{10}\text{H}_8\text{O}_3$ $[\text{M}]^+$ 176.0473. Found 176.0473.

(E)- α -Methyl-cinnamaldehyde (16a), [15174-47-7].

$^1\text{H-NMR}$ (CDCl_3) δ 2.08 (s, 3 H), 7.25 (s, 1 H), 7.53-7.83 (m, 5 H), 9.57 (s, 1 H); $^{13}\text{C-NMR}$ (CDCl_3) δ 10.9, 128.4, 128.6, 129.0, 129.4, 129.9, 135.0, 138.1, 149.6, 195.2; IR (GC-IR) 696, 779, 849, 1015, 1184, 1356, 1711, 2706, 2807, 3070; MS 91, 115, 117, 145, 146; HRMS (EI) calcd for $\text{C}_{10}\text{H}_{10}\text{O}$ $[\text{M}]^+$ 146.0732. Found 146.0731.

(Z)- α -Methyl-cinnamaldehyde (16b), [66051-14-7].

IR (GC-IR) 698, 744, 799, 1019, 1187, 1222, 1699, 2362, 2848, 3072; MS 91, 115, 117, 145, 146;

HRMS (EI) calcd for C₁₀H₁₀O [M]⁺ 146.0732. Found 146.0740.

α -Methylene-benzenepropanal (16c), [30457-88-6].

¹H-NMR (CDCl₃) δ 3.56 (s, 2 H), 6.06 (s, 1 H), 6.09 (s, 1 H), 7.31-7.15 (m, 5 H), 9.59 (s, 1 H);

¹³C-NMR (CDCl₃) δ 34.2, 126.3, 128.4, 129.0, 135.1, 138.0, 149.6, 193.8; IR (GC-IR) 697, 735, 852, 938, 1495, 1715, 2361, 2695, 2808, 3036, 3074; MS 78, 91, 115, 116, 117, 128, 145, 146.

HRMS (EI) calcd for C₁₀H₁₀O [M]⁺ 146.0732. Found 146.0734.

α -Benzyl-cinnamaldehyde (17), [838-87-9].

¹H-NMR (CDCl₃) δ 3.70 (s, 2 H), 7.21-7.42 (m, 11 H), 9.90 (s, 1 H); ¹³C-NMR (CDCl₃) δ 36.2, 126.3, 128.3, 128.4, 128.9, 129.1, 129.8, 133.9, 138.7, 141.0, 146.8, 192.2; IR (GC-IR) 697, 752, 868, 1028, 1134, 1362, 1447, 1495, 1628, 1705, 2706, 2809, 3034, 3070; MS 91, 97, 115, 116, 131, 222; HRMS (EI) calcd for C₁₆H₁₄O [M]⁺ 222.1045. Found 222.1053.

¹H-NMR (CDCl₃) δ 3.94 (s, 2 H), 7.13-7.49 (m, 11 H), 9.67 (s, 1 H); ¹³C-NMR (CDCl₃) δ 30.5, 126.1, 127.8, 128.5, 128.7, 129.6, 129.8, 134.4, 138.2, 140.4, 151.3, 194.9; MS 91, 97, 115, 116, 131, 222; HRMS (EI) calcd for C₁₆H₁₄O [M]⁺ 222.1045. Found 222.1057.

(18)

(4-*tert*-Butylphenyl)isobutyraldehyde (**18**) was obtained as a mixture of *m*- and *p*-**18'**. But IR and MS spectra could be measured. *p*-**18** could be isolated.

(1,1-Dimethylethyl)- α -methyl-benzenepropanal (*m*-18), [62518-65-4].

IR (GC-IR) 705, 783, 911, 1100, 1175, 1271, 1371, 1487, 1604, 1742, 2703, 2968, 3045; MS 57, 91, 117, 131, 147, 161, 189, 204. HRMS (EI) calcd for C₁₄H₂₀O [M]⁺ 204.1514. Found 204.1530.

3-(4-*tert*-Butylphenyl)isobutyraldehyde (p-18), [80-54-6].

$^1\text{H-NMR}$ (CDCl_3) δ 1.07 (d, $J=6.71$ Hz, 3 H), 1.30 (s, 9 H), 2.52-2.71 (m, 2H), 3.00-3.07 (m, 1H), 7.08 (d, $J=8.38$ Hz, 2 H), 7.29 (d, $J=8.83$ Hz, 2 H), 9.68 (d, $J=1.11$ Hz, 2 H); $^{13}\text{C-NMR}$ (CDCl_3) δ 13.3, 31.3, 34.3, 36.1, 48.0, 125.2, 128.5, 135.5, 148.9, 204.2; IR (GC-IR) 833, 921, 1102, 1175, 1266, 1372, 1459, 1515, 1604, 1742, 2703, 2968, 3027; MS 57, 91, 117, 131, 147, 189, 204. HRMS (EI) calcd for $\text{C}_{14}\text{H}_{20}\text{O}$ $[\text{M}]^+$ 204.1514. Found 204.1530.

Unfortunately, the compound **18'** could not be isolated as a pure form. The NMR spectra of these mixtures have been presented.

$^1\text{H-NMR}$ spectra of a mixture of *m*- and *p*-**18'**

¹³C-NMR spectra of a mixture of *m*- and *p*-**18'**

3-(1,1-Dimethylethyl)- α -methyl-benzenepropanoic acid (*m*-18'**), [New compound]**

IR (GC-IR) 705, 787, 1102, 1271, 1371, 1487, 1604, 1771, 2968, 3046, 3573; MS 91, 105, 117, 131, 147, 159, 205, 220; HRMS (EI) calcd for C₁₄H₂₀O₂ [M]⁺ 220.1463. Found 220.1493.

4-(1,1-Dimethylethyl)- α -methyl-benzenepropanoic acid (*p*-18'**) 66735-04-4**

IR (GC-IR) 787, 1017, 1102, 1267, 1371, 1464, 1513, 1772, 2967, 3027, 3573; MS 117, 131, 147, 159, 205, 220; HRMS (EI) calcd for C₁₄H₂₀O [M]⁺ 220.1463. Found 220.1478.

Methyl-*o*-isopropylhydrocinnamaldehyde (20) was obtained as a mixture of *o*-, *m*- and *p*-**20'**. But MS spectra could be measured

α -Methyl-*o*-isopropylhydrocinnamaldehyde (*o*-20**), [6502-20-1].**

HRMS (EI) calcd for C₁₃H₁₈O [M]⁺ 190.1358. Found 190.1387.

α -Methyl-*m*-(1-methylethyl)-benzenepropanal (*m*-20**), [56643-99-3].**

MS 91, 105, 117, 133, 147, 175, 190. HRMS (EI) calcd for C₁₃H₁₈O [M]⁺ 190.1358. Found 190.1387.

α -Methyl-*p*-isopropylhydrocinnamaldehyde (*p*-20), [103-95-7].

MS 91, 105, 117, 133, 147, 175, 190. HRMS (EI) calcd for C₁₃H₁₈O [M]⁺ 190.1358. Found 190.1390.

Unfortunately, the compound **20'** could not be isolated as a pure form. The NMR spectra of these mixtures have been presented.

¹H-NMR spectra of a mixture of *o*-, *m*- and *p*-20'

¹³C-NMR spectra of a mixture of *o*-, *m*- and *p*-20'

***o*-Isopropyl- α -methyl-hydrocinnamic acid (*o*-20'),** [New Compound].

MS 77, 91, 105, 119, 133, 145, 191, 206; HRMS (EI) calcd for C₁₃H₁₈O₂ [M]⁺ 206.1307. Found 206.1324.

***m*-Isopropyl- α -methyl-hydrocinnamic acid (*m*-20'),** [New Compound].

IR (GC-IR) 704, 786, 1098, 1133, 1385, 1465, 1605, 1772, 2878, 2969, 3028, 3057, 357; MS 77, 91, 105, 117, 133, 145, 160, 191, 206; HRMS (EI) calcd for C₁₃H₁₈O₂ [M]⁺ 206.1307. Found 206.1317.

***p*-Isopropyl- α -methyl-hydrocinnamic acid (*p*-20'),** [66735-06-6].

IR (GC-IR) 814, 1017, 1101, 1134, 1305, 1383, 1464, 1513, 1772, 2968, 3573; MS 91, 105, 117, 133, 145, 160, 191, 206; HRMS (EI) calcd for C₁₃H₁₈O₂ [M]⁺ 206.1307. Found 206.1318.

α ,2,5-Trimethyl-benzenepropanal (22), [6884-40-8].

¹H-NMR (CDCl₃) δ 1.10 (d, *J*=6.71 Hz, 3 H), 2.26 (s, 1 H), 2.28 (s, 1 H), 2.48-2.69 (m, 2H), 3.02-3.09 (m, 1H), 6.91 (s, 1H), 6.93 (d, *J*=7.83 Hz, 1 H), 7.03 (d, *J*=7.83 Hz, 1 H), 9.70 (s, 1 H);

¹³C-NMR (CDCl₃) δ 13.5, 19.1, 21.0, 33.9, 46.9, 127.1, 130.3, 130.3, 132.8, 135.3, 136.8, 204.3; IR

(NaCl) 805, 884, 1176, 1385, 1459, 1505, 1613, 1742, 2742, 2927, 2987 ;MS 91, 106, 119, 143, 158, 176; HRMS (EI) calcd for C₁₂H₁₆O [M]⁺ 176.1201. Found 176.1212.

2-Methoxy- α -methyl-benzenepropanal (*o*-23), [246529-82-8].

¹H-NMR (CDCl₃) δ 1.05 (d, J =6.76 Hz, 3 H), 2.59-2.77 (m, 2 H), 3.04-3.11 (m, 1 H), 3.80 (s, 3 H), 6.85 (dd, J =8.07 and 0.99 Hz, 1 H), 6.88 (ddd, J =7.58, 7.42 and 0.99 Hz, 1 H), 7.10 (dd, J =7.42 and 1.65 Hz, 1 H), 7.22 (ddd, J =8.07, 7.58 and 1.65 Hz, 1 H), 9.69 (d, J =1.65 Hz, 1 H); ¹³C-NMR (CDCl₃) δ 13.38, 31.78, 46.46, 55.10, 110.141, 120.24, 126.93, 127.71, 130.83, 157.17, 204.82; IR (GC-IR) 750, 923, 1047, 1101, 1124, 1244, 1465, 1493, 1593, 1741, 2703, 2802, 2846, 2943, 3077 ;MS 77, 91, 108, 121, 178; HRMS (EI) calcd for C₁₁H₁₄O₂ [M]⁺ 178.0994. Found 178.1000.

3-Methoxy- α -methyl-Benzenepropanal (*m*-23), [68383-30-2].

IR (GC-IR) 695, 778, 876, 898, 1049, 1161, 1264, 1453, 1488, 1601, 1743, 2704, 2801, 2848, 2947, 3006; MS 77, 91, 108, 121, 135, 150, 160, 178; HRMS (EI) calcd for C₁₁H₁₄O₂ [M]⁺ 178.0994. Found 178.1914.

2-Methyl-3-(*p*-methoxyphenyl)propionaldehyde (*p*-23), [5462-06-6].

¹H-NMR (CDCl₃) δ 1.16 (d, J =6.93 Hz, 3 H), 2.51-2.67 (m, 2 H), 2.99-3.05 (m, 1 H), 3.77 (s, 3 H), 6.83 (d, J =8.74 Hz, 2 H), 7.09 (d, J =8.74 Hz, 2 H), 9.70 (d, J =1.48 Hz, 1 H); ¹³C-NMR (CDCl₃) δ 13.19, 35.75, 48.24, 55.19, 113.73, 129.73, 130.562, 157.894, 204.48; IR (GC-IR) 834, 922, 1041, 1176, 1249, 1465, 1512, 1611, 1742, 2702, 2798, 2845, 2940, 3004; MS 77, 121, 178; HRMS (EI) calcd for C₁₁H₁₄O₂ [M]⁺ 178.0994. Found 178.1002.

α -Methyl-1,3-benzodioxole-4-propanal (24a), [New compound].

Light yellow liquid. ¹H-NMR (CDCl₃) δ 1.10 (d, J =6.71 Hz, 3 H), 2.57-2.79 (m, 2H), 2.98-3.05 (m, 1 H), 5.91 (s, 2 H), 6.62-6.62 (m, 3 H), 9.70 (s, 1 H); ¹³C-NMR (CDCl₃) δ 13.3, 30.5, 46.5, 100.5, 106.9, 120.2, 121.4, 123.0, 145.5, 146.9, 204.0; IR (GC-IR) 726, 836, 942, 1061, 1171, 1250, 1351,

1460, 1743, 2706, 2880, 2939, 2979; MS 77, 135, 164, 192; HRMS (EI) calcd for C₁₁H₁₂O₃ [M]⁺ 192.0786. Found 192.0797.

α -Methyl-1,3-benzodioxole-5-propanal (24b), [1205-17-0].

¹H-NMR (CDCl₃) δ 0.98 (d, J =6.71 Hz, 3 H), 2.39-2.55 (m, 2H), 2.86-2.93 (m, 1 H), 5.81 (s, 2 H), 6.51 (d, J =7.83 Hz, 1 H), 6.55 (s, 1 H) 6.62 (d, J =7.83 Hz, 1 H), 9.58 (s, 1 H); ¹³C-NMR (CDCl₃) δ 13.1, 36.2, 48.1, 100.7, 108.0, 109.1, 121.7, 132.2, 145.8, 147.4, 204.1; IR (GC-IR) 812, 862, 944, 1049, 1193, 1248, 1357, 1444, 1491, 1743, 2704, 2880, 2937; MS 77, 135, 192; HRMS (EI) calcd for C₁₁H₁₂O₃ [M]⁺ 192.0786. Found 192.0799.