Palladium-Catalyzed Asymmetric Allylic α -Alkylation of Acyclic Ketones

Barry M. Trost* and Jiayi Xu

*Department of Chemistry, Stanford University, Stanford, California
94305-5080. bmtrost@stanford.edu*

Supplemental Information

Experimental

General

All reactions were carried out under an atmosphere of nitrogen or argon in oven-dried glassware with magnetic stirring, unless otherwise indicated. Solvents for Pd-catalyzed reactions were distilled from sodium-benzophenone and degassed by freeze-pump-thaw techniques under vacuum¹. Solvents for other reactions were freshly purified by passage through an aluminum column before use. Tris(dibenzylideneacetone)palladium(0) monochloroform complex, Pd₂dba₃•CHCl₃ (**2**) was prepared by the procedure of Ibers². Ligands **1** were prepared by literature procedures^{3,4}. All other reagents were used as obtained unless otherwise noted.

Flash Chromatography was performed with EM Science silica gel (0.040-0.063 μ m grade). Analytical thin-layer chromatography was performed with 0.2 mm coated commercial silica gel plates (E. Merck, DC-Plasrikfolien, kieselgel 60 F254). Melting points were obtained on a Thomas-Hoover apparatus in open capillary tubes and are uncorrected. Kugelrohr distillations were performed on a Büchi GKR-50 glass tube oven. Proton nuclear magnetic resonance (¹H-NMR) data were acquired on a Mercury 400 (400 MHz) or on a Varian Unity Inova-500 (500 MHz) spectrometer. Chemical shifts are reported in delta (δ) units, in parts per million (ppm) downfield from tetramethylsilane. Splitting patterns are designated as s, singlet; d, doublet; t, triplet; q, quartet; p, pentet, m, multiplet, br, broad. Carbon-13 nuclear magnetic resonance (¹³C-NMR) data were acquired at 100 MHz on a Mercury 400 or at 125 MHz on a Varian Unity Inova 500 spectrometer. Chemical shifts are reported in ppm relative to the center line of a triplet at 77.1 ppm for chloroform-*d*. Infrared (IR) data were recorded as films on sodium chloride plates or a potassium bromide (KBr) pellets on a Perkin-Elmer Paragon 500 FT-IR spectrometer. Absorbance frequencies are reported in reciprocal centimeters (cm⁻¹). Elemental analyses (Anal.) were performed by M.-H.-W. Laboratories of Phoenix, AZ. Chiral GC analyses were performed on a HP 6890 series GC system using a Cyclosil B chiral column. GC analyses were performed on a HP 6850 series GC system using a Agilent Technologies HP-1 GC column (30 m length, 0.32 mm I.D., 0.25 μ m film). Chiral HPLC analyses were performed on a Thermo Separation Products Spectra Series P-100 or 200 and UV100 (254 nm) using Chiralcel[®] columns (OD-H, OB-H, OJ, AD, As, or OC) eluting with heptane / *iso*-propanol mixtures indicated. Optical rotations were measured on a Jasco DIP-1000 digital polarimeter using 5 cm cells and the

sodium D line (589 nm) at ambient temperature in the solvent and concentration indicated.

General Procedure for the Preparation of Allyl Enol Carbonates⁵.

Method A: To a clean dry 100 ml flask with a magnetic stirring bar was loaded 1.15 g (6.0 mmol) sodium hexamethyldisilazane (Aldrich, 97%) in a glove box under nitrogen. The flask was cooled in a dry-ice-acetone bath for 5 min and 20 ml dry THF was transferred slowly to this flask with a syringe. The flask was warmed to 0 °C to make a clear solution and 0.91 ml (6.0 mmol) TMEDA was added. The solution was cooled to -78 °C again and a solution of ketone (5.0 mmol) in 5 ml THF was added slowly in 5 min. The reaction mixture was stirred for 1 h and transferred through a cannula into a solution of 0.64 ml (6 mmol) allyl chloroformate in 5 ml THF at -78 °C and stirred for 5 min. The reaction was quenched with aqueous ammonium chloride solution and transferred into a separation funnel. Diluted with 50 ml diethyl ether and washed with brine once, the organic layer was dried over anhydrous magnesium sulfate and concentrated *in vacuo*. The product was purified with column chromatography on silica gel eluted with 5% petroleum ether / diethyl ether. Analytic pure sample was prepared by Kugelrohr distillation under high vacuum.

Method B: 0.64 ml (6 mmol) allyl chloroformate was added in one portion to the enolate solution prepared by the same procedure as described in Method A at -78 °C. The reaction was worked up and purified by using the same procedure described in Method A.

(Z)-Allyl 1-phenyl-1-propenyl carbonate: Method A; starting from 2.68 g of propiophenone (20 mmol), 2.22 g product was obtained (51%) as colorless oil. $R_f = 0.50$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max} = 3061$ (m), 2920 (m), 1760 (s), 1673 (m), 1496 (s), 1446 (s), 1366 (s), 1227 (s), 1186 (s), 966 (s), 765 (s), 693 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.5\text{--}7.2$ (m, 5H), 5.94 (m, 1H), 5.86 (q, $J = 7.2$ Hz, 1H), 5.40 (dq, $J_1 = 17.2$ Hz, $J_2 = 1.2$ Hz, 1H), 5.29 (dq, $J_1 = 10.4$ Hz, $J_2 = 1.2$ Hz, 1H), 4.68 (dt, $J_1 = 5.6$ Hz, $J_2 = 1.3$ Hz, 2H), 1.77 (d, $J = 7.2$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 152.3, 147.4, 134.8, 131.3, 128.6, 128.2, 124.3, 119.2, 113.0, 69.0, 11.3$. Anal. Calcd. for $\text{C}_{13}\text{H}_{14}\text{O}_3$: C, 71.54; H, 6.47; Found: C, 71.67; H, 6.33.

To an oven-dried flask with a stirring bar was added 0.84 ml of diisopropylamine (6 mmol) and 20 ml dry THF. The solution was cooled to -78 °C and 2.4 ml 2.5 M *n*-BuLi in hexane was added. The flask was taken out of the dry-ice bath after 5 min and put in an ice-water bath for 10 min with stirring. 0.91 ml (6.0 mmol) TMEDA was added and the solution then was cooled to -78 °C again. 701 mg (5 mmol) 1-cyclohexyl-1-propanone in 5 ml THF was slowly dropped into the solution in 10 min. The reaction was stirred for 1 h at -78 °C and 0.64 ml allyl chloroformate was added in one portion. The reaction mixture was stirred for 15 min and warmed to ambient temperature before the quenching with ammonium chloride saturated aqueous solution. Extracted with ether and the organic layer was combined and dried over anhydride magnesium sulfate. After concentration, the crude product was analyzed to be a mixture of *Z/E* isomers with a ratio of 1:0.6 by ¹H-NMR. The crude product was purified by column chromatography eluted with 2% diethyl ether in petroleum ether to yield 300 mg of *Z* isomer and 270 mg of *E* isomer and 450 mg of mixture. The total yield is 1.02 g (91%).

(*Z*)-Allyl 1-cyclohexylprop-1-enyl carbonate (*Z*-7): colorless oil. *R*_f = 0.55 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\text{max}}$ = 2929 (s), 2855 (s), 1760 (s), 1695 (m), 1450 (s), 1364 (m), 1296 (s), 1234 (s), 1187 (s), 1167 (s), 971 (s), 780 cm⁻¹ (m); ¹H NMR (500 MHz, CDCl₃): δ = 5.97 (ddt, *J*₁ = 17.1 Hz, *J*₂ = 10.4 Hz, *J*₃ = 5.8 Hz, 1H), 5.40 (dq, *J*₁ = 17.2 Hz, *J*₂ = 1.2 Hz, 1H), 5.30 (dq, *J*₁ = 10.4 Hz, *J*₂ = 1.2 Hz, 1H), 5.06 (qd, *J*₁ = 6.9 Hz, *J*₂ = 1.0 Hz, 1H), 4.67 (dt, *J*₁ = 5.8 Hz, *J*₂ = 1.2 Hz, 2H), 2.10 (m, 1H), 1.87 (m, 2H), 1.75 (m, 2H), 1.66 (m, 1H), 1.52 (dd, *J*₁ = 6.8 Hz, *J*₂ = 1.2 Hz, 3H), 1.20 (m, 5H). ¹³C NMR (125 MHz, CDCl₃): δ = 153.8, 152.9, 131.5, 119.0, 109.0, 68.7, 41.7, 30.6, 26.2, 26.1, 10.5. Anal. Calcd. for C₁₃H₂₀O₃: C, 69.61; H, 8.99; Found: C, 69.47; H, 8.79.

(*E*)-Allyl 1-cyclohexylprop-1-enyl carbonate (*E*-7): colorless oil. *R*_f = 0.50 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\text{max}}$ = 2931 (s), 2856 (s), 1760 (s), 1688 (w), 1451 (m), 1386 (m), 1363 (m), 1295 (s), 1249 (s), 1229 (s), 1173 (s), 1058 (m), 977 (m), 781 cm⁻¹ (m); ¹H NMR (500 MHz, CDCl₃): δ = 5.96 (ddt, *J*₁ = 17.1 Hz, *J*₂ = 10.4 Hz, *J*₃ = 5.8 Hz, 1H), 5.38 (dq, *J*₁ = 17.2 Hz, *J*₂ = 1.2 Hz, 1H), 5.29 (dq, *J*₁ = 10.4 Hz, *J*₂ = 1.2 Hz, 1H), 5.17 (q, *J* = 7.2 Hz, 1H), 4.64 (dt, *J*₁ = 5.7 Hz, *J*₂ = 1.2 Hz, 2H), 2.50 (m, 1H), 1.76 (m, 2H), 1.68 (d, *J*₁ = 6.8 Hz, 3H), 1.66 (m, 3H), 1.4-1.1 (m, 5H). ¹³C NMR (125 MHz, CDCl₃): δ = 154.4, 153.1, 131.6, 119.0, 111.5, 68.7, 38.5, 29.7, 26.3, 25.8, 11.4. Anal. Calcd. for C₁₃H₂₀O₃: C, 69.61; H, 8.99; Found: C, 69.71; H, 8.72.

(*Z*)-Allyl 1-phenyl-1-butenyl carbonate: Method A; starting from 1.48 g of

butyrophenone (10 mmol), 2.13 g product was obtained (91%) as colorless oil. $R_f = 0.51$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max} = 3061$ (m), 2970 (s), 2937 (m), 1760 (s), 1668 (w), 1496 (m), 1448 (m), 1300 (m), 1225 (s), 1186 (m), 973 (s), 775 (m), 693 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.5\text{--}7.2$ (m, 5H), 5.94 (m, 1H), 5.78 (t, $J = 7.2$ Hz, 1H), 5.38 (dq, $J_1 = 17.2$ Hz, $J_2 = 1.2$ Hz, 1H), 5.28 (dq, $J_1 = 10.4$ Hz, $J_2 = 1.2$ Hz, 1H), 4.67 (dt, $J_1 = 5.6$ Hz, $J_2 = 1.3$ Hz, 2H), 2.22 (q, $J = 7.2$ Hz, 2H), 1.07 (t, $J = 7.2$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 153.0, 146.0, 134.8, 131.3, 128.6, 128.2, 124.3, 120.1, 119.2, 69.0, 19.4, 13.5$. Anal. Calcd. for $\text{C}_{14}\text{H}_{16}\text{O}_3$: C, 72.39; H, 6.94; Found: C, 72.59; H, 6.81.

(Z)-Allyl 1-phenyl-1-heptenyl carbonate: Method A; starting from 951 mg of 1-phenyl-1-heptanone (5 mmol), 1.29 g product was obtained (91%) as colorless oil. $R_f = 0.49$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max} = 2957$ (s), 2930 (s), 2859 (s), 1761 (s), 1668 (w), 1496 (m), 1448 (m), 1365.1 (m), 1226 (s), 1186 (m), 965 (s), 770 (m), 693 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.5\text{--}7.2$ (m, 5H), 5.95 (m, 1H), 5.79 (t, $J = 7.2$ Hz, 1H), 5.38 (dq, $J_1 = 17.2$ Hz, $J_2 = 1.2$ Hz, 1H), 5.29 (dq, $J_1 = 10.4$ Hz, $J_2 = 1.2$ Hz, 1H), 4.67 (dt, $J_1 = 5.6$ Hz, $J_2 = 1.3$ Hz, 2H), 2.20 (q, $J = 7.2$ Hz, 2H), 1.46 (p, $J = 7.2$ Hz, 2H), 1.33 (m, 4H), 0.90 (t, $J = 7.2$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 153.0, 146.4, 134.9, 131.3, 128.6, 128.3, 124.4, 119.2, 118.8, 69.0, 31.6, 28.7, 25.9, 22.5, 14.1$. Anal. Calcd. for $\text{C}_{17}\text{H}_{22}\text{O}_3$: C, 74.42; H, 8.08; Found: C, 74.62; H, 8.26.

(Z)-Allyl 3-methyl-1-phenyl-1-butenyl carbonate: Method A; starting from 811 mg of 3-methyl-1-phenyl-1-butanone (5 mmol), 1.13 g product was obtained (92%) as colorless oil. $R_f = 0.54$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max} = 2961$ (s), 2871 (w), 1761 (s), 1669 (w), 1496 (w), 1448 (w), 1300 (m), 1228 (s), 1188 (m), 973 (s), 771 (m), 693 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.5\text{--}7.2$ (m, 5H), 5.95 (m, 1H), 5.61 (d, $J = 9.6$ Hz, 1H), 5.38 (dq, $J_1 = 17.2$ Hz, $J_2 = 1.2$ Hz, 1H), 5.29 (dq, $J_1 = 10.4$ Hz, $J_2 = 1.2$ Hz, 1H), 4.67 (dt, $J_1 = 5.6$ Hz, $J_2 = 1.3$ Hz, 2H), 2.74 (m, 1H), 1.07 (d, $J = 6.0$ Hz, 6H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 153.1, 144.8, 134.9, 131.3, 128.6, 128.2, 125.4, 124.5, 119.2, 69.0, 26.0, 22.6$. Anal. Calcd. for $\text{C}_{15}\text{H}_{18}\text{O}_3$: C, 73.15; H, 7.37; Found: C, 72.90; H, 7.20.

(Z)-Allyl 1,3-diphenyl-1-propenyl carbonate: Method A; starting from 1.05 g of β -phenylpropiophenone (5 mmol), 1.27 g product was obtained (86%) as colorless oil. R_f = 0.36 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max}$ = 3061 (m), 3028 (m), 2949 (w), 1760 (s), 1668 (w), 1495 (s), 1454 (s), 1365 (s), 1294 (s), 1227 (s), 1186 (s), 956 (s), 768 (s), 745 (s), 698 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): δ = 7.5-7.2 (m, 10H), 5.94 (m, 1H), 5.86 (q, J = 7.2 Hz, 1H), 5.38 (dq, J_1 = 17.2 Hz, J_2 = 1.2 Hz, 1H), 5.29 (dq, J_1 = 10.4 Hz, J_2 = 1.2 Hz, 1H), 4.68 (dt, J_1 = 5.6 Hz, J_2 = 1.3 Hz, 2H), 3.56 (d, J = 7.2 Hz, 2H). ^{13}C NMR (100 MHz, CDCl_3): δ = 152.9, 146.9, 139.5, 134.5, 131.2, 128.6, 128.5, 126.4, 124.6, 119.4, 117.1, 69.2, 32.2. Anal. Calcd. for $\text{C}_{19}\text{H}_{18}\text{O}_3$: C, 77.53; H, 6.16; Found: C, 77.70; H, 5.91.

(Z)-Allyl 1-(6'-methoxy-2-naphthalenyl)-1-propenyl carbonate: Method A; starting from 1.07 g of 1-(6-methoxy-2-naphthalenyl)-1-propanone (5 mmol), 1.45 g product was obtained (97%) as white crystals. Mp: 86 $^{\circ}\text{C}$; R_f = 0.33 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max}$ = 2941 (w), 1761 (s), 1228 (s), 1192 (s), 1168 (m), 981 (m), 858 (m), 820 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): δ = 7.77 (s, 1H), 7.70 (dd, J_1 = 11.6 Hz, J_2 = 8.8 Hz, 2H), 7.54 (dd, J_1 = 8.4 Hz, J_2 = 1.6 Hz, 1H), 7.12 (m, 2H), 5.98 (m, 1H), 5.95 (q, J = 7.6 Hz, 1H), 5.42 (dq, J_1 = 17.2 Hz, J_2 = 0.8 Hz, 1H), 5.31 (dq, J_1 = 10.4 Hz, J_2 = 0.8 Hz, 1H), 4.71 (d, J = 6.0 Hz, 2H), 3.90 (s, 3H), 1.82 (d, J = 7.2 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ = 158.0, 152.9, 147.5, 134.4, 131.3, 129.95, 129.91, 128.7, 127.2, 123.05, 123.00, 119.3, 119.2, 112.7, 105.7, 69.1, 55.4, 11.5. Anal. Calcd. for $\text{C}_{18}\text{H}_{18}\text{O}_3$: C, 72.47; H, 6.08; Found: C, 72.62; H, 6.15.

(Z)-Allyl 1-(2'-fluorophenyl)-1-propenyl carbonate: Method A; starting from 761 mg of 2'-fluoropropiophenone (5 mmol), 1.09 g product was obtained (93%) as colorless oil. R_f = 0.45 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max}$ = 3080 (w), 2921 (w), 1763 (s), 1491 (m), 1452 (m), 1365 (w), 1228 (s), 968 (m), 769 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): δ = 7.4-7.0 (m, 4H), 5.95 (m, 2H), 5.38 (dq, J_1 = 17.2 Hz, J_2 = 1.2 Hz, 1H), 5.29

(dq, $J_1 = 10.4$ Hz, $J_2 = 1.2$ Hz, 1H), 4.68 (dt, $J_1 = 6.0$ Hz, $J_2 = 1.3$ Hz, 2H), 1.80 (dd, $J_1 = 6.8$ Hz, $J_2 = 0.8$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 160.8, 158.3, 152.7, 142.14, 142.10, 131.2, 129.65, 129.56, 127.75, 127.72, 124.18, 124.14, 123.0, 119.2, 118.4, 118.3, 116.4, 116.2, 69.1, 11.5$. ^{19}F NMR (376 MHz, CDCl_3): $\delta = -114.6$. Anal. Calcd. for $\text{C}_{13}\text{H}_{13}\text{FO}_3$: C, 66.09; H, 5.55; F, 8.04; Found: C, 66.00; H, 5.77; F, 8.26.

(Z)-Allyl 1-(3'-chlorophenyl)-1-propenyl carbonate: Method A; starting from 843 mg of 3'-chloropropiophenone (5 mmol), 1.12 g product was obtained (89%) as colorless oil. $R_f = 0.35$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\text{max}} = 3068$ (w), 2921 (w), 1761 (s), 1596 (m), 1568 (m), 1365 (m), 1226 (s), 976 (m), 785 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.5\text{--}7.2$ (m, 4H), 5.97 (m, 1H), 5.89 (q, $J = 7.6$ Hz, 1H), 5.40 (dq, $J_1 = 17.2$ Hz, $J_2 = 1.2$ Hz, 1H), 5.31 (dq, $J_1 = 10.4$ Hz, $J_2 = 1.2$ Hz, 1H), 4.70 (dt, $J_1 = 5.6$ Hz, $J_2 = 1.3$ Hz, 2H), 1.78 (d, $J = 6.8$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 152.7, 146.1, 136.7, 134.7, 131.1, 129.9, 128.3, 124.6, 122.5, 119.4, 114.6, 69.2, 11.4$. Anal. Calcd. for $\text{C}_{13}\text{H}_{13}\text{ClO}_3$: C, 61.79; H, 5.19; Cl, 14.03; Found: C, 61.85; H, 4.93; Cl, 13.88.

(Z)-Allyl 1-(4'-bromophenyl)-1-propenyl carbonate: Method A; starting from 1.06 g of 4'-bromopropiophenone (5 mmol), 1.42 g product was obtained (98%) as colorless oil. $R_f = 0.34$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\text{max}} = 3086$ (w), 2919 (w), 1760 (s), 1671 (w), 1590 (m), 1487 (s), 1365 (m), 1224 (s), 966 (s), 825 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.46$ (dt, $J_1 = 9.2$ Hz, $J_2 = 2.2$ Hz, 2H), 7.29 (dt, $J_1 = 9.2$ Hz, $J_2 = 2.2$ Hz, 2H), 5.96 (m, 1H), 5.86 (q, $J = 7.6$ Hz, 1H), 5.40 (dq, $J_1 = 17.2$ Hz, $J_2 = 1.2$ Hz, 1H), 5.31 (dq, $J_1 = 10.4$ Hz, $J_2 = 1.2$ Hz, 1H), 4.68 (dt, $J_1 = 5.6$ Hz, $J_2 = 1.3$ Hz, 2H), 1.76 (d, $J = 6.8$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 152.7, 146.5, 133.8, 131.7, 131.1, 126.0, 122.3, 119.4, 113.9, 69.2, 11.4$. Anal. Calcd. for $\text{C}_{13}\text{H}_{13}\text{BrO}_3$: C, 52.55; H, 4.41; Br, 26.89; Found: C, 52.70; H, 4.41; Br, 27.00.

(Z)-Allyl 1-(2'-methoxyphenyl)-1-propenyl carbonate: Method A; starting from 1.27 g of 2'-methoxypropiofenone (7.7 mmol), 1.12 g product was obtained (59%) as colorless oil. $R_f = 0.22$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max} = 3078$ (m), 2947 (s), 1760 (s), 1599 (m), 1494 (s), 1456 (s), 1366 (m), 1223 (s), 964 (m), 754 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.34$ (dd, $J_1 = 7.6\text{ Hz}$, $J_2 = 2.0\text{ Hz}$, 1H), 7.26 (m, 1H), 6.90 (m, 2H), 5.94 (m, 2H), 5.37 (dq, $J_1 = 17.2\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 1H), 5.27 (dq, $J_1 = 10.4\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 1H), 4.66 (dt, $J_1 = 5.6\text{ Hz}$, $J_2 = 1.3\text{ Hz}$, 2H), 3.17 (s, 3 H), 1.77 (d, $J = 6.8\text{ Hz}$, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 156.8$, 152.8, 144.8, 131.5, 129.4, 128.5, 124.0, 120.5, 118.9, 116.9, 111.2, 68.8, 55.6, 11.5. Anal. Calcd. for $\text{C}_{14}\text{H}_{16}\text{O}_4$: C, 67.73; H, 6.50; Found: C, 67.57; H, 6.65.

(Z)-Allyl 1-(2-pyridinyl)-1-propenyl carbonate: Method A; starting from 676 mg of 1-(2-pyridinyl)-1-propanone⁶ (5 mmol), 1.12 g product was obtained (59%) as colorless oil. $R_f = 0.36$ (30% EtOAc in petroleum ether); IR (film): $\tilde{\nu}_{\max} = 3056$ (w), 2949 (w), 1760 (s), 1675 (m), 1585 (m), 1471 (m), 1434 (m), 1282 (s), 1230 (s), 976 (s), 781 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 8.56$ (m, 1H), 7.66 (td, $J_1 = 7.6\text{ Hz}$, $J_2 = 2.0\text{ Hz}$, 1H), 7.36 (d, $J = 8.2\text{ Hz}$, 1H), 7.17 (ddd, $J_1 = 7.6\text{ Hz}$, $J_2 = 5.0\text{ Hz}$, $J_3 = 1\text{ Hz}$, 1H), 6.64 (q, $J = 7.4\text{ Hz}$, 1 H), 5.98 (m, 1H), 5.42 (dq, $J_1 = 17.2\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 1H), 5.31 (dq, $J_1 = 10.4\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 1H), 4.72 (dt, $J_1 = 5.6\text{ Hz}$, $J_2 = 1.3\text{ Hz}$, 2H), 1.84 (d, $J = 7.0\text{ Hz}$, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 152.9$, 151.9, 149.4, 146.5, 136.7, 122.7, 119.3, 118.3, 116.5, 69.2, 11.4. Anal. Calcd. for $\text{C}_{12}\text{H}_{13}\text{NO}_3$: C, 65.74; H, 5.98; N, 6.39; Found: C, 65.62; H, 5.89; N, 6.52.

(Z)-Allyl 1-(3'-nitrophenyl)-1-propenyl carbonate: Method A; starting from 896 mg of 3'-nitropropiofenone (5 mmol), 1.23 g product was obtained (94%) as colorless oil. $R_f = 0.17$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max} = 3089$ (w), 2901 (w), 1760 (s), 1671 (w), 1532 (s), 1447 (m), 1352 (s), 1223 (s), 976 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 8.28$ (t, $J = 2.0\text{ Hz}$, 1H), 8.14 (ddd, $J_1 = 8.3\text{ Hz}$, $J_2 = 2.1\text{ Hz}$, $J_2 = 1\text{ Hz}$, 1H), 7.74 (ddd, $J_1 = 7.8\text{ Hz}$, $J_2 = 1.8\text{ Hz}$, $J_2 = 1.1\text{ Hz}$, 1H), 7.52 (t, $J = 8.0\text{ Hz}$, 1H), 6.05 (q, $J = 7.0\text{ Hz}$, 1H), 5.88 (m, 1H), 5.42 (dq, $J_1 = 17.2\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 1H), 5.33 (dq, $J_1 = 10.4\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 1H), 4.72 (dt, $J_1 = 5.6\text{ Hz}$, $J_2 = 1.3\text{ Hz}$, 2H), 1.83 (d, $J = 7.0\text{ Hz}$, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 152.6$, 148.5, 145.3, 136.6, 131.0, 130.0, 129.7, 122.8,

119.6, 119.3, 116.3, 69.4, 11.5. Anal. Calcd. for $C_{13}H_{13}NO_5$: C, 59.31; H, 4.98; N, 5.32; Found: C, 59.10; H, 4.88; N, 5.12.

(Z)-Allyl 1-(2-furanyl)-1-propenyl carbonate: Method A; starting from 621 mg of 1-(2-furanyl)-1-propanone (5 mmol), 0.86 g product was obtained (83%) as colorless oil. R_f = 0.43 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max}$ = 3156 (w), 2987 (w), 2920 (w), 2862 (w), 1770 (s), 1686 (w), 1494 (m), 1448 (m), 1365 (m), 1224 (s), 1162 (s), 978 (s), 739 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): δ = 7.34 (d, J = 1.8 Hz, 1H), 6.36 (dd, J_1 = 3.4 Hz, J_2 = 1.8 Hz, 1H), 6.26 (d, J = 3.4 Hz, 1H), 5.97 (m, 1H), 5.89 (q, J = 7.3 Hz, 1H), 5.41 (dq, J_1 = 17.2 Hz, J_2 = 1.2 Hz, 1H), 5.32 (dq, J_1 = 10.4 Hz, J_2 = 1.2 Hz, 1H), 4.72 (dt, J_1 = 5.6 Hz, J_2 = 1.3 Hz, 2H), 1.74 (d, J = 7.2 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ = 152.7, 148.8, 142.5, 139.5, 131.2, 119.4, 111.9, 111.3, 106.8, 69.3, 10.7. Anal. Calcd. for $C_{11}H_{12}O_4$: C, 63.45; H, 5.81; Found: C, 63.24; H, 5.62.

(Z)-Allyl 1-(2'-(trifluoromethyl)phenyl)-1-propenyl carbonate: Method A; starting from 1.01 g of 2'-trifluoromethylpropiophenone (5 mmol), 1.24 g product was obtained (87%) as colorless oil. R_f = 0.32 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max}$ = 3078 (w), 2921 (w), 1760 (s), 1450 (s), 1366 (s), 1316 (s), 1232 (s), 1171 (s), 1133 (s), 968 (s), 769 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): δ = 7.7-7.4 (m, 4H), 5.89 (m, 1H), 5.49 (q, J = 7.0 Hz, 1H), 5.33 (dq, J_1 = 17.2 Hz, J_2 = 1.2 Hz, 1H), 5.25 (dq, J_1 = 10.4 Hz, J_2 = 1.2 Hz, 1H), 4.60 (dt, J_1 = 6.0 Hz, J_2 = 1.3 Hz, 2H), 1.79 (d, J = 7.0 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ = 152.8, 145.6, 132.1, 131.9, 131.4, 129.0, 126.54, 126.48, 126.43, 122.4, 119.3, 118.1, 69.2, 11.5. ^{19}F NMR (376 MHz, CDCl_3): δ = -60.1. Anal. Calcd. for $C_{14}H_{13}F_3O_3$: C, 58.74; H, 4.58; F, 19.91; Found: C, 59.00; H, 4.36; F, 20.21.

(Z)-Allyl 1-mesityl-1-propenyl carbonate: To an oven-dried flask with a stirring bar was added 2.54 ml of HMDS (12 mmol) and 20 ml dry THF. The solution was cooled to -78°C and 5.5 ml 2.2 M *n*-BuLi in hexane was added. The flask was taken out of the dry-ice bath after 5 min and put in an ice-water bath for 10 min with stirring. The solution

then was cooled to $-78\text{ }^{\circ}\text{C}$ again and 1.76 g (10 mmol) 2',4',6'-trimethylpropiophenone in 5 ml THF was slowly dropped into the solution in 10 min. The reaction was stirred for 1 h at $-78\text{ }^{\circ}\text{C}$ and 1.28 ml allyl chloroformate was added in one portion. The reaction mixture was stirred for 1 h and warmed to ambient temperature for another 20 min before the quenching with ammonium chloride saturated aqueous solution. Extracted with ether and the organic layer was combined and dried over anhydride magnesium sulfate. Concentrated and purified by column chromatography to give 1.37 g (53%) colorless oil as a mixture of isomers with a *Z/E* ratio of 96/4 by proton NMR. $R_f = 0.47$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\text{max}} = 2921$ (s), 2863 (w), 1758 (s), 1446 (m), 1224 (s), 1162 (m), 966 (s), 926 (m), 852 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 6.86$ (s, 2H), 5.90 (m, 1H), 5.32 (dq, $J_1 = 17.4\text{ Hz}$, $J_2 = 1.4\text{ Hz}$, 1H), 5.29 (dq, $J_1 = 10.4\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 1H), 5.22 (q, $J = 7.0\text{ Hz}$, 1H), 4.58 (dt, $J_1 = 5.8\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 2H), 2.37 (s, 6H), 2.27 (s, 3H), 1.80 (d, $J = 6.9\text{ Hz}$, 3H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 152.5$, 145.4, 138.1, 137.8, 131.9, 131.5, 128.4, 119.0, 117.0, 68.7, 21.1, 20.4, 11.1. Anal. Calcd. for $\text{C}_{16}\text{H}_{20}\text{O}_3$: C, 73.82; H, 7.74; Found: C, 74.00; H, 7.86.

(E)-Allyl 1-mesityl-1-propenyl carbonate: To an oven-dried flask with a stirring bar was added 0.84 ml of diisopropylamine (6 mmol) and 5 ml dry THF. The solution was cooled to $-78\text{ }^{\circ}\text{C}$ and 2.6 ml 2.3 M *n*-BuLi in hexane was added. The flask was taken out of the dry-ice bath after 5 min and put in an ice-water bath for 10 min with stirring. The solution then was cooled to $-78\text{ }^{\circ}\text{C}$ again and 881 mg (5 mmol) 2',4',6'-trimethylpropiophenone in 2 ml THF was slowly dropped into the solution in 10 min. The reaction was stirred for 1 h at $-78\text{ }^{\circ}\text{C}$ and 1.28 ml allyl chloroformate was added in one portion. The reaction mixture was stirred for 2 h and warmed to $0\text{ }^{\circ}\text{C}$ for another 20 min before the quenching with ammonium chloride saturated aqueous solution. Extracted with ether and the organic layer was combined and dried over anhydride magnesium sulfate. Concentrated and purified by column chromatography eluted with 2% diethyl ether in petroleum ether to yield 0.83 g (64%) colorless oil as a mixture of isomers with a *Z/E* ratio of 5/95 by proton NMR. $R_f = 0.47$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\text{max}} = 2920$ (s), 2862 (w), 1758 (s), 1612 (m), 1449 (m), 1381 (m), 1363 (m), 1239 (s), 1198 (s), 1158 (s), 993 (s), 932 (s), 851 cm^{-1} (m); ^1H NMR (500 MHz, CDCl_3): $\delta = 6.87$ (s, 2H), 5.87 (m, 1H), 5.73 (q, $J = 7.2\text{ Hz}$, 1H), 5.31 (dq, $J_1 = 17.4\text{ Hz}$, $J_2 = 1.4\text{ Hz}$, 1H), 5.23 (dq, $J_1 = 10.4\text{ Hz}$, $J_2 = 1.2\text{ Hz}$, 1H), 4.55 (dt, $J_1 = 5.7\text{ Hz}$, $J_2 = 1.3\text{ Hz}$, 2H), 2.32 (s, 6H), 2.28 (s, 3H), 1.46 (d, $J = 7.2\text{ Hz}$, 3H). ^{13}C NMR (125 MHz, CDCl_3): $\delta = 153.3$, 146.0, 138.4, 138.1, 131.5, 129.1, 128.3, 119.0, 115.6, 68.6, 21.2, 19.9, 12.2. Anal. Calcd.

for C₁₆H₂₀O₃: C, 73.82; H, 7.74; Found: C, 73.94 ; H, 7.68.

(Z)-Allyl 1-cyclohexenyl-1-propenyl carbonate: Method B. starting from 520 mg of 1-cyclohexenyl-1-propanone (3.76 mmol), 0.70 g product was obtained (84%) as colorless oil. R_f = 0.41 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max}$ = 3047 (w), 2931 (s), 2861 (s), 1760 (s), 1664 (m), 1448 (s), 1364 (s), 1304 (s), 1248 (s), 1216 (s), 977 (s), 928 (s), 805 (s), 781 cm⁻¹ (s); ¹H NMR (400 MHz, CDCl₃): δ = 5.97 (ddt, J_1 = 17.1 Hz, J_2 = 10.4 Hz, J_3 = 5.8 Hz, 1H), 5.87 (bs, 1H), 5.40 (dq, J_1 = 17.2 Hz, J_2 = 1.4 Hz, 1H), 5.35 (qd, J_1 = 7.0 Hz, J_2 = 0.8 Hz, 1H), 5.30 (dq, J_1 = 10.5 Hz, J_2 = 1.2 Hz, 1H), 4.69 (dt, J_1 = 5.8 Hz, J_2 = 1.4 Hz, 2H), 2.13 (m, 4H), 1.71-1.54 (m, 4H), 1.64 (dd, J_1 = 7.2 Hz, J_2 = 0.6 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃): δ = 153.1, 148.7, 131.7, 130.4, 123.7, 119.2, 110.2, 69.0, 25.4, 24.8, 22.6, 11.2. Anal. Calcd. for C₁₃H₁₈O₃: C, 70.24; H, 8.16; Found: C, 70.50; H, 7.97.

Allyl (1E,3Z)-1-phenylpenta-1,3-dien-3-yl carbonate: Method B. starting from 801 mg of (*E*)-1-phenyl-1-penten-3-one (5 mmol), 1.10 g product was obtained (90%) as colorless oil. The Z/E ratio is 25/1 by ¹H-NMR. R_f = 0.33 (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\max}$ = 3026 (m), 2948 (m), 1760 (s), 1660 (m), 1449 (s), 1366 (s), 1231 (s), 957 cm⁻¹ (s); ¹H NMR (400 MHz, CDCl₃): δ = 7.39 (m, 2H), 7.31 (m, 2H), 7.22 (m, 1H), 6.62 (d, J = 16 Hz, 1H), 6.54 (d, J = 16 Hz, 1H), 6.00 (ddt, J_1 = 17.1 Hz, J_2 = 10.4 Hz, J_3 = 5.8 Hz, 1H), 5.52 (q, J = 7.2 Hz, 1H), 5.44 (dq, J_1 = 17.2 Hz, J_2 = 1.4 Hz, 1H), 5.33 (dq, J_1 = 10.5 Hz, J_2 = 1.2 Hz, 1H), 4.74 (dt, J_1 = 5.8 Hz, J_2 = 1.4 Hz, 2H), 1.73 (d, J = 7.2 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃): δ = 152.5, 147.2, 136.5, 131.3, 128.7, 127.9, 127.3, 126.7, 122.8, 119.3, 117.8, 69.1, 11.5. Anal. Calcd. for C₁₅H₁₆O₃: C, 73.75; H, 6.60; Found: C, 73.78; H, 6.41.

General Procedure for the Pd-catalyzed Reaction of Carbonates.

The catalyst is sensitive to air and water and thus thorough degassing of the solvent needs to be performed. Two test tubes were connected with a double-end needle. One test tube was loaded with 7.8 mg (0.75% mmol) Pd₂(dba)₃CHCl₃ (**2**) and 1.65% mmol ligand and the other one was loaded with 0.30 mmol carbonate. The system was evacuated and flushed with argon three times at which point 1.5 ml degassed solvent was added to both of the test tubes. After stirred for 20 min the orange solution of catalyst was transferred

into the test tube containing carbonate substrate. In most cases, the color of the reaction solution turned to light yellow within a few minutes and then turned back to orange in different time depending on the choice of ligand, substrate, solvent and temperature, indicating the completion of the reaction. For GC analysis, the reaction mixture was mixed with certain amount of decane as inner reference before use. Otherwise, the reaction mixture was concentrated *in vacuo* and purified by column chromatography on silica gel eluted with diethyl ether in petroleum ether

(S)-2-Methyl-1-phenyl-4-penten-1-one (4)⁶: 96%; Colorless oil. $R_f = 0.59$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{25} = +39.7$ ($c = 1.5$, CH_2Cl_2 , 94% ee); HPLC (Chiralcel[®] OD-H column; 2000:1 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 14.68$ min (minor), $t_2 = 16.25$ min (major)); IR (film): $\tilde{\nu}_{\text{max}} = 3078$ (w), 2976 (m), 2933 (w), 1682 (s), 1642 (w), 1448 (m), 1209 (s), 976 (s), 917 (m), 704 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.96$ (d, $J = 7.2$ Hz, 2H), 7.56 (t, $J = 7.6$ Hz, 1H), 7.47 (t, $J = 7.6$ Hz, 2H), 5.79 (m, 1H), 5.03 (m, 2H), 3.54 (sextet, $J = 6.8$ Hz, 1H), 2.56 (m, 1H), 2.20 (m, 1H), 1.21 (d, $J = 7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 203.7$, 136.5, 135.9, 133.0, 128.7, 128.3, 116.8, 40.5, 37.7, 17.1.

(S)-2-Methyl-1-phenylpentan-1-one⁷: 50 mg **4** (0.29 mmol) was dissolved in 2 ml methanol and was added 10 mg 5% Pd on CaCO_3 . The reaction flask was sealed and purged with H_2 three times. The reaction mixture was stirred under H_2 balloon at room temperature for 2 hours. Most solvent was removed *in vacuo* and the crude product was purified by silica gel column chromatography eluted with 2% diethyl ether in petroleum ether. 42 mg (82%) of colorless oil was obtained. $R_f = 0.55$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{25} = +25.7$ ($c = 2.83$, EtOH, 94% ee); IR (film): $\tilde{\nu}_{\text{max}} = 3064$ (w), 2961 (s), 2933 (s), 2874 (m), 1682 (s), 1596 (m), 1448 (m), 1241 (m), 1212 (s), 973 (s), 704 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.96$ (m, 2H), 7.55 (m, 1H), 7.46 (m, 2H), 3.49 (sextet, $J = 6.9$ Hz, 1H), 1.78 (m, 1H), 1.36 (m, 3H), 1.19 (d, $J = 6.9$ Hz, 3H), 0.90 (d, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 204.8$, 1367.0, 133.0, 128.8, 128.5, 40.5, 36.1, 20.8, 17.4, 14.4. (*R*)-enantiomer was reported having $[\alpha]_D^{25} = -15.8$ (EtOH).⁸

2-Ethyl-1-phenyl-4-penten-1-one⁹: 94%; Colorless oil. $R_f = 0.58$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{26} = -0.93$ ($c = 1.95$, CH_2Cl_2 , 94% ee); HPLC (Chiralcel[®] OD-H column; 2000:1 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 12.289$ min (minor), $t_2 = 13.465$ min (major)); IR (film): $\tilde{\nu}_{\text{max}} = 3078$ (w), 2965 (m), 2929 (m), 1682 (s), 1448 (m), 1222 (m), 1208 (m), 913 (m), 734 (m), 703 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.95$ (d, $J = 7.2$ Hz, 2H), 7.56 (t, $J = 7.6$ Hz, 1H), 7.47 (t, $J = 7.6$ Hz, 2H), 5.75 (m, 1H), 5.00 (m, 2H), 3.45 (pentet, $J = 6.0$ Hz, 1H), 2.52 (m, 1H), 2.27 (m, 1H), 1.81 (m, 1H), 1.61 (m, 1H), 0.88 (t, $J = 7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 203.7$, 137.5, 136.0, 133.0, 128.7, 128.3, 116.7, 47.4, 35.9, 25.0, 11.7. Spectral data was identical to the known compound.

1-Cyclohexyl-2-methyl-4-penten-1-one¹⁰: 94%; Colorless oil. $R_f = 0.58$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{22} = -30.8$ ($c = 2.35$, CH_2Cl_2 , 97% ee); IR (film): $\tilde{\nu}_{\text{max}} = 3078$ (w), 2932 (s), 2855 (s), 1708 (s), 1642 (w), 1450 (s), 1375 (m), 993 (s), 914 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 5.70$ (m, 1H), 5.01 (m, 2H), 2.74 (sextet, $J = 6.9$ Hz, 1H), 2.45 (m, 1H), 2.37 (m, 1H), 2.05 (m, 1H), 1.78 (m, 4H), 1.67 (m, 1H), 1.30 (m, 5H), 1.05 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 217.0$, 136.1, 116.7, 50.0, 44.3, 37.4, 28.6, 28.4, 25.9, 25.81, 25.78, 16.5. Anal. Calcd. for $\text{C}_{12}\text{H}_{20}\text{O}$: C, 79.94; H, 11.18; Found: C, 80.20; H, 10.99. ee was determined by the following procedure.

To the solution of 46 mg 1-cyclohexyl-2-methyl-4-penten-1-one (0.26 mmol) in 1 ml methanol was added 48 mg sodium borohydride at room temperature and stirred for 5 min. Most solvent was concentrated and the crude product was purified by column chromatography on silica gel eluted with 10% ether in petroleum ether to yield a mixture of *syn*- and *anti*- (1:1.1) 1-cyclohexyl-2-methylpent-4-en-1-ol 35 mg (75%) as colorless oil.

20 mg 1-cyclohexyl-2-methylpent-4-en-1-ol, 22 mg of (*S*)-(+)-2-methoxy-2-phenylacetic acid (aldrich, 99+% ee), 3.4 mg DMAP and 27 mg DCC was mixed with 1 ml dichloromethane. The reaction was stirred at room temperature for 1 h and filtrated through a short pad silica gel eluted with 10% diethyl ether in petroleum ether to yield 34 mg (94%) colorless oil as a mixture of *O*-methyl mandelic esters, which has no separation on TLC plate. HPLC (Chiralcel[®] AD column; 97:3 Heptane / Isopropanol; flow rate = 1 mL / min; t_1 = 5.444 min, t_2 = 6.741 min, t_3 = 7.525 min, t_4 = 9.140 min). de of 11-a and 11-d or of 11-b and 11-c reflects the ee of 1-cyclohexyl-2-methyl-4-penten-1-one.

2-Allyl-1-phenyl-1-heptanone: 93%; Colorless oil. R_f = 0.61 (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{23}$ = +1.64 (c = 1.35, CH_2Cl_2 , 92% ee); IR (film): $\tilde{\nu}_{\text{max}}$ = 3078 (m), 2927 (s), 2858 (s), 1682 (s), 1642 (m), 1448 (m), 1226 (m), 1207 (m), 915 (m), 707 (s), 688 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): δ = 7.95 (m, 2H), 7.56 (tt, J_1 = 7.2 Hz, J_2 = 1.2 Hz, 1H), 7.47 (m, 2H), 5.75 (m, 1H), 4.99 (m, 2H), 3.51 (m, 1H), 2.51 (m, 1H), 2.26 (m, 1H), 1.77 (m, 1H), 1.53 (m, 1H), 1.25 (m, 6H), 0.84 (m, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ = 203.9, 137.5, 136.0, 133.0, 128.7, 128.3, 116.7, 45.9, 36.4, 32.02, 31.99, 27.1, 22.5, 14.1. Anal. Calcd. for $\text{C}_{16}\text{H}_{22}\text{O}$: C, 83.43; H, 9.63; Found: C, 83.50; H, 9.89.

To the solution of 44 mg 2-allyl-1-phenyl-1-heptanone (0.19 mmol) in 1 ml ethyl acetate was added 96 mg sodium periodate (0.45 mmol), 0.13 ml 4% (w/w) OsO_4 in water and 1 ml water. The reaction mixture was stirred at room temperature for 2 h and then transferred into a separation funnel. Extracted with ether and the organic layer was combined, dried over magnesium sulfate and concentrated. The crude product was

purified by column chromatography on silica gel eluted with 10% ethyl acetate in petroleum ether to yield 30 mg 3-benzoyloctanal (67%) as colorless oil. $R_f = 0.45$ (30% ethyl acetate in petroleum ether); $[\alpha]_D^{25} = -51.3$ ($c = 1.9$, CH_2Cl_2 , 92% ee); HPLC (Chiralcel[®] OC column; 90:10 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 8.625$ min (major), $t_2 = 10.461$ min (minor)); IR (film): $\tilde{\nu}_{\text{max}} = 3063$ (w), 2931 (s), 2859 (s), 1723 (s), 1682 (s), 1597 (w), 1448 (m), 1385 (m), 1239 (m), 976 (m), 698 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 9.80$ (s, 1H), 7.98 (m, 2H), 7.58 (tt, $J_1 = 7.2$ Hz, $J_2 = 1.2$ Hz, 1H), 7.48 (m, 2H), 3.95 (m, 1H), 3.17 (dd, $J_1 = 18.6$ Hz, $J_2 = 8.8$ Hz, 1H), 2.66 (dd, $J_1 = 18.6$ Hz, $J_2 = 4.2$ Hz, 1H), 1.70 (m, 1H), 1.48 (m, 1H), 1.24 (m, 6H), 0.84 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 202.6$, 200.8, 136.4, 133.2, 128.7, 128.4, 45.4, 40.1, 32.3, 31.7, 26.8, 22.4, 14.0. Anal. Calcd. for $\text{C}_{15}\text{H}_{20}\text{O}_2$: C, 77.55; H, 8.68; Found: C, 77.40; H, 8.47.

2-Allyl-3-methyl-1-phenyl-1-butanone¹¹: Colorless oil. $R_f = 0.60$ (Diethyl ether/petroleum ether 1:9); IR (film): $\tilde{\nu}_{\text{max}} = 3078$ (w), 2963 (s), 2874 (m), 1682 (s), 1448 (m), 1209 (m), 1002 (m), 913 (m), 696 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.94$ (m, 2H), 7.55 (tt, $J_1 = 7.2$ Hz, $J_2 = 1.2$ Hz, 1H), 7.46 (m, 2H), 5.71 (m, 1H), 5.01 (m, 1H), 4.91 (m, 1H), 3.36 (m, 1H), 2.56 (m, 1H), 2.32 (m, 1H), 2.06 (m, 1H), 0.94 (t, $J = 7.0$ Hz, 6H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 203.9$, 138.3, 136.3, 132.9, 128.7, 128.3, 116.4, 52.3, 33.1, 30.5, 29.8, 21.3, 19.5.

To the solution of 36 mg 2-allyl-3-methyl-1-phenyl-1-butanone (0.18 mmol) in 1 ml ethyl acetate was added 85 mg sodium periodate (0.40 mmol), 0.12 ml 4% (w/w) OsO_4 in water and 1 ml water. The reaction mixture was stirred at room temperature for 2 h and then transferred into a separation funnel. Extracted with ether and the organic layer was combined, dried over magnesium sulfate and concentrated. The crude product was purified by column chromatography on silica gel eluted with 10% ethyl acetate in petroleum ether to yield 35 mg 3-benzoyl-4-methylpentanal (95%) as colorless oil. $R_f = 0.44$ (30% ethyl acetate in petroleum ether); $[\alpha]_D^{22} = -30.9$ ($c = 1.1$, CH_2Cl_2 , 32% ee); HPLC (Chiralcel[®] OC column; 90:10 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 10.658$ min (major), $t_2 = 13.146$ min (minor)); IR (film): $\tilde{\nu}_{\text{max}} = 3062$ (w), 2964 (s), 2875 (m), 2729 (m), 1716 (s), 1682 (s), 1597 (m), 1448 (m), 1392 (s), 1248 (s), 1200 (s), 978

(m), 696 cm⁻¹ (s); ¹H NMR (400 MHz, CDCl₃): δ = 9.82 (s, 1H), 7.98 (d, *J* = 7.6 Hz, 2H), 7.57 (t, *J* = 7.2 Hz, 1H), 7.48 (t, *J* = 7.3 Hz, 2H), 3.87 (ddd, *J*₁ = 10.2 Hz, *J*₂ = 4.8 Hz, *J*₃ = 3.2 Hz, 1H), 3.22 (dd, *J*₁ = 18.4 Hz, *J*₂ = 10.2 Hz, 1H), 2.61 (dd, *J*₁ = 18.4 Hz, *J*₂ = 3.0 Hz, 1H), 2.10 (m, 1H), 0.98 (d, *J* = 6.9 Hz, 3H), 0.81 (d, *J* = 6.9 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃): δ = 202.4, 201.1, 137.1, 133.0, 128.7, 128.4, 45.9, 41.5, 29.8, 21.3, 18.4. Anal. Calcd. for C₁₃H₁₆O₂: C, 76.44; H, 7.90; Found: C, 76.35; H, 7.90.

2-Benzyl-1-phenyl-4-penten-1-one: 75%; Colorless oil. *R*_f = 0.49 (Diethyl ether/petroleum ether 1:9); [α]_D²⁴ = -44.2 (c = 1.93, CHCl₃, 88% ee); HPLC (Chiralcel[®] AD column; 99:1 Heptane / Isopropanol; flow rate = 1 mL / min; *t*₁ = 7.873 min (minor), *t*₂ = 8.698 min (major)); IR (film): $\tilde{\nu}_{\text{max}}$ = 3063 (w), 2923 (w), 2874 (m), 1682 (s), 1448 (m), 1236 (m), 919 (m), 699 cm⁻¹ (s); ¹H NMR (400 MHz, CDCl₃): δ = 7.86 (m, 2H), 7.52 (tt, *J*₁ = 7.2 Hz, *J*₂ = 1.2 Hz, 1H), 7.41 (m, 2H), 7.3-7.1 (m, 5H), 5.73 (m, 1H), 5.01 (m, 2H), 3.80 (m, 1H), 3.10 (dd, *J*₁ = 13.7 Hz, *J*₂ = 7.6 Hz, 1H), 2.81 (dd, *J*₁ = 13.7 Hz, *J*₂ = 6.4 Hz, 1H), 2.53 (m, 1H), 2.30 (m, 1H); ¹³C NMR (100 MHz, CDCl₃): δ = 203.1, 139.7, 137.2, 135.3, 133.0, 129.1, 128.6, 128.46, 128.27, 126.3, 117.3, 48.1, 37.7, 36.3. Anal. Calcd. for C₁₈H₁₈O: C, 86.36; H, 7.25; Found: C, 86.50; H, 7.24.

1-(6'-Methoxy-2-naphthalenyl)-2-methyl-4-penten-1-one: 90%; Colorless oil. *R*_f = 0.25 (Diethyl ether/petroleum ether 1:9); [α]_D²³ = +7.4 (c = 1.54, CH₂Cl₂, 95% ee); HPLC (Chiralcel[®] OJ column; 90:10 Heptane / Isopropanol; flow rate = 1 mL / min; *t*₁ = 14.17 min (major), *t*₂ = 18.47 min (minor)); IR (film): $\tilde{\nu}_{\text{max}}$ = 3076 (w), 2974 (m), 2935 (m), 1674 (s), 1626 (s), 1482 (s), 1269 (s), 1196 (s), 1030 (m), 916 (m), 812 cm⁻¹ (m); ¹H NMR (400 MHz, CDCl₃): δ = 8.40 (d, *J* = 1.6 Hz, 1H), 8.01 (dd, *J*₁ = 8.4 Hz, *J*₂ = 1.6 Hz, 1H), 7.85 (d, *J* = 9.2 Hz, 1H), 7.77 (d, *J* = 8.8 Hz, 1H), 7.20 (dd, *J*₁ = 8.8 Hz, *J*₂ = 1.8 Hz, 1H), 7.15 (d, *J* = 1.8 Hz, 1H), 5.83 (m, 1H), 5.04 (m, 2H), 3.94 (s, 3H), 3.68 (sextet, *J* = 6.8 Hz, 1H), 2.62 (m, 1H), 2.25 (m, 1H), 1.26 (d, *J* = 7.6 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃): δ = 203.4, 159.8, 137.3, 136.0, 131.8, 131.2, 129.7, 127.9, 127.3, 125.1, 119.8, 116.8, 105.7, 55.5, 40.3, 37.9, 17.4. Anal. Calcd. for C₁₇H₁₈O₂: C, 80.28; H, 7.13; Found: C, 80.48; H, 6.99.

1-(2'-Fluorophenyl)-2-methyl-4-penten-1-one: 80%; Colorless oil. $R_f = 0.51$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{24} = +22$ ($c = 1.55$, CH_2Cl_2 , 94% ee); HPLC (Chiralcel[®] OB-H column; Heptane; flow rate = 1 mL / min; $t_1 = 9.53$ min (major), $t_2 = 13.96$ min (minor)); IR (film): $\tilde{\nu}_{\text{max}} = 3080$ (w), 2979 (w), 2936 (w), 1688 (s), 1610 (s), 1480 (m), 1452 (s), 1274 (m), 1211 (m), 976 (s), 977 (m), 756 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.77$ (td, $J_1 = 7.6$ Hz, $J_2 = 2.0$ Hz, 1H), 7.50 (m, 1H), 7.24 (m, 1H), 7.12 (ddd, $J_1 = 11.6$ Hz, $J_2 = 8.4$ Hz, $J_3 = 2.0$ Hz, 1H), 5.78 (m, 1H), 5.04 (m, 2H), 3.42 (m, 1H), 2.56 (m, 1H), 2.16 (m, 1H), 1.19 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 202.63$, 202.59, 162.5, 160.0, 135.8, 134.2, 134.1, 131.0, 130.9, 126.1, 126.0, 124.61, 124.58, 116.9, 116.8, 116.5, 45.23, 45.17, 37.3, 16.1. ^{19}F NMR (376 MHz, CDCl_3): $\delta = -111.6$. Anal. Calcd. for $\text{C}_{12}\text{H}_{13}\text{FO}$: C, 74.98; H, 6.82; F, 9.88; Found: C, 75.30; H, 6.62; F, 9.69.

1-(3'-Chlorophenyl)-2-methyl-4-penten-1-one: 97%; Colorless oil. $R_f = 0.65$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{23} = +32.7$ ($c = 2.2$, CH_2Cl_2 , 93% ee); HPLC (Chiralcel[®] OJ column; Heptane; flow rate = 1 mL / min; $t_1 = 8.87$ min (minor), $t_2 = 9.839$ min (major)); IR (film): $\tilde{\nu}_{\text{max}} = 3077$ (w), 2977 (w), 2934 (w), 1688 (s), 1571 (w), 1237 (w), 1205 (m), 993 cm^{-1} (w); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.92$ (t, $J = 2.0$ Hz, 1H), 7.82 (m, 1H), 7.53 (ddd, $J_1 = 8.0$ Hz, $J_2 = 2.4$ Hz, $J_3 = 1.2$ Hz, 1H), 7.42 (m, 1H), 5.77 (m, 1H), 5.04 (m, 2H), 3.48 (sextet, $J = 6.4$ Hz, 1H), 2.55 (m, 1H), 2.20 (m, 1H), 1.21 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 202.4$, 138.1, 135.5, 135.1, 132.9, 130.1, 128.5, 126.4, 117.1, 40.7, 37.5, 17.0. Anal. Calcd. for $\text{C}_{12}\text{H}_{13}\text{ClO}$: C, 69.07; H, 6.28; Cl, 16.99; Found: C, 68.96; H, 6.13; Cl, 17.16.

1-(4'-Bromophenyl)-2-methyl-4-penten-1-one: 94%; Colorless oil. $R_f = 0.56$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{26} = +21.8$ ($c = 1.5$, CH_2Cl_2 , 93% ee); HPLC (Chiralcel[®] OJ column; Heptane; flow rate = 1 mL / min; $t_1 = 12.35$ min (minor), $t_2 = 14.35$ min (major)); IR (film): $\tilde{\nu}_{\text{max}} = 3077$ (w), 2977 (w), 2934 (w), 1682 (s), 1585 (s), 1397 (m), 1206 (m), 1071 (m), 976 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 7.82$ (m, 2H), 7.62 (m, 2H), 5.76 (m, 1H), 5.04 (m, 2H), 3.47 (sextet, $J = 6.8$ Hz, 1H), 2.54 (m, 1H), 2.20 (m, 1H), 1.20 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 202.7$, 135.6, 135.2,

132.0, 129.9, 128.2, 117.0, 40.5, 37.6, 17.0. Anal. Calcd. for $C_{12}H_{13}BrO$: C, 56.94; H, 5.18; Br, 31.57; Found: C, 57.08; H, 5.08; Br, 31.80.

1-(2-Methoxyphenyl)-2-methyl-4-penten-1-one: 99%; Colorless oil. $R_f = 0.32$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{23} = -0.33$ ($c = 5.0$, CH_2Cl_2 , 98% ee); IR (film): $\tilde{\nu}_{max} = 3076$ (m), 2976 (m), 2921 (m), 1674 (s), 1597 (s), 1485 (s), 1464 (s), 1285 (s), 1245 (s), 1024 (m), 976 (m), 755 cm^{-1} (m); 1H NMR (400 MHz, $CDCl_3$): $\delta = 7.23$ (dd, $J_1 = 7.6$ Hz, $J_2 = 2.0$ Hz, 1H), 7.43 (m, 1H), 6.97 (m, 2H), 5.77 (m, 1H), 5.00 (m, 2H), 3.88 (s, 3H), 3.50 (m, 1H), 2.52 (m, 1H), 1.12 (m, 1H), 1.14 (d, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$): $\delta = 207.1$, 157.9, 136.4, 132.9, 130.2, 129.0, 120.8, 116.4, 111.4, 55.5, 45.1, 37.5, 16.0. Anal. Calcd. for $C_{13}H_{16}O_2$: C, 76.44; H, 7.90; Found: C, 76.46; H, 8.01.

To the solution of 50 mg 1-(2-methoxyphenyl)-2-methyl-4-penten-1-one (0.24 mmol) in 1 ml ethyl acetate was added 113 mg sodium periodate (0.53 mmol), 0.15 ml 4% (w/w) OsO_4 in water and 1 ml water. The reaction mixture was stirred at room temperature for 2 h and then transferred into a separation funnel. Extracted with ether and the organic layer was combined, dried over magnesium sulfate and concentrated. The crude product was purified by column chromatography on silica gel eluted with 10% ethyl acetate in petroleum ether to yield 21 mg 3-benzoyl-4-methylpentanal (42%) as colorless oil. $R_f = 0.31$ (30% ethyl acetate in petroleum ether); $[\alpha]_D^{23} = +38.3$ ($c = 2.1$, CH_2Cl_2 , 98% ee); HPLC (Chiralcel[®] OC column; 90:10 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 22.601$ min (major), $t_2 = 29.465$ min (minor)); IR (film): $\tilde{\nu}_{max} = 2972$ (m), 2934 (m), 1726 (s), 1667 (s), 1598 (s), 1485 (s), 1463 (s), 1285 (s), 1245 (s), 1022 (m), 979 (m), 756 cm^{-1} (s); 1H NMR (400 MHz, $CDCl_3$): $\delta = 9.82$ (t, $J = 1.2$ Hz, 1H), 7.65 (dd, $J_1 = 7.6$ Hz, $J_2 = 2.0$ Hz, 1H), 7.47 (ddd, $J_1 = 10$ Hz, $J_2 = 7.6$ Hz, $J_3 = 2.0$ Hz, 1H), 7.00 (m, 2H), 3.98 (sextet, $J = 7.2$ Hz, 1H), 3.90 (s, 3H), 3.01 (ddd, $J_1 = 17.2$ Hz, $J_2 = 10.8$ Hz, $J_3 = 1.2$ Hz, 1H), 2.50 (ddd, $J_1 = 17.2$ Hz, $J_2 = 6.4$ Hz, $J_3 = 1.6$ Hz, 1H), 1.19 (d, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$): $\delta = 204.8$, 201.2, 158.0, 133.5, 130.7, 127.5, 120.9, 111.5, 55.6, 46.9, 40.3, 16.9. Anal. Calcd. for $C_{12}H_{14}O_3$: C, 69.88; H, 6.84; Found: C, 69.68; H, 6.69.

2-Methyl-1-(2-pyridinyl)-4-penten-1-one: 95%; Colorless oil. $R_f = 0.33$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{24} = +26.3$ ($c = 2.4$, CH_2Cl_2 , 73% ee); IR (film): $\tilde{\nu}_{\text{max}} = 3077$ (m), 2976 (s), 2934 (m), 1694 (s), 1583 (m), 1455 (m), 1436 (m), 1355 (m), 1266 (m), 1216 (m), 996 (s), 981 (s), 746 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 8.69$ (m, 1H), 8.04 (m, 1H), 7.84 (ddd, $J_1 = 7.6\text{ Hz}$, $J_2 = 7.6\text{ Hz}$, $J_3 = 1.8\text{ Hz}$, 1H), 7.47 (m, 1H), 5.80 (m, 1H), 5.00 (m, 2H), 4.18 (sextet, $J = 7.2\text{ Hz}$, 1 H), 2.56 (m, 1H), 2.23 (m, 1H), 1.20 (d, $J = 6.8\text{ Hz}$, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 204.9$, 153.0, 149.0, 137.0, 136.1, 127.0, 122.5, 116.5, 38.9, 37.2, 16.5. Anal. Calcd. for $\text{C}_{11}\text{H}_{13}\text{NO}$: C, 75.40; H, 7.48; N, 7.99; Found: C, 75.21; H, 7.45; N, 7.76.

To the solution of 23 mg 2-methyl-1-(2-pyridinyl)-4-penten-1-one (0.13 mmol) in 1 ml ethyl acetate was added 61mg sodium periodate (0.28 mmol), 0.085 ml 4% (w/w) OsO_4 in water and 1 ml water. The reaction mixture was stirred at room temperature for 2 h and then transferred into a separation funnel. Extracted with ether and the organic layer was combined, dried over magnesium sulfate and concentrated. The crude product was purified by column chromatography on silica gel eluted with 10% ethyl acetate in petroleum ether to yield 19 mg 3-methyl-4-oxo-4-(pyridin-2-yl)butanal (87%) as colorless oil. $R_f = 0.28$ (30% ethyl acetate in petroleum ether); $[\alpha]_D^{23} = +1.6$ ($c = 0.7$, CH_2Cl_2 , 73% ee); HPLC (Chiralcel[®] OC column; 90:10 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 18.79$ min (major), $t_2 = 22.55$ min (minor)); IR (film): $\tilde{\nu}_{\text{max}} = 3056$ (w), 2972 (m), 2934 (m), 2829 (m), 2727 (m), 1716 (s), 1698 (s), 1583 (s), 1456 (s), 1378 (s), 1267 (m), 1233 (s), 996 (s), 747 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 9.80$ (s, 1H), 8.71 (dq, $J_1 = 4.8\text{ Hz}$, $J_2 = 0.9\text{ Hz}$, 1H), 8.05 (m, $J_1 = 8.0\text{ Hz}$, $J_2 = 0.9\text{ Hz}$, 1H), 7.85 (td, $J_1 = 7.6\text{ Hz}$, $J_2 = 1.8\text{ Hz}$, 1H), 7.48 (ddd, $J_1 = 7.5\text{ Hz}$, $J_2 = 4.8\text{ Hz}$, $J_3 = 1.2\text{ Hz}$, 1H), 4.54 (m, 1H), 3.12 (ddd, $J_1 = 18.2\text{ Hz}$, $J_2 = 8.4\text{ Hz}$, $J_3 = 0.8\text{ Hz}$, 1H), 2.63 (ddd, $J_1 = 18.2\text{ Hz}$, $J_2 = 5.4\text{ Hz}$, $J_3 = 1.2\text{ Hz}$, 1H), 1.27 (d, $J = 8.2\text{ Hz}$, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 203.4$, 200.8, 152.3, 149.1, 137.1, 127.6, 122.7, 47.2, 34.3, 17.4. Anal. Calcd. for $\text{C}_{10}\text{H}_{11}\text{NO}_2$: C, 67.78; H, 6.26; N, 7.90; Found: C, 68.00; H, 6.50; N, 8.06.

2-Methyl-1-(3'-nitrophenyl)-4-penten-1-one: 91%; Colorless oil. $R_f = 0.31$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{24} = +27.2$ ($c = 1.3$, CHCl_3 , 82% ee); HPLC (Chiralcel[®] AD column; 99:1 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 10.65$ min (minor), $t_2 = 12.69$ min (major)); IR (film): $\tilde{\nu}_{\text{max}} = 3082$ (w), 2978 (w), 2935 (w), 1694 (s), 1614

(m), 1532 (s), 1350 (s), 1208 (m), 1001 (m), 919 (m), 716 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): δ = 8.28 (t, J = 2.0 Hz, 1H), 8.13 (ddd, J_1 = 8.2 Hz, J_2 = 2.3 Hz, J_3 = 1.0 Hz, 1H), 7.74 (ddd, J_1 = 7.9 Hz, J_2 = 1.8 Hz, J_3 = 1.0 Hz, 1H), 7.52 (t, J = 8.2 Hz, 1H), 6.05 (q, J = 7.0 Hz, 1H), 5.96 (m, 1H), 5.42 (dq, J_1 = 17.2 Hz, J_2 = 1.4 Hz, 1H), 5.33 (dq, J_1 = 10.4 Hz, J_2 = 1.2 Hz, 1H), 4.72 (dt, J_1 = 5.8 Hz, J_2 = 1.4 Hz, 1H), 1.83 (d, J = 7.2, 1H); ^{13}C NMR (100 MHz, CDCl_3): δ = . Anal. Calcd. for $\text{C}_{12}\text{H}_{13}\text{NO}_3$: C, 65.74; H, 5.98; N, 6.39;; Found: C, 66.00; H, 5.86; N, 6.45.

1-(2-Furanyl)-2-methyl-4-penten-1-one: 89%; Colorless oil. R_f = 0.29 (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{24}$ = +30.2 (c = 2.64, CH_2Cl_2 , 88% ee); GC (Cyclosil B[®] column; 90 °C isothermal; 1.2 flow rate; t_1 = 47.567 min (minor), t_2 = 48.202 min (major)); IR (film): $\tilde{\nu}_{\text{max}}$ = 3134 (w), 3079 (w), 2976 (m), 2934 (m), 1672 (s), 1567 (s), 1467 (s), 1395 (m), 1262 (m), 1015 (m), 983 (m), 761 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): δ = 7.57 (dd, J_1 = 1.7 Hz, J_2 = 0.8 Hz, 1H), 7.18 (dd, J_1 = 3.5 Hz, J_2 = 0.8 Hz, 1H), 6.51 (dd, J_1 = 3.5 Hz, J_2 = 1.7 Hz, 1H), 5.74 (m, 1H), 5.00 (m, 2H), 3.30 (sextet, J = 7.0 Hz, 1 H), 2.52 (m, 1H), 2.18 (m, 1H), 1.18 (d, J = 6.8 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ = 192.7, 152.4, 146.4, 135.7, 117.4, 116.9, 112.2, 41.3, 37.4, 16.7. Anal. Calcd. for $\text{C}_{10}\text{H}_{12}\text{O}_2$: C, 73.15; H, 7.37; Found: C, 72.96; H, 7.37.

2-Methyl-1-(2'-(trifluoromethyl)phenyl)-4-penten-1-one: 94%; Colorless oil. R_f = 0.37 (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{22}$ = +11.1 (c = 2.40, CHCl_3 , 92% ee); HPLC (Chiralcel[®] OD-H column; 2000:1 Heptane / Isopropanol; flow rate = 1 mL / min; t_1 = 22.792 min (minor), t_2 = 24.434 min (major)); IR (film): $\tilde{\nu}_{\text{max}}$ = 3080 (w), 3079 (w), 2980 (m), 2937 (w), 1705 (s), 1458 (m), 1314 (s), 1274 (m), 1170 (s), 1135 (s), 1063 (m), 975 (m), 769 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): δ = 7.72 (dd, J_1 = 7.6 Hz, J_2 = 0.5 Hz, 1H), 7.58 (m, 1H), 7.43 (dd, J_1 = 7.2 Hz, J_2 = 1.0 Hz, 1H), 5.76 (m, 1H), 5.08 (m, 2H), 3.16 (sextet, J = 7.0 Hz, 1 H), 2.54 (m, 1H), 2.15 (m, 1H), 1.16 (d, J = 7.0 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ = 207.2, 135.7, 131.9, 130.3, 127.6, 127.23, 127.17, 127.13, 122.4, 117.5, 45.8, 36.9, 16.0. ^{19}F NMR (376 MHz, CDCl_3): δ = -58.3. Anal. Calcd. for $\text{C}_{13}\text{H}_{13}\text{F}_3\text{O}$: C, 64.46; H, 5.41; F, 23.53; Found: C, 64.52; H, 5.26; F, 23.68.

1-Mesityl-2-methyl-4-penten-1-one: 99%; Colorless oil. $R_f = 0.47$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{23} = +0.5$ ($c = 2.47$, CHCl_3 , 96% ee); IR (film): $\tilde{\nu}_{\text{max}} = 2969$ (s), 2921 (s), 1694 (s), 1611 (m), 1456 (m), 1378 (w), 973 cm^{-1} (m); ^1H NMR (400 MHz, CDCl_3): $\delta = 6.82$ (s, 2H), 5.76 (m, 1H), 5.05 (m, 2H), 2.92 (m, 1H), 2.54 (m, 1H), 2.26 (s, 3H), 2.18 (s, 6H), 2.10 (m, 1H), 1.12 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 213.1$, 138.8, 138.5, 136.0, 133.5, 128.8, 117.1, 47.3, 36.3, 21.1, 19.8, 15.2. Anal. Calcd. for $\text{C}_{15}\text{H}_{20}\text{O}$: C, 83.28; H, 9.32; Found: C, 83.56; H, 9.07.

To the solution of 33 mg 1-mesityl-2-methyl-4-penten-1-one (0.15 mmol) in 1 ml ethyl acetate was added 71 mg sodium periodate (0.33 mmol), 0.1 ml 4% (w/w) OsO_4 in water and 1 ml water. The reaction mixture was stirred at room temperature for 2 h and then transferred into a separation funnel. Extracted with ether and the organic layer was combined, dried over magnesium sulfate and concentrated. The crude product was purified by column chromatography on silica gel eluted with 10% ethyl acetate in petroleum ether to yield 24 mg 3-methyl-4-oxo-4-(pyridin-2-yl)butanal (73%) as colorless oil. $R_f = 0.58$ (30% ethyl acetate in petroleum ether); $[\alpha]_D^{24} = +24$ ($c = 24$, CH_2Cl_2 , 96% ee); HPLC (Chiralcel[®] OJ column; 99:1 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 21.43$ min (minor), $t_2 = 26.84$ min (major)); IR (film): $\tilde{\nu}_{\text{max}} = 2973$ (s), 2920 (s), 2729 (m), 1732 (s), 1694 (s), 1611 (s), 1456 (m), 1381 (m), 1234 (m), 975 (s), 853 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 9.86$ (s, 1H), 6.95 (s, 1H), 3.50 (sextet, $J = 7.2$ Hz, 1H), 3.02 (ddd, $J_1 = 17.7$ Hz, $J_2 = 6.4$ Hz, $J_3 = 0.8$ Hz, 1H), 2.51 (ddd, $J_1 = 17.7$ Hz, $J_2 = 6.4$ Hz, $J_3 = 1.2$ Hz, 1H), 2.28 (s, 3H), 2.22 (s, 6H), 1.17 (d, $J = 7.3$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 211.4$, 200.5, 138.8, 137.5, 133.6, 128.9, 45.5, 42.5, 21.1, 19.8, 16.1. Anal. Calcd. for $\text{C}_{14}\text{H}_{18}\text{O}_2$: C, 77.03; H, 8.31; Found: C, 76.89; H, 8.17.

1-Cyclohexenyl-2-methyl-4-penten-1-one: 94%; Colorless oil. $R_f = 0.57$ (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{22} = +29.8$ ($c = 3.1$, CH_2Cl_2 , 88% ee); HPLC (Chiralcel[®] OB-H column; 97:3 Heptane / Isopropanol; flow rate = 1 mL / min; $t_1 = 4.710$ min (major), $t_2 = 5.709$ min (minor)); IR (film): $\tilde{\nu}_{\text{max}} = 3076$ (w), 2973 (s), 2934 (s), 2861 (s), 1665 (s), 1637 (s), 1450 (m), 1435 (m), 1387 (w), 1197 (s), 988 (s), 912 cm^{-1} (s); ^1H NMR (400 MHz, CDCl_3): $\delta = 6.90$ (m, 1H), 5.72 (m, 1H), 5.00 (m, 2H), 3.24 (sextet, $J = 6.9$ Hz, 1H), 2.40 (m, 1H), 2.24 (m, 4H), 2.07 (m, 1H), 1.62 (m, 4H), 1.07 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3): $\delta = 204.8$, 139.4, 138.6, 136.3, 116.3, 38.8, 38.1, 26.2,

23.4, 22.1, 21.6, 17.6. Anal. Calcd. for C₁₂H₁₈O: C, 80.85; H, 10.18; Found: C, 81.08; H, 10.40.

(*E*)-4-Methyl-1-phenyl-1,6-heptadien-3-one: 93%; Colorless oil. R_f = 0.37 (Diethyl ether/petroleum ether 1:9); $[\alpha]_D^{23}$ = -20.0 (c = 3.3, CH₂Cl₂, 91% ee); HPLC (Chiralcel[®] OD-H column; 99.5:0.5 Heptane / Isopropanol; flow rate = 1 mL / min; t_1 = 18.680 min (minor), t_2 = 21.179 min (major)); IR (film): $\tilde{\nu}_{\max}$ = 3079 (m), 2974 (s), 2932 (m), 1688 (s), 1661 (s), 1613 (s), 1577 (s), 1450 (s), 1187 (s), 1054 (s), 990 (s), 916 (s), 762 cm⁻¹ (s); ¹H NMR (400 MHz, CDCl₃): δ = 7.61 (d, J = 16.0 Hz, 1H), 7.56 (m, 2H), 7.39 (m, 3H), 6.81 (d, J = 16.0 Hz, 1H), 5.78 (m, 1H), 5.05 (m, 2H), 2.93 (sextet, J = 6.9 Hz, 1H), 2.51 (m, 1H), 2.18 (m, 1H), 2.07 (m, 1H), 1.18 (d, J = 7.0 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃): δ = 203.0, 142.7, 135.8, 134.7, 130.5, 129.0, 128.4, 124.9, 116.8, 44.4, 37.3, 16.4. Anal. Calcd. for C₁₄H₁₆O: C, 83.96; H, 8.05; Found: C, 83.71; H, 8.25.

Reference:

- (1) Perrin, D. D. Armarego, W. L. F. *Purification of Laboratory Chemicals*; Pergamon Press: New York, 1988.
- (2) Ukai, T.; Kawazura, H.; Ishii, Y.; Bonnet, J. J.; Ibers, J. A. *J. Organometallic Chem.* **1974**, 65, 253-266.
- (3) Trost, B. M.; Van Vranken, D. L.; Bingel, C. *J. Am. Chem. Soc.* **1992**, 114, 9327-9343.
- (4) Trost, B. M.; Bunt, R. C.; Lemoine, R. C.; Calkins, T. L. *J. Am. Chem. Soc.* **2000**, 122, 5968-5976.
- (5) Harwood, L. M.; Houminer, Y.; Manage, A.; Seeman, J. I. *Tetrahedron* **1994**, 35, 8027-8030.
- (6) Crotti, P.; Di Bussolo, V.; Favero, L.; Macchia, F.; Pineschl, M.; Napolitano, E. *Tetrahedron* **1999**, 55, 5853-5866.
- (7) Henin, F.; M'Boungou-M'Passi, A.; Muzart, J.; Pete, J. P. *Tetrahedron* 1994, 50, 2849-2864.
- (8) Baldwin, J. E.; Patrick, J. E. *J. Am. Chem. Soc.* **1971**, 93, 3556.
- (9) Wagner, P. J.; Liu, K. C.; Noguchi, Y. *J. Am. Chem. Soc.* **1981**, 103, 3837-3841.
- (10) Gansaeuer, A.; Fielenbach, D.; Stock, C.; Geich-Gimbel, D. *Adv. Synth. Catal.* **2003**, 345, 1017-1030.
- (11) Barentsen, Helma M.; Sieval, Alex B.; Cornelisse, Jan. *Tetrahedron* **1995**, 51, 7495-520.