The Palladium-catalyzed Conversion of β,γ - Unsaturated Silyl sulfinates into (*E*)-Alkenes: Asymmetric Synthesis of Polypropionate Fragments

Xiaogen Huang, Cotinica Craita and Pierre Vogel*

*Laboratory of Glycochemistry and Asymmetric Synthesis, Swiss Federal Institute of
Technology, EPFL-BCH, CH-1015 Lausanne-Dorigny, Switzerland*

pierre.vogel@epfl.ch

Fax +41 21 693 9375

Phone +41 21 693 9371

General remarks: Commercial reagents (Fluka, Aldrich) were used without purification. Solvents were distilled prior to use: THF from Na and benzophenone; MeOH from Mg and I₂; DMF and CH₂Cl₂ from CaH₂. Liquid/solid flash chromatography (FC): columns of silica gel (0.040-0.63 mm, Merck No.9385 silica gel 60, 240-400 mesh). TLC for reaction monitoring: Merck silica gel 60 F₂₅₄ plates; detection by UV light; Pancaldi reagent or KMnO₄. IR spectra: Perkin-Elmer-1420 spectrometer. ¹H-NMR spectra : Bruker-ARX-400 spectrometer (400 MHz); δ (H) in ppm relative to the solvent's residual ¹H signal [CHCl₃, δ (H) 7.27; CH₃OD, δ (H) 3.31; C₆H₆, δ (H) 7.3] as internal reference; all ¹H assignments were confirmed by 2D-

COSY spectra. ^{13}C NMR spectra: same instrument as above (100.6 MHz); $\delta(\text{C})$ in ppm relative to solvent's C-signal [CDCl_3 , $\delta(\text{C})$ 77.0; CD_3OD , $\delta(\text{C})$ 49.2; C_6D_6 $\delta(\text{C})$ 128.5] as internal reference; coupling constants J in Hz. MS: Nermag R-10-10C, chemical ionization (NH_3) mode m/z (amu) [% relative base peak (100%)], HRMS: Jeol AX-505. Elemental analyses: Ilse Beetz, D-96301 Kronach, Germany.

1. NMR study of the reaction of trimethylsilyl 2-methylprop-2-enesulfinate (**19**) with palladium complexes.

1.1. Reaction of **19** with 0.5 equiv. 1:1 $\text{Pd}(\text{OAc})_2/\text{PPh}_3$

To CDCl_3 (0.5ml) in a 5mm NMR tube was added **19** (68.4mg, 0.35mmol, 1equiv.), $\text{Pd}(\text{OAc})_2$ (40mg, 0.175mmol, 0.5equiv.) and PPh_3 (46mg, 0.175mmol, 0.5equiv.). The reaction was followed by ^1H -NMR at 25°C. The intermediate complex **20** was formed first. It was then converted slowly and quantitatively into sulfone **21** at 25°C (see ^1H NMR spectra below: **Fig. 1**).

*Data for intermediate **20** (2'-methylallyl triphenylphosphine palladium 2-methylprop-2-enesulfinate):*

^1H -NMR (CDCl_3 , 400MHz): 7.8~6.8(m, H arom.), 4.90(m, H-C(3)), 4.66(m, H-C(3)), 4.36(t, H-C(1')), 3.53(d, H-C(1')), 3.34(m, H-C(1)), 3.00(s, H-C(3')), 2.87(s, H-C(3')), 1.90(m, $\text{CH}_3\text{-C}(2')$), 1.74(m, $\text{CH}_3\text{-C}(2)$).

*Data for compound **21** (bis(methallyl)sulfone):*

MS (CI) m/z : 192(100, $[\text{M}+18]^+$), 175(3), 109(6), 95(13).

^1H -NMR (CDCl_3 , 400MHz): 5.19(m, H-C(3)), 5.03(m, H-C(3)), 3.65(s, H-C(1)), 1.96(s, $\text{CH}_3\text{-C}(2)$).

Figure 1: ^1H -NMR spectra taken during reaction **19**+1:1 $\text{Pd}(\text{OAc})_2/\text{PPh}_3$ (0.5equiv.).

1.2. Reaction of **19** with one equivalent of 1:1 $\text{Pd}(\text{OAc})_2/\text{PPh}_3$

To CDCl_3 (0.5ml) in a 5mm NMR tube was added **19** (20mg, 0.11mmol, 1equiv.), $\text{Pd}(\text{OAc})_2$ (25mg, 0.11mmol, 1equiv.) and PPh_3 (29mg, 0.11mmol, 1equiv.) at room temperature. The reaction was followed by ^1H -NMR at 25°C . The intermediate **22** was formed first. It was then converted into the stable complex bis (methallyl) (triphenylphosphine) palladium complex **23** (see **Fig. 2**).

Data for **23**:

^1H -NMR (CDCl_3 , 400MHz): 7.8~6.8(m, H arom.), 4.53(m, H-C(3)), 3.67(d, H-C(1)), 2.10(s, CH_3 -C(2)).

Figure 2: ^1H -NMR spectra taken during the reaction of **19**+1equiv. of 1:1 $\text{Pd}(\text{OAc})_2/\text{PPh}_3$.

2. Kinetic studies

2.1. Desulfitation of prop-2-enesulfinic acid (**26**, **26-d**), kinetic deuterium isotopic effect in CDCl_3 .

One standard solution was prepared. Trimethylsilyl prop-2-enesulfinate **26** (460mg, 2.6mmol) and toluene (22mg, internal standard) were dissolved into 5ml of CDCl_3 . Two parallel reaction flasks containing 1ml of the above solution were prepared and to each of them 13mg of $\text{Pd}(\text{OAc})_2$ and 14mg of PPh_3 were added. The reaction flasks were cooled down to -40°C and 0.25ml of anhydrous *i*-PrOH or *i*-PrOD was introduced. The resulted mixtures were stirred at the same temperature for half an hour, allowing complete desilylation of the silyl sulfinate. The solutions so-obtained were transferred into 5mm NMR tubes. The reactions were followed at 25°C (2 min's interval, one single scan). The results are shown in **Fig. 3**. It was found that the reactions follow first-order rate laws. A kinetic deuterium isotopic effect $k_{\text{H}}/k_{\text{D}}=2.2\pm0.2$ was evaluated at 25°C .

Figure 3: Kinetics of the reactions of prop-2-enesulfinic acid: (A) in 1/4 *i*-PrOH/CDCl₃, (B) in 1/4 *i*-PrOD/CDCl₃ in the presence of 9% of 1:1 Pd(OAc)₂/PPh₃.

2.2. Solvent effect on the rate and kinetic deuterium isotope effect of desulfitation of prop-2-enesulfinic acid in the presence or absence of Pd(OAc)₂/PPh₃.

Instead of using CDCl₃ as solvent in the preparation of trimethylsilyl prop-2-enesulfinate, CD₃CN was used. The same method of sample preparation as above was applied and the kinetics followed by ¹H-NMR (**Fig. 4**). It was found that at 25°C the desulfitation in 1/4 *i*-PrOH/CD₃CN is about twice as fast as in 1/4 *i*-PrOH/CDCl₃. A significant increase of the kinetic deuterium isotopic effect was found as k_H/k_D in CD₃CN was evaluated to 6.9±0.2 at 25°C.

Figure 4. Kinetics of the desulfonation of prop-2-enesulfonic acid in CD_3CN in the presence of 9% $\text{Pd}(\text{OAc})_2/\text{PPh}_3$, at 25°C

Trimethylsilyl prop-2-enesulfinate (230mg, 1.3mmol) and toluene (12mg) were dissolved into 2.5ml of CD_3CN . Two parallel reaction flasks containing 1ml of the above solution were prepared and to each of them 0.25ml of anhydrous $i\text{-PrOH}/i\text{-PrOD}$ at -40°C . The resulting mixtures were stirred at the same temperature for 15 mins, allowing complete desilylation of the silyl sulfinate. The solutions were transferred into 5mm NMR tubes quickly for kinetic study (**Fig. 5**). The kinetic deuterium isotopic effect was calculated to be $k_{\text{H}}/k_{\text{D}}=5.9 \pm 0.2$.

Figure 5: Kinetics of the desulfonation of prop-2-enesulfonic acid in CD_3CN in the absence of $\text{Pd}(\text{OAc})_2/\text{PPh}_3$, at 25°C .

2.3. Role of the palladium catalyst on the rate of retro-ene desulfitation.

Silyl sulfinate **15** was prepared by the oxyallylation reaction of diene **13** and enoxysilane **14**. After 12h at -80°C , one equivalent of Et_3N to Lewis acid was introduced. SO_2 and CH_2Cl_2 were then removed completely under vacuum at -78°C . The resulting oil was dissolved into CD_3CN and transferred into a 5 mm NMR tube. The spectrum showed the nearly quantitative formation of **15**. A large excess of isopropanol was then added and the reaction was followed by ^1H -NMR. Desilylation occurred at -20°C , giving the corresponding sulfinic acid **16** which was relatively stable as the retro-ene reaction at room temperature was rather slow. Heating to 50°C led to fast decomposition of **16** and the formation of alkene **17**. This ^1H -NMR study showed that the rate of retro-ene reaction is very sensitive to the structure of the substrate (compare rate constants of the reactions of prop-2-enesulfonic acid and 2-methylprop-2-enesulfonic acid), but it is not affected significantly by the palladium catalyst.

A standard solution of silyl sulfinate **15** in CD_3CN was prepared as above. Two parallel reaction flasks were prepared. To each of them was added the same volume of isopropanol but to one of them was added a catalytic amount of $\text{Pd}(\text{OAc})_2$ and PPh_3 (1/1, 10%). The reactions of both solutions were followed by ^1H -NMR at 40°C . The results are shown in **Fig. 6**. It was found that the reactions were first-order and in the presence of palladium species the reaction was only 1.3 times faster than the blank. Interestingly, however, the presence of palladium species led to a cleaner reaction mixture.

Figure 6: Kinetics of the desulfitation of **16** (A) in the absence of palladium catalyst, (B) in the presence of 0.1 equiv. of 1:1 $\text{Pd}(\text{OAc})_2/\text{PPh}_3$.

3.1. General procedure for the preparations of (±)-**17**, (±)-**28/29**, (-)-**31/32**, (-)-**34/35**, (-)-**37**, (+)-**39/40**, (+)-**41/42** and **44/45**

A two-necked flask was dried by flame and filled with Ar, then 20 mL of CH₂Cl₂ and (CF₃SO₂)₂NH (1mL, ~0.5M in CH₂Cl₂ solution, 0.5mmol, 0.19equiv.) were introduced. The mixture was cooled by liquid N₂ and connected to vacuum. About 10 mL of SO₂ was condensed into the flask, which was allowed to warm to -80 °C by using acetone-dry ice cooling system. After 30 min., the mixture of diene (**13**, (-)-**30**, (+)-**38**: 2.65mmol, 1equiv.) and enoxysilane (2.65mmol, 1equiv.) in 5 mL CH₂Cl₂ was introduced dropwise. Stirring was continued overnight at -84 °C. Then all the SO₂ and CH₂Cl₂ were evaporated under vacuum. The resulting viscous mixture was dissolved into anh. CH₃CN. In another flask, a mixture of Pd(OAc)₂ (60mg, 0.26mmol, 0.1equiv.), PPh₃ (70mg, 0.26mmol, 0.1equiv.) and anh. K₂CO₃ (70mg, 1 equiv. with respect to the acid promoter) was prepared. The solution of the resulting mixture in CH₃CN (16mL) was added, followed by the addition of isopropanol (4mL). The resulting mixture was heated to 80 °C for 30 min. Then sat. aq. NaHCO₃ solution was added and the mixture was extracted by ether. The organic phase was separated and washed with brine and dried (Na₂SO₄). After solvent evaporation, flash column chromatography on silica gel gave the corresponding product.

3.2. Complete characterisation of new products.

(4*RS*,5*SR*,6*E*)-4-Benzyloxy-5-methyloct-6-en-2-one (±**17**):¹

Colorless oil (72%).

¹H NMR (CDCl₃, 400MHz): 7.35-7.24 (*m*, 5 arom H), 5.47 (*dqd*, ³*J*(H-C(7), H-C(6))=15.3, ³*J*(H-C(7), H-C(8))=6.0, ³*J*(H-C(7), H-C(5))=0.8, H-C(7)), 5.38 (*ddq*, ³*J*(H-C(6), H-C(7))=15.3, ³*J*(H-C(6), H-C(5))=7.1, ³*J*(H-C(6), H-C(8))=1.3, H-C(6)), 4.53 (*m*, 2H,

CH₂(Bn)), 3.91 (*ddd*, ³*J* (H-C(4), H-C(3))=4.0, ³*J* (H-C(4), H-C(3))=8.5, ³*J* (H-C(4), H-C(5))=3.9, H-C(4)), 2.67 (*dd*, ²*J*=16.2, ³*J* (H-C(3), H-C(4))=8.5, H-C(3)), 2.47-2.50 (*m*, H-C(5)); 2.44 (*dd*, ³*J* (H-C(3), H-C(4))=4.0, ²*J*=16.2, H-C(3)), 2.14(*s*, H-C(1)), 68 (*ddd*, ³*J* (H-C(8), H-C(7))=6.0, ⁴*J* (H-C(8), H-C(6))=1.3, ⁵*J* (H-C(8), H-C(5))=0.9, CH₃(8)), 1.03 (*d*, ³*J* (CH₃-C(5), H-C(6))=6.9, CH₃-C(5)).

45:55 Mixture of (4*RS*,5*SR*,6*SR*,7*E*)- and (4*RS*,5*RS*,6*RS*,7*E*)-5-benzyloxy-4,6-Dimethylnon-7-en-3-one ((±)-28 and (±)-29

Colorless oil (64%); **R_f**=0.35 (PE/EA=10/1); **UV** (MeCN): 195 (1290); **IR** (film): 2970, 2934, 1715, 1455, 1376, 1069, 972, 698; **CI-MS** (NH₃): 275 (2, [M+1]⁺), 251 (0.4), 207 (7), 189 (3), 151 (2), 131 (3.5); **¹H-NMR**(CDCl₃, 400MHz) of the mixture: 7.37-7.23 (*m*, 5H arom.); 5.57-5.34 (*m*, H-C((7)+ H-C(8))); 4.46 (*m*, CH₂(Bn)); 4.58 (*m*, CH₂(Bn)); 3.62 (*dd*, ³*J*(H-C(5), H-C(4))=7.5, ³*J*(H-C(5), H-C(6))=4.0, H-C(5)); 3.57 (*dd*, ³*J*(H-C(5), H-C(4))=9.6, ³*J*(H-C(5), H-C(6))=2.4, H-C(5)); 2.83 (*dq*, ³*J*(H-C(4), CH₃-C(4))=7.0, ³*J*(H-C(4), H-C(5))=9.6, H-C(4)); 2.5 (*q*, ³*J*(H-C(2), H-C(1))=7.3, H-C(2)); 2.39 (*m*, H-C(6)); 2.28 (*m*, H-C(6)); 1.7 (*d*, ³*J*(CH₃-C(9), H-C(8))=5.6, CH₃-C(9)); 1.67 (*d*, ³*J*(CH₃-C(9), H-C(8))=5.6, CH₃-C(9)); 1.18 (*d*, ³*J*(CH₃-C(4), H-C(4))=7.0, CH₃-C(4)); 1.14 (*d*, ³*J*(CH₃-C(6), H-C(6))=7.0, CH₃-C(6)); 1.07 (*d*, ³*J*(CH₃-C(6), H-C(6))=6.9, CH₃-C(6)); 1.03 (*t*, ³*J*(CH₃-C(1), H-C(2))=7.1, CH₃-C(1)); 1.0 (*t*, ³*J*(CH₃-C(1), H-C(2))=7.2, CH₃-C(1)); 0.96 (*d*, ³*J*(CH₃-C(4), H-C(4))=7.0, CH₃-C(4)); **¹³C-NMR**(CDCl₃, 100.61MHz) of the mixture: 215.4 (*s*, C(3)); 214.4(*s*, C(3)); 138.7, 138.5 (2*s*, 2C arom.); 132.8, 131.9 (2*d*, ¹*J*(C, H)=151.8, C(7) or C(8)); 128.3, 128.2, 127.7, 127.6 (4*d*, ¹*J*(C, H)=160, 159, 158, C arom.); 125.6 (*d*, ¹*J*(C, H)=151, C(7) or C(8)); 86.0, 84.2 (2*d*, ¹*J*(C, H)=144, 142, C(5)); 75.0, 74.9 (2*t*, ¹*J*(C, H)=143, 142, CH₂(Bn)); 49.3, 48.6 (2*d*, ¹*J*(C,

H)=129, 130.2, C(4)); 40.9, 40.8 (2*d*, $^1J(\text{C}, \text{H})=127, 125$, C(6)); 39.5, 37.4 (2*t*, $^1J(\text{C}, \text{H})=125$, C(2)); 18.6, 18.5 (2*q*, $^1J(\text{C}, \text{H})=127$, CH₃-C(6)); 17.9, 17.9 (2*q*, $^1J(\text{C}, \text{H})=125$, CH₃-C(9)); 13.6, 13.6 (2*q*, $^1J(\text{C}, \text{H})=129, 129$, CH₃-C(4)); 7.6, 7.4 (2*q*, $^1J(\text{C}, \text{H})=128, 127.5$, CH₃-C(1));
Anal. Calcd. for C₁₈H₂₆O₂ (274.39): C 78.79, H 9.55; Found: C 78.71, H 9.56.

(4*S*,5*R*,6*E*)-5-Methyl-4-[[*(1S)*-1-phenylethyl]oxy]oct-6-en-2-one (31) and (4*R*,5*S*,6*E*)-5-Methyl-4-[[*(1S)*-1-phenylethyl]oxy]oct-6-en-2-one (32)

Colorless oil (66%), inseparable mixture

31: IR (film): 2936, 1735, 1459, 1411, 1259, 914, 837; **^1H NMR (CDCl₃, 400 MHz):**
 δ 7.32- 7.25 (*m*, 5 arom. H), 5.38 (*dq*, $^3J(\text{H-C}(7), \text{H-C}(6))=15.5$, $^3J(\text{H-C}(7), \text{CH}_3(8))=5.9$, H-C(7)), 5.34 (*ddq*, $^3J(\text{H-C}(6), \text{H-C}(7))=15.5$, $^3J(\text{H-C}(6), \text{H-C}(5))=6.8$, $^4J(\text{H-C}(6), \text{CH}_3(8))=1.5$, H-C(6)), 4.47 (*q*, $^3J(\text{H-C}(1'), \text{CH}_3(2'))=6.5$, H-C(1')), 3.76 (*ddd*, $^3J(\text{H-C}(4), \text{H}_a\text{-C}(3))=7.4$, $^3J(\text{H-C}(4), \text{H}_b\text{-C}(3))=4.6$, $^3J(\text{H-C}(4), \text{HC}(5))=3.8$, H-C(4)), 2.5 (*qdd*, $^3J(\text{H-C}(5), \text{CH}_3\text{-C}(5))=6.9$, $^3J(\text{H-C}(5), \text{H-C}(6))=6.9$, $^3J(\text{H-C}(5), \text{H-C}(4))=3.8$, H-C(5)), 2.44 (*dd*, $^2J(\text{H}_a\text{-C}(3), \text{H}_b\text{-C}(3))=15.4$, $^3J(\text{H}_a\text{-C}(3), \text{H-C}(4))=7.4$, H_a-C(3)), 2.3 (*dd*, $^2J(\text{H}_b\text{-C}(3), \text{H}_a\text{-C}(3))=15.4$, $^3J(\text{H}_b\text{-C}(3), \text{H-C}(4))=4.6$, H_b-C(3)), 1.88 (*s*, CH₃(1)), 1.64 (*dd*, $^3J(\text{CH}_3(8), \text{H-C}(7))=5.9$, $^3J(\text{CH}_3(8), \text{H-C}(6))=1.5$, CH₃(8)), 1.4 (*d*, $^3J(\text{CH}_3(2'), \text{H-C}(1'))=6.5$, CH₃(2')), 1.04 (*d*, $^3J(\text{CH}_3(5), \text{H-C}(5))=6.9$, CH₃-C(5)); **^{13}C NMR (CDCl₃, 100.61MHz):** 207.9 (*s*, C(2)), 143.9 (*s*, C arom.), 132.6 (*d*, $^1J(\text{C}, \text{H})=148$, C(6)), 128.2, 127.5, 126.5 (3*d*, $^1J(\text{C}, \text{H})=160, 160, 157$, 5 arom. C), 125.8 (*d*, $^1J(\text{C}, \text{H})=157$, C(7)), 76.9 (*d*, $^1J(\text{C}, \text{H})=136$, C(4)), 76.2 (*d*, $^1J(\text{C}, \text{H})=141$, C(1')), 45.6 (*t*, $^1J(\text{C}, \text{H})=126$, C(3)), 39.2 (*d*, $^1J(\text{C}, \text{H})=126$, C(5)), 30.7 (*s*, C(1)); 23.9 (*q*, $^1J(\text{C}, \text{H})=127$,

C(2')), 18.1 (*q*, $^1J(\text{C},\text{H})=126$, C(8)), 14.7 (*d*, $^1J(\text{C},\text{H})=126$, $\underline{\text{C}}\text{H}_3\text{-C}(5)$); **HRMS (EI)** calcd for $\text{C}_{17}\text{H}_{24}\text{O}_2\text{Na}^+$ [$\text{M}+\text{Na}^+$] 283.1674, found 283.1679. **Anal.** Calcd for $\text{C}_{17}\text{H}_{24}\text{O}_2$: C, 78.42; H, 9.29. Found: C, 78.32; H, 9.30.

32: IR (film): 2930, 2858, 2363, 1778, 1719, 1472, 1391, 1325, 1258, 1189, 1131, 1079, 1006, 940, 905, 866, 836, 774, 668; **^1H NMR (CDCl_3 , 400MHz):** δ 7.33 (*m*, 5 arom. H), 5.34 (*dq*, $^3J(\text{H-C}(7), \text{H-C}(6))=15.4$, $^3J(\text{H-C}(7), \text{CH}_3(8))=5.8$, H-C(7)), 5.25 (*ddq*, $^3J(\text{H-C}(6), \text{H-C}(7))=15.4$, $^3J(\text{H-C}(6), \text{H-C}(5))=7.0$, $^4J(\text{H-C}(6), \text{CH}_3(8))=1.3$, H-C(6)), 4.48 (*q*, $^3J(\text{H-C}(1'), \text{CH}_3(2'))=6.4$, H-C(1')), 3.83 (*ddd*, $^3J(\text{H-C}(4), \text{Ha-C}(3))=7.7$, $^3J(\text{H-C}(4), \text{Hb-C}(3))=3.8$, $^3J(\text{H-C}(4), \text{HC}(5))=3.8$, H-C(4)), 2.69 (*dd*, $^2J(\text{Ha-C}(3), \text{Hb-C}(3))=16.0$, $^3J(\text{Ha-C}(3), \text{H-C}(4))=7.7$, Ha-C(3)), 2.44 (*dd*, $^2J(\text{Hb-C}(3), \text{Ha-C}(3))=16.0$, $^3J(\text{Hb-C}(3), \text{H-C}(4))=4.5$, Hb-C(3)), 2.2 (*s*, $\text{CH}_3(1)$), 2.16 (*m*, H-C(5)), 1.64 (*d*, $^3J(\text{CH}_3(8), \text{H-C}(7))=5.8$, $\text{CH}_3(8)$), 1.39 (*d*, $^3J(\text{CH}_3(2'), \text{H-C}(1'))=6.4$, $\text{CH}_3(2')$), 0.85 (*d*, $^3J(\text{CH}_3(5), \text{H-C}(5))=7.0$, $\text{CH}_3\text{-C}(5)$); **^{13}C NMR (CDCl_3 , 100.61MHz):** 208.5 (*s*, C(2)), 144.3 (*s*, C arom.), 132.8 (*d*, $^1J(\text{C},\text{H})=152$, C(6)), 128.2, 127.3, 126.5 (*3d*, $^1J(\text{C},\text{H})=160$ 162, 159, 5 arom C), 125.5 (*d*, $^1J(\text{C},\text{H})=152$, C(7)), 77.5 (*d*, $^1J(\text{C},\text{H})=148$, C(4)), 76.9 (*d*, $^1J(\text{C},\text{H})=146$, C(1')), 45.0 (*t*, $^1J(\text{C},\text{H})=126$, C(3)), 39.9 (*d*, $^1J(\text{C},\text{H})=127$, C(5)), 31.5 (*q*, $^1J(\text{C},\text{H})=127$, C(1)); 23.7 (*q*, $^1J(\text{C},\text{H})=127$, C(2')), 18.0 (*q*, $^1J(\text{C},\text{H})=127$, C(8)), 14.7 (*d*, $^1J(\text{C},\text{H})=126$, $\underline{\text{C}}\text{H}_3\text{-C}(5)$); **HRMS (EI)** calcd for $\text{C}_{17}\text{H}_{24}\text{O}_2\text{Na}^+$ [$\text{M}+\text{Na}^+$] 283.1674, found 283.1679. **Anal.** Calcd for $\text{C}_{17}\text{H}_{24}\text{O}_2$: C, 78.42; H, 9.29. Found: C, 78.32; H, 9.30.

(3*S*,4*R*,5*E*)-4-Methyl-1-phenyl-3-([(1*S*)-1-phenylethyl]oxy)hept-5-en-1-one ((-)-34) :

Colorless oil (58%).

$[\alpha]_D^{25} = -57$ ($c=0.5$, CHCl_3); **IR (film)**: 2972, 1685, 1598, 1449, 1359, 1265, 1086, 1026, 973, 760, 691; **^1H NMR (CDCl_3 , 400MHz)**: 7.8-7.1 (m, 10 arom H); 5.43 (m, H-C(5)+H-C(6)), 4.52 (q , $^3J(\text{H-C}(1'), \text{CH}_3(2'))=6.4$, H-C(1')), 3.98 (ddd , $^3J(\text{H-C}(3), \text{H}_a\text{-C}(2))=6.7$, $^3J(\text{H-C}(3), \text{H}_b\text{-C}(2))=5.8$, $^3J(\text{H-C}(3), \text{H-C}(4))=4.5$, H-C(3)), 3.0 (dd , $^2J(\text{H}_a\text{-C}(2), \text{H}_b\text{-C}(2))=16.0$, $^3J(\text{H}_a\text{-C}(2), \text{H-C}(3))=6.7$, $\text{H}_a\text{-C}(2)$), 2.91 (dd , $^2J(\text{H}_b\text{-C}(2), \text{H}_a\text{-C}(2))=16.0$, $^3J(\text{H}_b\text{-C}(2), \text{H-C}(3))=5.8$, $\text{H}_b\text{-C}(2)$), 2.54 (m, H-C(4)), 1.66 (d , $^3J(\text{CH}_3(7), \text{H-C}(6))=4.5$, $\text{CH}_3(7)$), 1.42 (d , $^3J(\text{CH}_3(2'), \text{H-C}(1'))=6.4$, $\text{CH}_3(2')$), 1.15 (d , $^3J(\text{CH}_3(4), \text{H-C}(4))=7.0$, $\text{CH}_3\text{-C}(4)$); **^{13}C -NMR (CDCl_3 , 100.6MHz)**: 199.3 (s, C(1)), 143.9 (s, C arom), 137.3 (s, C arom), 132.7 (d, $^1J(\text{C}, \text{H})=160$, C arom), 132.5 (d, $^1J(\text{C}, \text{H})=161$, C(5)), 128.3, 128.2, 128.1, 127.2, 126.3, 126.1 (9 arom C + C(6)), 77.4 (d, $^1J(\text{C}, \text{H})=142$, C(1')), 76.5 (d, $^1J(\text{C}, \text{H})=142$, C(3)), 40.9 (d, $^1J(\text{C}, \text{H})=123$, C(4)), 39.7 (t, $^1J(\text{C}, \text{H})=126$, C(2)), 23.9 (q , $^1J(\text{C}, \text{H})=126$, C(7)), 18.0 (q , $^1J(\text{C}, \text{H})=126$, $\text{CH}_3(4)$), 15.2 (q , $^1J(\text{C}, \text{H})=126$, C(2')); **MS (MALDI)**: 361.57 $[\text{M}+\text{K}]^+$; **Anal.** Calcd for $\text{C}_{22}\text{H}_{26}\text{O}_2$: C, 81.95; H, 8.13. Found: C, 81.84; H, 8.01.

(3R,4S,5E)-4-Methyl-1-phenyl-3-([(1S)-1-phenylethyl]oxy)hept-5-en-1-one ((-)-35) :

Colorless oil (12 %).

$[\alpha]_D^{25} = -12$ ($c=0.5$, CHCl_3); **IR (film)**: **^1H NMR (CDCl_3 , 400MHz)**: 8.1-7.0 (m, 10 arom H); 5.3 (m, H-C(5)+H-C(6)), 4.47 (q , $^3J(\text{H-C}(1'), \text{CH}_3(2'))=6.5$, H-C(1')), 4.0 (ddd , $^3J(\text{H-C}(4), \text{H}_a\text{-C}(3))=7.4$, $^3J(\text{H-C}(4), \text{H}_b\text{-C}(3))=4.3$, $^3J(\text{H-C}(4), \text{H-C}(5))=4.3$, H-C(3)), 3.24 (dd , $^2J(\text{H}_a\text{-C}(2), \text{H}_b\text{-C}(2))=16.0$, $^3J(\text{H}_a\text{-C}(2), \text{H-C}(3))=7.4$, $\text{H}_a\text{-C}(2)$), 2.95 (dd , $^2J(\text{H}_b\text{-C}(2), \text{H}_a\text{-C}(2))=16.0$,

$^3J(\text{H}_b\text{-C}(2), \text{H-C}(3))=4.3$, $\text{H}_b\text{-C}(2))$, 2.23 (*m*, $\text{H-C}(4))$, 1.6 (*d*, $^3J(\text{CH}_3(7), \text{H-C}(6))=4.9$, $\text{CH}_3(7))$, 1.28 (*d*, $^3J(\text{CH}_3(2'), \text{H-C}(1'))=6.5$, $\text{CH}_3(2')$), 0.85 (*d*, $^3J(\text{CH}_3(4), \text{H-C}(4))=6.8$, $\text{CH}_3\text{-C}(4))$;
 ^{13}C NMR (CDCl_3 , 100.6MHz): 199.9 (*s*, $\text{C}(1))$, 144.2 (*s*, C arom), 137.6 (*s*, C arom), 132.9 (*d*, $^1J(\text{C}, \text{H})=161$, C arom), 132.8 (*d*, $^1J(\text{C}, \text{H})=160$, $\text{C}(5))$, 128.5, 128.2, 128.1, 127.3, 126.6, (5 *arom C*), 125.7 (*d*, $^1J(\text{C}, \text{H})=151$, $\text{C}(6))$, 77.8 (*d*, $^1J(\text{C}, \text{H})=151$, $\text{C}(1')$), 77.1 (*d*, $^1J(\text{C}, \text{H})=141$, $\text{C}(3))$, 40.5 (*d*, $^1J(\text{C}, \text{H})=128$, $\text{C}(4))$, 40.5 (*t*, $^1J(\text{C}, \text{H})=125$, $\text{C}(2))$, 23.6 (*q*, $^1J(\text{C}, \text{H})=128$, $\text{C}(7))$, 18.0 (*q*, $^1J(\text{C}, \text{H})=125$, $\text{CH}_3\text{-C}(4))$, 15.2 (*q*, $^1J(\text{C}, \text{H})=128$, $\text{C}(2')$); **MS (MALDI)**: 361.57 $[\text{M}+\text{K}]^+$; **Anal.** Calcd for $\text{C}_{22}\text{H}_{26}\text{O}_2$: C, 81.95; H, 8.13. Found: C, 81.84; H, 8.01.

(5*S*,6*R*,7*E*)-2,2,6-Trimethyl-5-[[*(1S)*-1-phenylethyl]oxy]non-7-en-3-one ((-)-37):

Colorless oil (71%).

$[\alpha]_D^{25} = -48$ ($c=1.00$, CHCl_3); **IR (film)** : 2970, 1704, 1453, 1367, 1088, 973, 700; **^1H NMR (CDCl_3 , 400 MHz)** : δ 7.34-7.23 (*m*, 5 *arom. H*), 5.41-5.37 (*m*, $\text{H-C}(7)+\text{H-C}(8))$, 4.47 (*q*, $^3J(\text{H-C}(1'), \text{CH}_3(2'))=6.2$, $\text{H-C}(1')$), 3.88 (*ddd*, $^3J(\text{H-C}(5), \text{H}_a\text{-C}(4))=6.2$, $^3J(\text{H-C}(5), \text{H}_b\text{-C}(4))=6.2$, $^3J(\text{H-C}(5), \text{HC-(6)})=3.7$, $\text{H-C}(5))$, 2.52 (*dd*, $^3J(\text{H}_a\text{-C}(4), \text{H}_b\text{-C}(4))=17.3$, $^3J(\text{H}_a\text{-C}(4), \text{H-C}(5))=6.2$, $\text{H}_a\text{-C}(4))$, 2.42 (*dd*, $^3J(\text{H}_b\text{-C}(4), \text{H}_a\text{-C}(4))=17.3$, $^3J(\text{H}_b\text{-C}(4), \text{H-C}(5))=6.2$, $\text{H}_b\text{-C}(4))$, 2.42 (*m*, $\text{H-C}(6))$, 1.66 (*d*, $^3J(\text{CH}_3(9), \text{H-C}(8))=4.3$, $\text{CH}_3(9))$, 1.41(*d*, $^3J(\text{CH}_3(2'), \text{H-C}(1'))=6.2$, $\text{CH}_3(2')$), 1.09 (*d*, $^3J(\text{CH}_3(6), \text{H-C}(6))=6.8$, $\text{CH}_3\text{-C}(6))$, 1.0 (*s*, $\text{CH}_3(1)+(\text{CH}_3)_2\text{C}(2))$; **^{13}C NMR (CDCl_3 , 100.61MHz)**: 214.3 (*s*, $\text{C}(3))$, 144.21 (*s*, C arom.); 132.6 (*d*, $^1J(\text{C,H})=151$, $\text{C}(7))$, 128.3, 127.2, 126.3 (3*d*, $^1J(\text{C,H})=161$, 161, 157, 5 *arom. C*),

125.7 (*d*, $^1J(\text{C,H})=151$, C(8)), 77.0 (*d*, $^1J(\text{C,H})=141$, C(5)), 76.7 (*d*, $^1J(\text{C,H})=138$, C(1')), 44.1 (*s*, C(2)); 39.9 (*d*, $^1J(\text{C,H})=125$, C(6)), 39.0 (*t*, $^1J(\text{C,H})=125$, C(4)), 25.9 (*q*, $^1J(\text{C,H})=132$, C(1)), 23.7 (*q*, $^1J(\text{C,H})=125$, C(2')), 18.0 (*q*, $^1J(\text{C,H})=125$, C(9)), 15.9 (*q*, $^1J(\text{C,H})=125$, $\underline{\text{CH}}_3$ -C(6)); **HRMS (EI)** calcd for $\text{C}_{20}\text{H}_{10}\text{O}_2$ [M^+] 302.2253, found 302.2245. **Anal.** Calcd for $\text{C}_{20}\text{H}_{10}\text{O}_2$: C, 79.42; H, 10.00. Found: C, 79.45; H, 10.07.

(4*R*,5*S*,6*S*,7*E*)-5-{[(1*R*)-1-(2,4,6-triisopropylphenyl)ethyl]oxy}-4,6-dimethylnon-7-en-3-one ((+)-39):²

Colorless oil (70%).

$[\alpha]_D^{25} = +2$; $[\alpha]_{577}^{25} = +3$; $[\alpha]_{546}^{25} = +2$; $[\alpha]_{435}^{25} = +5$; $[\alpha]_{405}^{25} = +5$ ($c=1.0$, CHCl_3); **IR (film)** : 2963, 1713, 1608, 1460, 1381, 1210, 1072, 1023, 974, 910, 878, 734; **$^1\text{H NMR}$ (CDCl_3 , 400 MHz)** : δ 7.09-6.94 (*m*, 2 arom H.), 5.46 (*ddq*, $^3J(\text{H-C}(7), \text{H-C}(8)) = 15.6$, $^3J(\text{H-C}(7), \text{H-C}(6)) = 8.3$, $^4J(\text{H-C}(7), \text{CH}_3(9)) = 1.3$, H-C(7)), 5.36 (*dq*, $^3J(\text{H-C}(8), \text{H-C}(7)) = 15.6$, $^3J(\text{H-C}(8), \text{CH}_3(9)) = 6.1$, H-C(8)), 5.06 (*q*, $^3J(\text{H-C}(1'), \text{CH}_3(2')) = 6.7$, H-C(1')), 3.96, 3.18 (2 *sept*, $^3J((\text{CH}_3)_2\text{CH-Ar}, (\text{CH}_3)_2\text{CH-Ar}) = 6.9$, 2($\text{CH}_3)_2\text{CH-Ar}$), 3.67 (*dd*, $^3J(\text{H-C}(5), \text{H-C}(4)) = 7.6$, $^3J(\text{H-C}(5), \text{H-C}(6)) = 3.0$, H-C(5)), 2.88 (*sept*, $^3J((\text{CH}_3)_2\text{CH-Ar}, (\text{CH}_3)_2\text{CH-Ar}) = 6.9$, $(\text{CH}_3)_2\text{CH-Ar}$), 2.72 (*dq*, $^3J(\text{H-C}(4), \text{H-C}(5)) = 7.6$, $^3J(\text{H-C}(4), \text{CH}_3\text{-C}(4)) = 7.1$, H-C(4)), 2.39 (*m*, $\text{CH}_2(2)$); 2.23 (*dqd*, $^3J(\text{H-C}(6), \text{H-C}(7)) = 8.3$, $^3J(\text{H-C}(6), \text{CH}_3\text{-C}(6)) = 6.9$, $^3J(\text{H-C}(6), \text{H-C}(5)) = 3.0$, H-C(6)), 1.68 (*dd*, $^3J(\text{CH}_3(9), \text{H-C}(8)) = 6.1$, $^3J(\text{CH}_3(9), \text{H-C}(7)) = 1.3$, $\text{CH}_3(9)$), 1.53 (*d*, $^3J(\text{CH}_3(2'), \text{H-C}(1')) = 6.7$, $\text{CH}_3(2')$), 1.34-1.2 (*m*, 2($\text{CH}_3)_2\text{CH-Ar}$), 1.22 (*d*, $^3J((\text{CH}_3)_2\text{CH-Ar}, (\text{CH}_3)_2\text{CH-Ar}) = 6.9$, $(\text{CH}_3)_2\text{CH-Ar}$), 1.16 (*d*, $^3J(\text{CH}_3\text{-C}(6), \text{H-C}(6)) = 6.9$, $\text{CH}_3\text{-C}(6)$), 0.97 (*d*, $^3J(\text{CH}_3\text{-C}(6), \text{H-C}(6)) = 6.9$, $\text{CH}_3\text{-C}(6)$).

C(4), H-C(4))=7.1, CH₃-C(4)), 0.97 (*t*, ³*J*(H-C(1), H-C(2))=7.2, CH₃ (1)); ¹³C NMR (CDCl₃, 100.61MHz): 215 (*s*, C(3)), 148.3, 147.0, 144.2, 135 (4*s*, 4 arom. C), 133.3 (*d*, ¹*J*(C,H)=151, C(7)), 125.3 (*d*, ¹*J*(C,H)=151, C(8)), 123.0, 120.3 (2*d*, ¹*J*(C,H)=151, 152, 2 arom. C), 83.7 (*d*, ¹*J*(C,H)=130, C(5)), 74.8 (*d*, ¹*J*(C,H)=138, C(1')), 49.8 (*d*, ¹*J*(C,H)=128, C(4)), 41.2 (*d*, ¹*J*(C,H)=128, C(6)), 35.8 (*t*, ¹*J*(C,H)=124, C(2)), 33.9, 29.1, 28.9 (3*d*, ¹*J*(C,H)=130, 127, 132, 3(CH₃)₂CH-Ar), 24.8, 24.0, 23.9, 23.3, 23.2, 22.3 (3(CH₃)₂CH-Ar+C(2')), 19.1(*q*, ¹*J*(C,H)=126, CH₃-C(6)), 18.0 (*q*, ¹*J*(C,H)=125, C(9)), 14.6 (*q*, ¹*J*(C,H)=129, CH₃-C(4)), 7.5 (*q*, ¹*J*(C,H)=128, C(1)); **CI-MS** (NH₃): 432 (18, [M+18]⁺), 359 (11), 284 (1), 231 (100), 185 (9), 147 (6), 91 (3); **HRMS** (EI) calcd for C₂₈H₄₆O₂ [M⁺] 414.3477, found 414.3497. **Anal.** Calcd for C₂₈H₄₆O₂: C, 81.10; H, 11.18. Found: C, 81.00; H, 11.10.

(4*S*,5*S*,6*S*,7*E*)-5-[(1*R*)-1-(2,4,6-triisopropylphenylethyl)oxy]-4,6-dimethylnon-7-en-3-one
((+)-40):

Colorless oil (15%).

[α]_D²⁵ = +41; [α]₅₇₇²⁵ = +42; [α]₅₄₆²⁵ = +50; [α]₄₃₅²⁵ = +94; [α]₄₀₅²⁵ = +120 (c=1.0, CHCl₃); **IR** (film) : 2962, 1715, 1608, 1461, 1371, 1067, 973, 758; ¹H NMR (CDCl₃, 400 MHz) : δ 7.03-6.91 (*m*, 2H arom.), 5.52 (*ddq*, ³*J*(H-C(7), H-C(8))=15.6, ³*J*(H-C(7), H-C(6))=8.0, ⁴*J*(H-C(7), CH₃(9))=1.3, H-C(7)), 5.46 (*dq*, ³*J*(H-C(8), H-C(7))=15.6, ³*J*(H-C(8), CH₃(9))=5.4, H-C(8)), 4.9 (*q*, ³*J*(H-C(1'), CH₃ (2'))=6.7, H-C(1')), 3.91, 3.13 (2 *sept*, ³*J*((CH₃)₂CH-Ar, (CH₃)₂CH-Ar)=6.8, 6.7, 2(CH₃)₂CH-Ar), 3.6 (*dd*, ³*J*(H-C(5), H-C(4))=8.1, ³*J*(H-C(5), H-C(6))=3.2, H-C(5)), 2.83 (*sept*, ³*J*((CH₃)₂CH-Ar, (CH₃)₂CH-Ar)=6.9, (CH₃)₂CH-Ar), 2.7 (*dq*, ³*J*(H-C(4), H-C(5))=8.1, ³*J*(H-C(4), CH₃-C(4))=7.0, H-C(4)), 2.38 (*m*, H-C(6)), 2.15 (*dq*, ²*J*(H_a-C(2), H_b-

C(2)=18.3, $^3J(\text{H}_a\text{-C}(2), \text{CH}_3(1))=7.0$, $\text{H}_a\text{-C}(2))$, 1.85 (*dq*, $^2J(\text{H}_b\text{-C}(2), \text{H}_a\text{-C}(2))=18.3$, $^3J(\text{H}_b\text{-C}(2), \text{CH}_3(1))=7.0$, $\text{H}_b\text{-C}(2))$, 1.7 (*dd*, $^3J(\text{CH}_3(9), \text{H-C}(8))=5.4$, $^3J(\text{CH}_3(9), \text{H-C}(7))=1.3$, $\text{CH}_3(9))$, 1.48 (*d*, $^3J(\text{CH}_3(2'), \text{H-C}(1'))=6.7$, $\text{CH}_3(2'))$, 1.28 -1.22 (*m*, $3(\text{CH}_3)_2\text{CH-Ar}$), 1.12 (*d*, $^3J(\text{CH}_3\text{-C}(6), \text{H-C}(6))=6.9$, $\text{CH}_3\text{-C}(6))$, 0.9 (*d*, $^3J(\text{CH}_3\text{-C}(4), \text{H-C}(4))=7.0$, $\text{CH}_3\text{-C}(4))$, 0.55 (*t*, $^3J(\text{CH}_3(1), \text{CH}_2(2))=7.0$, $\text{CH}_3(1))$; ^{13}C NMR (CDCl_3 , 100.61MHz): 214.8, (s, C(3)), 148.0, 146.8, 144.5, 135.7 (4s, 4 arom. C), 133.3 (*d*, $^1J(\text{C,H})=153$, C(7)), 124.9 (*d*, $^1J(\text{C,H})=151$, C(8)), 122.7, 120.4 (2*d*, $^1J(\text{C,H})=152$, 158, 2 arom. C), 85.7 (*d*, $^1J(\text{C,H})=138$, C(5)), 75.6 (*d*, $^1J(\text{C,H})=140$, C(1')), 49.5 (*d*, $^1J(\text{C,H})=125$, C(4)), 39.6 (*d*, $^1J(\text{C,H})=127$, C(6)), 36.6 (*t*, $^1J(\text{C,H})=124$, C(2)), 34.0, 29.1, 28.9 (3*d*, $^1J(\text{C,H})=129$, 127, 132, $(\text{CH}_3)_2\text{CH-Ar}$), 25.6, 25.1, 25.0, 24.1, 23.9, 23.3, 22.9 ($3(\text{CH}_3)_2\text{CH-Ar}+\text{C}(2')$), 18.9 (*q*, $^1J(\text{C,H})=128$, $\text{CH}_3\text{-C}(6))$, 18.1 (*q*, $^1J(\text{C,H})=126$, C(9)), 13.4 (*q*, $^1J(\text{C,H})=127$, $\text{CH}_3\text{-C}(4))$, 7.0 (*q*, $^1J(\text{C,H})=132$, C(1)); **HRMS (MALDI)** calcd for $\text{C}_{28}\text{H}_{46}\text{O}_2\text{Na}^+ [\text{M}+\text{Na}^+]$, 437.6534 found 437.6538; **Anal.** Calcd for $\text{C}_{28}\text{H}_{46}\text{O}_2$: C, 81.10; H, 11.18. Found: C, 81.00; H, 11.10.

(3*R*,4*S*,5*E*)-4-Methyl-1-phenyl-3-[[*(1R)*-1-(2,4,6-triisopropylphenylethyl)oxy]hep-5-en-1-one ((+)-41):

Colorless oil (58%).

$[\alpha]_D^{25} = +29$ ($c=1.18$, CHCl_3); **IR (film)**: 2960, 2920, 2870, 1685, 1610, 1450, 1370, 1325, 1210, 1100, 1170, 1025, 995, 975, 880, 750, 690; **UV (MeCN)**: 210 (13000), 195 (9400); ^1H NMR (CDCl_3 , 400 MHz) : δ 7.77-7.34 (*m*, 5 arom. H), 7.05-6.88 (*m*, 2 arom. H), 5.45 (*ddq*, $^3J(\text{H-C}(5), \text{H-C}(6))=15.4$, $^3J(\text{H-C}(5), \text{H-C}(4))=8.5$, $^4J(\text{H-C}(5), \text{CH}_3(7))=1.6$, $\text{H-C}(5))$,

5.31 (*dq*, $^3J(\text{H-C}(6), \text{H-C}(5))=15.4$, $^3J(\text{H-C}(6), \text{CH}_3(7))=6.3$, $\text{H-C}(6)$), 5.05 (*q*, $^3J(\text{H-C}(1'), \text{CH}_3(2'))=6.7$, $\text{H-C}(1')$), 4.15 (*ddd*, $^3J(\text{H-C}(3), \text{H}_a\text{-C}(2))=7.5$, $^3J(\text{H-C}(3), \text{H}_b\text{-C}(2))=4.4$, $^3J(\text{H-C}(3), \text{H-C}(4))=3.0$, $\text{H-C}(3)$), 3.97, 3.15 (*2m*, $2(\text{CH}_3)_2\text{CH-Ar}$), 3.03 (*dd*, $^2J(\text{H}_a\text{-C}(2), \text{H}_b\text{-C}(2))=17.0$, $^3J(\text{H}_a\text{-C}(2), \text{H-C}(3))=7.5$, $\text{H}_a\text{-C}(2)$), 2.93 (*dd*, $^2J(\text{H}_b\text{-C}(2), \text{H}_a\text{-C}(2))=17.0$, $^3J(\text{H}_b\text{-C}(2), \text{H-C}(3))=4.4$, $\text{H}_b\text{-C}(2)$), 2.87 (*sept*, $^3J((\text{CH}_3)_2\text{CH-Ar}, (\text{CH}_3)_2\text{CH-Ar})=6.9$, $(\text{CH}_3)_2\text{CH-Ar}$), 2.36 (*m*, $\text{H-C}(4)$), 1.60 (*dd*, $^3J(\text{CH}_3(7), \text{H-C}(6))=6.3$, $^3J(\text{CH}_3(7), \text{H-C}(5))=1.6$, $\text{CH}_3(7)$), 1.56 (*d*, $^3J(\text{CH}_3(2'), \text{H-C}(1'))=6.7$, $\text{CH}_3(2')$), 1.27-1.06 (*m*, $3(\text{CH}_3)_2\text{CH-Ar}$), 1.18 (*d*, $^3J(\text{CH}_3\text{-C}(4), \text{H-C}(4))=6.8$, $\text{CH}_3\text{-C}(4)$); ^{13}C NMR (CDCl_3 , **100.61MHz**): 199.5 (*s*, $\text{C}(1)$), 148.5, 147.0, 146.8, 137.2, 135.4 (*5s*, 5 arom. C), 132.9 (*d*, $^1J(\text{C,H})=161$, C arom.), 132.8 (*d*, $^1J(\text{C,H})=161$, $\text{C}(5)$), 128.4, 128.0 (*2d*, $^1J(\text{C,H})=162$, 161, 2 arom. C), 125.9 (*d*, $^1J(\text{C,H})=151$, $\text{C}(6)$), 123.1, 120.4 (*2d*, $^1J(\text{C,H})=154$, 2 arom. C), 78.8 (*d*, $^1J(\text{C,H})=143$, $\text{C}(3)$), 72.7 (*d*, $^1J(\text{C,H})=141$, $\text{C}(1')$), 41.6 (*t*, $^1J(\text{C,H})=122$, $\text{C}(2)$), 41.5 (*d*, $^1J(\text{C,H})=127$, $\text{C}(4)$), 34.0, 29.1, 29.0 (*3d*, $^1J(\text{C,H})=127$, 127, 131, $3(\text{CH}_3)_2\text{CH-Ar}$), 25.2, 25.0, 24.6, 24.0, 23.9 ($3(\text{CH}_3)_2\text{CH-Ar}$), 22.0 (*q*, $^1J(\text{C,H})=126$, $\text{C}(2')$), 18.0 (*q*, $^1J(\text{C,H})=126$, $\text{C}(7)$), 17.6 (*q*, $^1J(\text{C,H})=128$, $\text{CH}_3\text{-C}(4)$); **CI-MS** (NH_3): 466 (0.4, $[\text{M}+18]^+$), 351 (1), 231 (100), 147 (3), 105 (25), 77 (7); **Anal.** Calcd for $\text{C}_{31}\text{H}_{44}\text{O}_2$: C, 82.98; H, 9.88. Found: C, 82.95; H, 9.81.

(3*S*,4*R*,5*E*)-4-Methyl-1-phenyl-3-{[(1*R*)-1-(2,4,6-triisopropylphenylethyl)oxy]hep-5-en-1-one ((+)-42):

Colorless oil (19%).

$[\alpha]_D^{25} = +10.7$ ($c=1.15$, CHCl_3); **IR (film)**: 2961, 2869, 1669, 1615, 1448, 1382, 1287, 1210, 1070, 1210, 1070, 1051, 970, 878, 733, 697; **$^1\text{H NMR}$ (CDCl_3 , 400 MHz)**: δ 8.0-7.48 (*m*, 5 arom. H), 6.99-6.92 (*m*, 2 arom. H), 5.38 (*m*, H-C(5)+H-C(6)), 5.1 (*q*, $^3J(\text{H-C}(1'), \text{CH}_3(2'))=6.4$, H-C(1')), 4.18 (*ddd*, $^3J(\text{H-C}(3), \text{H}_a\text{-C}(2))=8.9$, $^3J(\text{H-C}(3), \text{H}_b\text{-C}(2))=3.8$, $^3J(\text{H-C}(3), \text{H-C}(4))=3.2$, H-C(3)), 3.7, 3.18 (*2m*, $2(\text{CH}_3)_2\text{CH-Ar}$), 3.28 (*dd*, $^2J(\text{H}_a\text{-C}(2), \text{H}_b\text{-C}(2))=15.4$, $^3J(\text{H}_a\text{-C}(2), \text{H-C}(3))=8.9$, $\text{H}_a\text{-C}(2)$), 2.85 (*sept*, $^3J((\text{CH}_3)_2\text{CH-Ar}, (\text{CH}_3)_2\text{CH-Ar})=6.9$, $(\text{CH}_3)_2\text{CH-Ar}$), 2.83 (*dd*, $^2J(\text{H}_b\text{-C}(2), \text{H}_a\text{-C}(2))=15.4$, $^3J(\text{H}_b\text{-C}(2), \text{H-C}(3))=3.8$, $\text{H}_b\text{-C}(2)$), 2.45 (*m*, H-C(4)), 1.64 (*dd*, $^4J(\text{CH}_3(7), \text{H-C}(6))=4.5$, $^3J(\text{CH}_3(7), \text{H-C}(5))=1.3$, $\text{CH}_3(7)$), 1.36 (*d*, $^3J(\text{CH}_3(2'), \text{H-C}(1'))=6.4$, $\text{CH}_3(2')$), 1.28-1.09 (*m*, $3(\text{CH}_3)_2\text{CH-Ar}$), 0.94 (*d*, $^3J(\text{CH}_3\text{-C}(4), \text{H-C}(4))=7.0$, $\text{CH}_3\text{-C}(4)$); **$^{13}\text{C NMR}$ (CDCl_3 , 100.61 MHz)**: 200.4, (*s*, C(1)), 148.6, 147.0, 144.4, 137.9, 135.0 (*5s*, 5 arom. C), 133.1 (*d*, $^1J(\text{C,H})=161$, C arom.), 132.8 (*d*, $^1J(\text{C,H})=161$, C(5)), 128.5, 128.3 (*2d*, $^1J(\text{C,H})=161$, 2 arom. C), 125.6 (*d*, $^1J(\text{C,H})=151$, C(6)), 122.9, 120.2 (*2d*, $^1J(\text{C,H})=151$, 2 arom. C), 79.2 (*d*, $^1J(\text{C,H})=142$, C(3)), 72.9 (*d*, $^1J(\text{C,H})=139$, C(1')), 39.96 (*t*, $^1J(\text{C,H})=123$, C(2)), 39.6 (*d*, $^1J(\text{C,H})=125$, C(4)), 33.9, 29.1, 28.7 (*3d*, $^1J(\text{C,H})=126$, 126, 129, $3(\text{CH}_3)_2\text{CH-Ar}$), 24.9, 24.8, 24.6, 24.3, 23.9 ($3(\text{CH}_3)_2\text{CH-Ar}$), 22.1 (*q*, $^1J(\text{C,H})=126$, C(2')), 18.1 (*q*, $^1J(\text{C,H})=126.3$, C(7)), 13.8 (*q*, $^1J(\text{C,H})=126$, $\text{CH}_3\text{-C}(4)$); **CI-MS (NH_3)**: 466 (0.4, $[\text{M}+18]^+$), 351 (1), 231 (100), 147 (3), 105 (25), 77 (7); **Anal.** Calcd for $\text{C}_{31}\text{H}_{44}\text{O}_2$: C, 82.98; H, 9.88. Found: C, 82.95; H, 9.81.

(2*R*)-2-[(1*R*, 2*S*, 3*E*)-2-Methyl-1-[(*R*)-1-(2,4,6-triisopropylphenylethyl)oxy]pent-3-en-cyclohexanone 44 and (-)-(2*S*)-2-[(1*S*, 2*R*, 3*E*)-2-Methyl-1-[(*R*)-1-(2,4,6-triisopropylphenylethyl)oxy]pent-3-en-cyclohexanone 45

Colorless oil (66%), inseparable mixture.

IR (film) : 3055, 2965, 2935, 2870, 1705, 1605, 1510, 1460, 1420, 1385, 1265, 1170, 1105, 1070, 1030, 895, 750, 705; **UV**: 251 (4700), 203 (18900); **¹H NMR (CDCl₃, 400 MHz)** : δ 7.04-6.94 (*m*, 4H arom.), 5.44 (*ddq*, $^3J(\text{H-C}(3'), \text{H-C}(4')) = 15.5$, $^3J(\text{H-C}(3'), \text{H-C}(2')) = 8.8$, $^4J(\text{H-C}(3'), \text{CH}_3(5')) = 1.5$, 2H-C(3')), 5.36 (*dq*, $^3J(\text{H-C}(4'), \text{H-C}(3')) = 15.5$, $^3J(\text{H-C}(4'), \text{CH}_3(5')) = 5.9$, 2H-C(4')), 4.94 (*q*, $^3J(\text{H-C}(1''), \text{CH}_3(2'')) = 7.0$, 2H-C(1'')), 4.04 (*dd*, $^3J(\text{H-C}(1'), \text{H-C}(2')) = 5.3$, $^3J(\text{H-C}(1'), \text{H-C}(2)) = 3.5$, 2H-C(1')), 3.97, 3.08, 2.85 (3 *sept*, $^3J((\text{CH}_3)_2\text{CH-Ar}, (\text{CH}_3)_2\text{CH-Ar}) = 6.9$, 6(CH₃)₂CH-Ar), 2.85 (*m*, 2H-C(2')), 2.45-1.85 (*m*, 18H, H-C(2), CH₂(3), CH₂(4), CH₂(5), CH₂(6)), 1.64 (*dd*, $^3J(\text{CH}_3(5'), \text{H-C}(4')) = 5.9$, $^4J(\text{CH}_3(5'), \text{H-C}(3')) = 1.5$, 2CH₃(5')), 1.48 (*d*, $^3J(\text{CH}_3(2''), \text{H-C}(1'')) = 7.0$, 2CH₃(2'')), 1.27-1.21 (*m*, 6(CH₃)₂CH-Ar), 1.13, (*d*, $^3J(\text{CH}_3\text{-C}(2'), \text{H-C}(2')) = 7.0$, 2CH₃-C(2')) ; **¹³C NMR (CDCl₃, 100.61MHz)**: 213.4 (*s*, C(1)), 212.5 (*s*, C(1)), 148.4, 148.9, 147.3, 146.6, 144.7, 144.2, 135.5, 135.1 (8*s*, 8 arom. C), 133.9, 134.4 (2*d*, $^1J(\text{C,H}) = 153$, 152, 2C(3')), 124.8 (2*d*, $^1J(\text{C,H}) = 153$, 2C(4')), 123.0, 120.4 (4*d*, $^1J(\text{C,H}) = 155$, 150, 4 arom. C), 80.13, 77.6 (2*d*, $^1J(\text{C,H}) = 142$, 141, 2C(1')), 73.3, 73.0 (2*d*, $^1J(\text{C,H}) = 141$, 139, 2C(1'')), 54.8, 54.6 (2*d*, $^1J(\text{C,H}) = 125$, 125, 2C(2)), 42.6 (2*t*, $^1J(\text{C,H}) = 126$, 2C(6)), 42.1 (2*d*, $^1J(\text{C,H}) = 128$, 2C(2')), 33.9, 29.0, 28.9 (6*d*, $^1J(\text{C,H}) = 129$, 129, 125, 6(CH₃)₂CH-Ar), 30.0, 28.0, 22.8 (6*t*, $^1J(\text{C,H}) = 126$, C(3), C(4), C(5)), 25.5, 25.1, 24.7, 24.5, 23.9 (10*q*, $^1J(\text{C,H}) = 126$, 6(CH₃)₂CH-Ar), 21.7 (2*q*, $^1J(\text{C,H}) = 126$, 2C(2'')), 18.8 (2*q*, $^1J(\text{C,H}) = 125$, 2C(5')) ; 18.0 (2*q*, $^1J(\text{C,H}) = 125$, 2CH₃-C(2')) ; **IC-SM (NH₃)**: 425 (0.2, [M⁺-H]), 411 (0.2, [M⁺-CH₃]), 381 (0.4), 340 (0.4), 231 (100), 215 (16), 131 (6), 91 (8); **Anal.** Calcd for C₂₉H₄₆O₂: C, 81.63; H, 10.87. Found: C, 81.70; H, 10.87.

4.1. Determination of enantiomeric excess :

The enantiomeric purities of the pure retro-ene compounds **34**, **37**, **39**, **41**, **44** were established by cleavage of chiral auxiliary and transformation of the corresponding alcohols into the Mosher's esters.

General procedure for the formation of Mosher esters. To a solution of the alcohol in CH_2Cl_2 (0.04 M in concentration) were added (*R*)-(+)- or (*S*)-(-)- α -methoxy- α -(trifluoromethyl)phenylacetic acid (3 eq.), DMAP (0.5 eq.) and DCC (4 eq.). The reaction mixture was stirred at room temperature for 2 h then diluted with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium bicarbonate, water and brine and dried (anh. Na_2SO_4). The solvent was removed under reduced pressure. The residue was purified by column chromatography on silica gel (PE/EtOAc:10/1) affording the corresponding Mosher's esters. The relatively high acidity of the benzyl ether cleavage reaction is believed to be responsible for practical racemization as the measured e.e.'s decreased with the exposure time to CF_3COOH .

(R)- MPTA ester of compound (+)-39:

To a solution of (+)-**39** (0.153 g, 0.37 mmol) in 5 ml of CH₂Cl₂ was added at 0°C 0.15 ml TFA (5 eq., 1.85 mmol). The mixture was stirred at this temperature for 4h and then a saturated aqueous solution of NaHCO₃ (2 ml) was added. The aqueous layer was extracted three times with CH₂Cl₂ (3 ml). The combined organic extracts were dried (anh. Na₂SO₄). After work-up and column chromatography the desired alcohol **46c** was obtained in 60% yield (41 mg). From **46c** (12 mg, 0.065 mmol) and (R)-α-methoxy-α-(trifluoromethyl)phenylacetic acid, **47c**, was obtained in 98 % (25.5 mg) as a mixture of diastereoisomers 21/1, **91 % ee** (based on ¹⁹F NMR studies). The same reaction was repeated with (S)-α-methoxy-α-(trifluoromethyl)phenylacetic acid for a correct attribution of the minor diastereoisomer signal in ¹⁹F NMR.

47c (major isomer): **IR (film)** : 2976, 2942, 1745, 1716, 1452, 1379, 1258, 1171, 1123, 1015, 974, 922, 765, 721, 699; **¹H NMR (CDCl₃, 400 MHz)** : δ 7.57-7.4 (m, 5 arom. H), 5.4 (dd, ³J(H-C(5), H-C(4)) = 6.8, ³J(H-C(5), H-C(6)) = 6.8, H-C(5)), 5.37 (dq, ³J(H-C(8), H-C(7)) = 15.4, ³J(H-C(8), CH₃(9)) = 6.8, H-C(8)), 5.24 (ddq, ³J(H-C(7), H-C(8)) = 15.4, ³J(H-C(7), H-C(6)) = 8.6, ³J(H-C(7), CH₃(9)) = 1.2, H-C(7)), 3.54 (s, OCH₃), 2.86 (dq, ³J(H-C(4), H-C(5)) = 6.8, ³J(H-C(4), CH₃-C(4)) = 6.8, H-C(4)), 2.49 (qd, ³J(H-C(2), H-C(1)) = 7.4, ³J(H_a-C(2), H_b-C(2)) = 3.1, H-C(2)), 2.48 (qd, ³J(H-C(2), H-C(1)) = 7.4, ³J(H_a-C(2), H_b-C(2)) = 3.1, H-C(2)), 2.43 (m, H-C(6)); 1.6 (dd, ³J(CH₃(9), H-C(8)) = 6.78, ⁴J(CH₃(9), H-C(7)) = 1.2, CH₃(9)), 1.07 (d, ³J(CH₃-C(6), H-C(6)) = 7.4, CH₃-C(6)), 1.05 (t, ³J(H-C(1), H-C(2)) = 7.4, CH₃(1)), 0.96 (d, ³J(CH₃-C(4), H-C(4)) = 6.8, CH₃-C(4)) ; **¹³C NMR (CDCl₃, 100.61 MHz)**: 212.73 (C(3)), 165.93 (OCOR), 131.8, 131.2, 129.5, 128.3, 127.7, 127.2, 124.7, 121.9, 118.9 (5 arom.

C+C(7)+C(8)), 79.82 (*d*, $^1J(\text{C,H})=150$, C(5)), 55.26 (*q*, $^1J(\text{C,H})=143$, OCH₃), 47.51 (*d*, $^1J(\text{C,H})=128$, C(4)), 39.62 (*d*, $^1J(\text{C,H})=127$, C(6)), 34.86 (*t*, $^1J(\text{C,H})=127$, C(2)), 17.84 (*q*, $^1J(\text{C,H})=126$, CH₃(9)), 17.52 (*q*, $^1J(\text{C,H})=129$, CH₃-C(6)), 12.16 (*q*, $^1J(\text{C,H})=130$, CH₃-C(4)), 7.53 (*q*, $^1J(\text{C,H})=129$, CH₃(1)); **^{19}F NMR**: δ : -71.56 (major), -71.6 (minor); **HRMS (MALDI)** calcd for C₂₄H₂₇F₃O₄Na⁺ [M+Na⁺], 423.1759, found 423.1709; **Anal.** Calcd for C₂₁H₂₇F₃O₄: C, 62.99; H, 6.80. Found: C, 63.0; H, 6.71.

(R)- MPTA ester of compound (+)-41:

To a solution of (+)-**41** (0.095 g, 0.21 mmol) in 10 ml of CH₂Cl₂ was added at -20°C 0.08 ml TFA (5 eq., 1.06 mmol). The mixture was stirred at this temperature for 3h and then a saturated aqueous solution of NaHCO₃ (5 ml) was added. The aqueous layer was extracted three times with CH₂Cl₂ (5 ml). The combined organic extracts were dried (anh. Na₂SO₄). After work-up and column chromatography the desired alcohol **ent 46a** was obtained in 96% yield (44 mg). From **46a** (15 mg, 0.068 mmol) and (*R*)- α -methoxy- α -(trifluoromethyl)phenylacetic acid, **47a**, was obtained in 95 % (28 mg) as a single compound in 99 % **ee** (based on ^{19}F NMR studies). The same reaction was repeated with ($\pm$)-**41** for a correct attribution of the minor diastereoisomer signal in ^{19}F NMR.

IR (film) : 2971, 1748, 1688, 1598, 1581, 1495, 1450, 1380, 1272, 1181, 1122, 1081, 1019, 1000, 922, 846, 821, 754, 720, 691; **^1H -NMR (CDCl₃, 400MHz)**: 7.9-7.34 (m, 10 arom H); 5.74 (*ddd*, $^3J(\text{H-C}(3), \text{H}_a\text{-C}(2))=7.4$, $^3J(\text{H-C}(3), \text{H}_b\text{-C}(2))=4.9$, $^3J(\text{H-C}(3), \text{H-C}(4))=6.8$, H-C(3)), 5.5(*dq*, $^3J(\text{H-C}(6), \text{H-C}(5))=15.4$, $^3J(\text{H-C}(6), \text{CH}_3(7))=6.2$, H-C(6)), 5.37 (*ddq*, $^3J(\text{H-C}(5), \text{H-C}(6))=15.4$, $^3J(\text{H-C}(5), \text{H-C}(4))=8.0$, $^3J(\text{H-C}(5), \text{CH}_3(7))=1.2$, H-C(5)), 3.52 (s,

OCH₃), 3.3 (*dd*, ²*J*(H_a-C(2), H_b-C(2))=17.3, ³*J*(H_a-C(2), H-C(3))=7.4, H_a-C(2)), 3.09 (*dd*, ²*J*(H_b-C(2), H_a-C(2))=17.3, ³*J*(H_b-C(2), H-C(3))=4.9, H_b-C(2)), 2.64 (*m*, H-C(4)), 1.66 (*d*, ³*J*(CH₃(7), H-C(6))=6.2, ³*J*(CH₃(7), H-C(5))=1.2, CH₃(7)) ; 1.1 (*d*, ³*J*(CH₃(4), H-C(4))=6.8, CH₃-C(4)) ; ¹³C NMR (CDCl₃, 100.61MHz): 196.7 (*s*, C(1)) , 165.9 (*s*, OCOR), 136.7, 133.4, 132.5, 130.6, 129.4, 128.6, 128.2, 127.8, 124.8, 121.9(10arom.C+C(5)+C(6)), 75.9 (*d*, ¹*J*(C,H)=153, C(3)), 55.4 (*q*, ¹*J*(C,H)=144, OCH₃), 39.6 (*d*, ¹*J*(C,H)=131, C(4)), 37.4 (*t*, ¹*J*(C,H)=127, C(2)), 18. (*q*, ¹*J*(C,H)=131, CH₃-C(7)), 15.4 (*q*, ¹*J*(C,H)=127, CH₃-C(4)); ¹⁹F NMR : δ: -71.56 (major), -72.01 (minor); HRMS (MALDI) calcd for C₂₄H₂₅F₃O₄Na⁺ [M+Na⁺], 457.1603, found 457.1695; Anal. Calcd for C₂₄H₂₅F₃O₄: C, 66.35; H, 5.80, Found: C, 66.31; H, 5.88.

(S)- MPTA ester of compound (-)-34:

Following the general procedure, to a solution of (-)-**34** (0.05 g, 0.15 mmol) in 1 ml of CH₂Cl₂ was added at 0°C 0.11 ml TFA (10 eq., 1.85 mmol) and 0.016 ml of anisole (1 eq., 0.15 mmol). The mixture was stirred at this temperature for 4h and then a saturated aqueous solution of NaHCO₃ (2 ml) was added. The aqueous layer was extracted three times with CH₂Cl₂ (3 ml). The combined organic extracts were dried (anh. Na₂SO₄). After work-up and column chromatography the desired alcohol **46a** was obtained in 83% yield (28 mg). From **46a** (10 mg, 0.045 mmol) and (*S*)-α-methoxy-α-(trifluoromethyl)phenylacetic acid, **47a**, was obtained in 95 % (18.8 mg) as a mixture of diastereoisomers 35/1, **94% ee** (based on ¹⁹F NMR studies).

¹⁹F NMR : δ : -71.56 (major), -72.01 (minor) ;

(R)-MPTA ester of compound (+)-37 :

To a solution of (-)-**37** (40 mg, 0.13 mmol) in 1 ml of CH₂Cl₂ was added at -10°C 0.05 ml TFA (5 eq., 0.66 mmol) and 0.014 ml of anisole (1 eq., 0.13 mmol). After work-up and column chromatography the desired alcohol **46b** was obtained in 69% yield (18 mg). From **46b** (9 mg, 0.046 mmol) and (*R*)- α -methoxy- α -(trifluoromethyl)phenylacetic acid, **47b**, was obtained in 94 % (18.0 mg) as a mixture of diastereoisomers 24/1, **92% ee** (based on ¹⁹F NMR studies). The same reaction was repeated with ($\pm$)-**37** for a correct attribution of minor diastereoisomer signal in ¹⁹F NMR.

IR (film): 2971, 1749, 1710, 1626, 1479, 1700, 1626, 1479, 1452, 1368, 1261, 1171, 1122, 1081, 1019, 993, 918, 838, 765, 719, 698 ; **¹H-NMR (CDCl₃, 400MHz)**: 7.53-7.37 (m, 5 arom H); 5.57 (ddd, ³*J*(H-C(5), H-C(4))=7.7, ³*J* (H-C(5), H-C(4))=4.5, ³*J* (H-C(5), H-C(6))=3.8, H-C(5)), 5.5(ddq, ³*J*(H-C(8), H-C(7))=15.3, ³*J*(H-C(8), CH₃(9))=6.4, H-C(8)), 5.33 (dd, ³*J*(H-C(7), H-C(8))=15.3, ³*J*(H-C(7), H-C(6))=7.0, H-C(7)), 3.5 (s, OCH₃), 2.85 (dd, ²*J*(H_a-C(4), H_b-C(4))=17.9, ³*J* (H_a-C(4), H-C(5))=7.7, H_a-C(4)), 2.54 (dd, ²*J*(H_b-C(4), H_a-C(4))=17.9, ³*J*(H_b-C(4), H-C(5))=4.5, H_b-C(4)), 2.6 (m, H-C(6)), 1.65 (d, ³*J*(CH₃(9), H-C(8))=6.4, CH₃(9)), 1.03 (d, ³*J*(CH₃(6), H-C(6))=7.0, CH₃(6)), 1.02 (s, H-C(1)+2CH₃-C(2)); **¹³C NMR (CDCl₃, 100.61MHz)**: 211.7 (C(3)) , 165.7 (OCOR), 132.4, 131, 129.5, 128.7, 127.7, 124.8, 122, 119 (6 arom.C+C(7)+C(8)), 75.9 (d, ¹*J*(C,H)=154, C(5)), 55.4 (q, ¹*J*(C,H)=144, OCH₃), 44.2 (s, C(2)), 40 (d, ¹*J*(C,H)=131, C(6)), 37.4 (t, ¹*J*(C,H)=127, C(4)),

26.3 (*q*, $^1J(\text{C},\text{H})=127.0$, $\text{CH}_3(1)+2\text{CH}_3(2)$), 18 (*q*, $^1J(\text{C},\text{H})=126$, $\underline{\text{C}}\text{H}_3\text{-C}(9)$), 16.3 (*q*, $^1J(\text{C},\text{H})=126$, $\underline{\text{C}}\text{H}_3\text{-C}(6)$); ^{19}F NMR : δ : -71.81 (major); -71.93 (minor); **HRMS (MALDI)** calcd for $\text{C}_{22}\text{H}_{29}\text{F}_3\text{O}_4\text{Na}^+$ [$\text{M}+\text{Na}^+$], 437.1916, found 457.1967; **Anal.** Calcd for $\text{C}_{22}\text{H}_{29}\text{F}_3\text{O}_4$: C, 63.75; H, 7.05. Found: C, 63.7; H, 7.08.

(*R*)-MPTA ester of compound 44 :

To a solution of (+)-**44** (50 mg, 0.11 mmol) in 1 ml of CH_2Cl_2 was added at -20°C 0.02 ml TFA (2 eq., 0.23 mmol) and 0.012 ml of anisole (1 eq., 0.11 mmol). The mixture was stirred at this temperature for 1h and then a saturated aqueous solution of NaHCO_3 (1 ml) was added. The aqueous layer was extracted three times with CH_2Cl_2 (3 ml). The combined organic extracts were dried (anh. Na_2SO_4). After work-up and column chromatography the desired alcohol **46d** was obtained in 74% yield (17 mg). From **46d** (8.5 mg, 0.043 mmol) and (*R*)- α -methoxy- α -(trifluoromethyl)phenylacetic acid, **47d**, was obtained in 95 % (16.8 mg) as a mixture of diastereoisomers 35/1, **95 % ee** (based on ^{19}F NMR studies). The same reaction was repeated with (*S*)- α -methoxy- α -(trifluoromethyl)phenylacetic acid for a correct assignment of minor diastereoisomers signal in ^{19}F NMR.

IR (film): 2927, 2856, 1747, 1711, 1451, 1260, 1184, 1123, 1018, 975, 916, 800, 765; **^1H -NMR (CDCl_3 , 400MHz):** 7.57-7.39 (m, 5 arom H); 5.6 (*ddd*, $^3J(\text{H-C}(1'), \text{H-C}(2'))=5.8$, $^3J(\text{H-C}(1'), \text{H-C}(3'))=5.8$, $\text{H-C}(1')$), 5.42 (*dq*, $^3J(\text{H-C}(4'), \text{H-C}(3'))=15.7$, $^3J(\text{H-C}(4'), \text{CH}_3(5'))=6.1$, $\text{H-C}(4')$), 5.31 (*ddq*, $^3J(\text{H-C}(3'), \text{H-C}(4'))=15.7$, $^3J(\text{H-C}(3'), \text{H-C}(2'))=8.9$,

$^4J(\text{H-C}(3'), \text{CH}_3(5'))=1.3$, $\text{H-C}(3')$, 3.6 (s, OCH_3), 2.56 (m, $\text{H-C}(2')$), 2.42-1.83 (m, 9H, $\text{H-C}(2)$, $\text{CH}_2(3)$, $\text{CH}_2(4)$, $\text{CH}_2(5)$, $\text{CH}_2(6)$), 1.6 (dd, $^3J(\text{CH}_3(5'), \text{H-C}(4'))=6.1$, $^4J(\text{CH}_3(5'), \text{H-C}(3'))=1.3$, $\text{CH}_3(5')$), 0.98 (d, $^3J(\text{CH}_3(2'), \text{H-C}(2'))=7.0$, $\text{CH}_3(2')$); ^{13}C NMR (CDCl_3 , 100.61MHz): 209.5 (s, C(1)), 174.8 (OCOR), 132.4, 131.98, 129.4, 128.1, 127.6, 127.2, 124.8, 121.9 (6 arom.C+C(3')+C(4')), 77. (d, $^1J(\text{C,H})=153$, C(1')), 55.5 (q, $^1J(\text{C,H})=145$, OCH_3), 51.4 (d, $^1J(\text{C,H})=124$, C(2)), 42.2 (t, $^1J(\text{C,H})=134$, C(6)), 39.6 (d, $^1J(\text{C,H})=126$, C(2')), 27.6, 27.0, 22.5 (3t, $^1J(\text{C,H})=133$, 131, 125, C(3), C(4), C(5)), 18.0 (q, $^1J(\text{C,H})=127$, C(5')); 17.8 (q, $^1J(\text{C,H})=126$, $\text{CH}_3\text{-C}(2')$); ^{19}F NMR : δ : -71.48 (major), -71.72 (minor); MS (MALDI): 423.1709 [M+K⁺]; Anal. Calcd for $\text{C}_{22}\text{H}_{27}\text{F}_3\text{O}_4$: C, 64.07; H, 6.60. Found: C, 64.11; H, 6.66.

4.2. Determination of relative configurations.

The relative configuration β,γ -anti in the retro-ene products was established by the following procedure: deprotection (TFA, CH_2Cl_2 , anisole) of chiral auxiliary from the compounds ($\pm$)-**31**, ($\pm$)-**34**, and ($\pm$)-**41**, reduction of the carbonyl group ($\text{BEt}_3/\text{NaBH}_4$) to the corresponding syn-1,3-diols **48a**, **48d** and ozonolysis ($\text{O}_3/\text{Me}_2\text{S}$). The corresponding so-obtained aldehydes were equilibrated with a mixture of known α - and β -pyranoses **49a**, and **49d**.³ The β,γ relative configurations were established by NOESY.

The relative configuration α,β -syn in the retro-ene products has been already established by V. Narkevitch in our group, which was based on X-ray analysis of the corresponding sulfones derived from TBAF/MeI quenching of silyl sulfinate intermediates.¹ We have synthesized compound **50** for the determination of the relative configuration.

(+)-(2*E*,4*R*,5*R*,6*R*,7*S*)-(5,7-isopropylidenedioxy)-4,6-dimethyl-non-2-ene ((+)-50)

To the solution of compound **(+)-ent-46c** (157 mg, 0.85 mmol, 1 equiv.) in the mixture of THF/MeOH (4/1, 1ml/0.25ml) was added 1M solution of diethylmethoxyborane in THF (1.1 ml, 1.1 mmol, 1.3 equiv.) at -78 °C. The mixture was stirred at -78 °C for 20 min. NaBH₄ (42 mg, 1.1 mmol, 1.3 equiv.) was added in one portion at the same temperature and the resulting mixture was stirred overnight at -80 °C. The reaction was quenched by acetic acid, extracted by ether (20 ml×3). The organic layer was washed with NH₄Cl (sat.) solution (20 ml), brine (30 ml), dried (anh. Na₂SO₄). After solvent evaporation under *vacuo*, the residue was taken up by MeOH (10 ml) and heated under reflux for 20 min. Then methanol was evaporated giving the crude diol **(+)-ent-48e**.

To the solution of the crude diol **(+)-ent-48e** in acetone/dimethoxypropane (1/1, 14 ml) was added pyridinium paratoluenesulfonate (PPTS) (47 mg, 0.2equiv.). The reaction mixture was stirred at 20 °C for 2 h. Solid NaHCO₃ was added, the suspension was filtered and the filtrate was evaporated under *vacuo*. The residue was purified by flash chromatography on silica gel. Yield: 173 mg (90%) of **(+)-ent-48e**. The coupling constant between H-C(5) and H-C(6) is 2.2 Hz, which proves the *syn*-relationship.

Colorless oil (90%).

$[\alpha]_{589}^{25} = +17$; $[\alpha]_{577}^{25} = +26$; $[\alpha]_{546}^{25} = +19$; $[\alpha]_{435}^{25} = +34$; $[\alpha]_{405}^{25} = +45$ (c=1.13, CHCl₃); **IR (film)**: 3353, 2966, 1613, 1453, 1377, 1260, 1200, 1182, 1140, 1009, 983, 893, 808; **¹H-NMR (CDCl₃, 400MHz)**: 5.47 (m, H(3)+H(2)); 3.72 (dt, ³J(H-C(7), H-C(6))=2.2, ³J(H-C(7), H-C(8))=7.4, H-C(7)); 3.47 (dd, ³J(H-C(5), H-C(6))=2.2, ³J(H-C(5), H-C(4))=9.6, H-C(5)); 2.23 (m, H-C(4)); 1.67 (d, ³J(H-C(1), H-C(2))=5.2, H-C(1)); 1.50 (m, H-C(8)+H-C(6)); 1.40 (s, CH₃-C-O); 1.39 (s, CH₃-C-O); 0.91 (t, ³J(H-C(9), H-C(8))=7.4, H-C(9)); 0.89 (d, ³J(CH₃-C(6), H-C(6))=6.8, CH₃-C(6)); 0.83 (d, ³J(CH₃-C(4), H-C(4))=6.8, CH₃-C(4)); **¹³C-NMR (CDCl₃, 100.6MHz)**: 134.7 (d, ¹J(C, H)=150, C(3)); 123.6 (d, ¹J(C, H)=150, C(2)); 98.7 (s, C(CH₃)₂); 77.6 (d, ¹J(C, H)=141, C(5)); 75.1 (d, ¹J(C, H)=142, C(7)); 37.1 (d, ¹J(C, H)=124, C(4)); 32.3 (d, ¹J(C, H)=124, C(6)); 30.1 (q, ¹J(C, H)=127, CH₃-C-O); 25.7 (t, ¹J(C, H)=127, C(8)); 19.6 (q, ¹J(C, H)=125, CH₃-C-O); 18.2 (q, ¹J(C, H)=132, C(1)); 15.4 (q, ¹J(C, H)=124, C(9)); 9.8 (q, ¹J(C, H)=126, CH₃-C(6)); 4.3 (q, ¹J(C, H)=126, CH₃-C(4)); **CI-MS**: 227 (24, [M+1]⁺), 211 (8), 186 (6), 169 (68), 151 (100), 139 (5), 125 (17), 111 (21), 99 (38)

¹³C-NMR (CDCl₃, 100.6MHz):

1. Narkevitch, V.; Megevand, S.; Schenk, K.; Vogel, P. *J. Org. Chem.* **2001**, *66*, 5080;
2. V. Narkevitch, *Nouvelle réaction de formation de liaison carbone-carbone asymétrique: applications synthétique des co-condensations du dioxyde de soufre avec les diènes at les*

alcènes riches en électrons, Travail de doctorat, Université de Lausanne, **2001**, 184. The reaction sequence for retro-aldol product formation is shown in the following scheme:

3. Roulet, J.-M.; Pühr, G.; Vogel, P. *Tetrahedron Lett.* **1997**, 38, 6201