Palladium Catalyzed Decarboxylation and Decarbonylation under Hydrothermal Condition: Decarboxylative Deuteration

Seijiro Matsubara,* Yutaka Yokota, and Koichiro Oshima

Department of Material Chemistry, Graduate School of Engineering, Kyoto University, Kyotodaigaku-Katsura, Nishikyo, Kyoto 615-8501, Japan.

General Information.

Water (distilled) and 10wt% Pd/C were purchased from Wako Co. Ltd.. Deuterium oxide (D, 99.9%) was purchased from Cambridge Isotope Laboratories, Inc.. All solvents for extraction were used as obtained from commercial suppliers. Chromatographic purification of products was accomplished using forced-flow chromatography on Kanto Chemical Co., INC. Silica gel 60 N (spherical, neutral).

^1H , ^2H and ^{13}C NMR spectra were recorded on Varian Gemini-2000 (300, 40, and 75 MHz, respectively) instrument. Data for ^1H , ^2H and ^{13}C NMR are reported as chemical shift (δ ppm).

Decarboxylation and Decarbonylation Procedure (Procedure I, Table 1):

Figure 1. 30mL Teflon[®] lined autoclave

See the following reference for the reaction vessel in Figure 1: Matsubara, S., Yokota, Y.; Oshima, K. *Chem. Lett.* **2004**, 34, 294; Yamasaki, Y.; Enomoto, H.; Yamasaki, N.; Nakahara, M. *Bull. Chem. Soc. Jpn.*, **2000**, 73, 2687. The vessel is designed to release the internal overpressure. It is commercially available from Shikokurika Co., Ltd., Kochi, Japan.). In a teflon vessel, 10wt% Pd on active carbon (100 mg, 0.1 mmol Pd), water (20 ml), and carbonyl

compound (carboxylic acid, aldehyde, lactone, or amide, 3.0 mmol) were added. After the vessel was placed in autoclave and sealed, the whole was placed in oven (250 °C). After the autoclave was heated for the period in Table 1, it was cooled to the room temperature. The obtained mixture was extracted with pentane and ether. Teflon vessel absorbs organic compounds. To extract the product completely, the vessel washed with hexane and water using ultra sonic cleaner for 30 min each. These solvents used for wash were extracted with ether. The combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. The obtained product was purified by a short silica-gel column chromatography. Instead of the vessel in Figure 1, glass sealed tube is also possible. In the sealed tube case, however, the reaction should be taken care concerning about explosion, as the internal pressure is quite high. The supplier of 10wt% Pd on active carbon was not crucial. We also used that from Nacarai Tesque and Aldrich.

Coupling-degradation Reaction Procedure (Procedure II, 4, 6, and 8):

In a teflon vessel, 10wt% Pd on active carbon (100 mg, 0.1 mmol Pd), 0.1 M NaOHaq (20 ml), and aldehyde (3.0 mmol) were added. After the vessel was placed in autoclave and sealed, the whole was placed in oven (250 °C). After the autoclave was heated for 4 h, it was cooled to the room temperature. The obtained mixture was extracted with pentane and ether. Teflon vessel absorbs organic compounds. To extract the product completely, the vessel washed with hexane and water using ultra sonic cleaner. These solvents used for wash were extracted with ether. The combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. The obtained product was purified by a short silica-gel column chromatography.

Decarboxylative Deuteration Procedure (Procedure III, 10, 12, 14, 16, and 18):

Deuteration was performed using D₂O instead of water (Procedure I) and 0.1 M NaOD/D₂O instead of 0.1 M NaOHaq (Procedure II). Deuteration ratio in the product was determined by ¹H nmr using CHBr₃ as an internal standard and ²H nmr using CDCl₂CDCl₂ as an internal standard. In the product **10**, **12**, **14**, **16**, and **18**, both calculated ratios from ¹H nmr and ²H nmr were coincident.

Compound 10:

Pulse Sequence: s2pul
 Solvent: CDCl₃
 Ambient temperature
 GEMINI-300BB "varian"
 yoke 4108
 Relax. delay 1.000 sec
 Pulse 45.0 degrees
 Acq. time 3.001 sec
 Width 4500.0 Hz
 8 repetitions
 OBSERVE H1, 300.0672956 MHz
 DATA PROCESSING
 FT size 32768
 Total time 0 min, 39 sec

Pulse Sequence: s2pul
 Solvent: CHCl₃
 Ambient temperature
 File:
 GEMINI-300BB "varian"

Pulse 21.6 degrees
 Acq. time 4.016 sec
 Width 2000.0 Hz
 54 repetitions
 OBSERVE H2, 46.0456643 MHz
 DATA PROCESSING
 FT size 4096
 Total time 7 min, 44 sec

[TIC]
Data : GC-EI019 Date : 21-Jan-104 19:50
Sample : -
Note : -
Inlet : GC Ion Mode : EI+
Ion Species : Normal Ion [MF-Linear]
TIC Range : m/z 50 to 500 Output RT Range : 3.10 to 6.23 min
453200384

[Mass Spectrum]
RT : 5.24 min Scan# : 631 Temp : 267.6 deg.C
Ion Mode : EI+ Int. : 1599.98
17116378

Compound 12:

γ-bL-4206

Pulse Sequence: s2pul
 Solvent: CDCl₃
 Ambient temperature
 GEMINI-300BB "varian"

Relax. delay 1.000 sec
 Pulse 45.0 degrees
 Acq. time 3.001 sec
 Width 4500.0 Hz
 8 repetitions
 OBSERVE H1, 300.0672956 MHz
 DATA PROCESSING
 FT size 32768
 Total time 0 min, 39 sec

¹H NMR

CD₃-(CD₂)₁₁-CD₃ D_a: 91%D
 D_b: 91%D

KYOTO UNIVERSITY

Pulse Sequence: s2pul
 Solvent:
 Ambient temperature
 File: γ-bL-4206
 GEMINI-300BB "varian"

Pulse 21.6 Degrees
 Acq. time 4.016 sec
 Width 2000.0 Hz
 54 repetitions
 OBSERVE H2, 46.0456643 MHz
 DATA PROCESSING
 FT size 4096
 Total time 7 min, 44 sec

²H NMR

CD₃-(CD₂)₁₁-CD₃ D_a: 91%D
 D_b: 91%D

KYOTO UNIVERSITY

[TIC]

Data : GC-EI098

Date : 15-Mar-104 11:56

Sample: -

Note: -

Inlet : GC

Ion Mode : EI+

Ion Species : Normal Ion [MF-Linear]

TIC Range : m/z 50 to 500

Output RT Range : 5.73 to 6.59 min

847709440

[Mass Spectrum]

RT : 5.93 min

Scan# : 714

Temp : 294.2 deg.C

Ion Mode : EI+

Int. : 1182.11

13033255

[Mass Spectrum]

RT : 5.93 min

Scan# : 714

Temp : 294.2 deg.C

Ion Mode : EI+

Int. : 1182.11

4648224

Mass :

Da: 91% D

Db: 91% B

Fw(0% D) 187

Fw(100% D) 212

KYOTO UNIVERSITY

Compound **14**:

[TIC]

Data : GC-EI095

Date : 13-Mar-104 12:29

Sample: -

Note: -

Inlet : GC

Ion Mode : EI+

Ion Species : Normal Ion [MF-Linear]

TIC Range : m/z 50 to 500

Output RT Range : 0.00 to 5.92 min

236039184

[Mass Spectrum]

RT : 4.84 min

Scan# : 583

Temp : 252.1 deg.C

Ion Mode : EI+

Int. : 571.31

6254952

[Mass Spectrum]

RT : 4.84 min

Scan# : 583

Temp : 252.1 deg.C

Ion Mode : EI+

Int. : 571.31

3832259

Mass:

D_a: >99%D
 D_b: >99%D
 D_c: 44%D
 D_d: 99%D
 D_e: 99%D
 D_{f,g}: 98%D
 D_h: 99%D

0%D; Fw148

100%D; Fw164

Main Product Fw162

KYOTO UNIVERSITY

Compound 16:

Compound 18:

¹³C OBSERVE

exp1 std13c

SAMPLE DEC. & VT
date Mar 12 2004 dfrq 300.163
solvent CDCl3 dn H1
file exp gpwr 42
ACQUISITION dof 0
sfrq 75.488 dm yyy
tn C13 dnm w
at 2.000 dmf 13200
np 75046 PROCESSING
aw 18761.7 lb 1.00
fb 10400 wtfile
bs 16 proc ft
tpwr 55 fn not used
pw 6.5
dl 1.000 werr
tof 0 weap procplot
nt 10000 wbs
ct 10000 wnt
alock n
gain not used
FLAGS
il n
in n
dp y
DISPLAY
sp -1826.7
wp 18761.7
vs 458
sc 0
wc 250
hmm 24.46
is 500.00
rfl 7638.7
rfp 5812.0
th 2
ins 100.000

¹³C NMR

D_a: 93% D
D_b: 95% D
D_c: 95% D
D_d: 95% D

KYOTO UNIVERSITY

220 200 180 160 140 120 100 80 60 40 20 0 ppm

[TIC]

Data : GC-EI094

Date : 13-Mar-104 12:15

Sample: -

Note: -

Inlet : GC

Ion Mode : EI+

Ion Species : Normal Ion [MF-Linear]

TIC Range : m/z 50 to 500

Output RT Range : 0.00 to 10.91 min

743463424

[Mass Spectrum]

RT : 9.53 min

Scan# : 1147

Temp : 439.4 deg.C

Ion Mode : EI+

Int. : 1546.90

17770892

[Mass Spectrum]

RT : 9.54 min

Scan# : 1148

Temp : 440.0 deg.C

Ion Mode : EI+

Int. : 1057.06

4971515

Mass

0%D: Fw224

100%D: Fw244

KYOTO UNIVERSITY